

Article

# The Algebraic Surfaces of the Enneper Family of Maximal Surfaces in Three Dimensional Minkowski Space

Erhan Güler 

Department of Mathematics, Faculty of Sciences, Kutlubey Campus, Bartın University, Bartın 74100, Turkey; eguler@bartin.edu.tr; Tel.: +90-378-5011000 (ext. 2275)

**Abstract:** We consider the Enneper family of real maximal surfaces via Weierstrass data  $(1, \zeta^m)$  for  $\zeta \in \mathbb{C}$ ,  $m \in \mathbb{Z}_{\geq 1}$ . We obtain the irreducible surfaces of the family in the three dimensional Minkowski space  $\mathbb{E}^{2,1}$ . Moreover, we propose that the family has degree  $(2m + 1)^2$  (resp., class  $2m(2m + 1)$ ) in the cartesian coordinates  $x, y, z$  (resp., in the inhomogeneous tangential coordinates  $a, b, c$ ).

**Keywords:** Weierstrass representation; Enneper maximal surface; algebraic surface; degree; class

**MSC:** primary 53A35; secondary 53C42, 65D18

## 1. Introduction

A minimal surface is a surface of vanishing mean curvature in three dimensional Euclidean space  $\mathbb{E}^3$ . There are many classical and modern minimal surfaces in the literature. See Darboux [1,2], Dierkes [3], Fomenko and Tuzhilin [4], Gray, Salamon, and Abbena [5], Nitsche [6], Osserman [7], Spivak [8] for some books, Lie [9], Schwarz [10], Small [11,12], and Weierstrass [13,14] for some papers related to minimal surfaces in Euclidean geometry.

Lie [9] studied the algebraic minimal surfaces and gave a table classifying these surfaces. See also Enneper [15], Güler [16], Nitsche [6], and Ribaucour [17] for details.

Weierstrass [13] revealed a representation for minimal surfaces in three dimensional Euclidean space  $\mathbb{E}^3$ . Almost one hundred years later, Kobayashi [18] gave an analogous Weierstrass-type representation for conformal spacelike surfaces with mean curvature identically 0, called maximal surfaces, in three dimensional Minkowski space  $\mathbb{E}^{2,1}$ .

In this paper, we consider the Enneper family of maximal surfaces  $\mathcal{E}_m$  for positive integers  $m \geq 1$  by using Weierstrass data  $(1, \zeta^m)$  for  $\zeta \in \mathbb{C}$ , and then show that these surfaces are algebraic in  $\mathbb{E}^{2,1}$ . See Güler [16] for a Euclidean case of Enneper's algebraic minimal surfaces family.

In Section 2, we give this family of real maximal surfaces in  $(r, \theta)$  and  $(u, v)$  coordinates by using Weierstrass representation in  $\mathbb{E}^{2,1}$ . In Section 3, we find irreducible algebraic equations defining surfaces  $\mathcal{E}_m(u, v)$  in terms of running coordinates  $x, y, z$ , and  $a, b, c$ , and also compute degrees and classes of  $\mathcal{E}_m(u, v)$ . Finally, we summarize all findings in tables in the last section, then give some open problems.

## 2. Family of Enneper Maximal Surfaces

Let  $\mathbb{E}^{n,1} := (\{x = (x_1, \dots, x_n, x_0)^t \mid x_i \in \mathbb{R}\}, \langle \cdot, \cdot \rangle)$  be the  $(n + 1)$ -dimensional Lorentz-Minkowski (for short, Minkowski) space with Lorentzian metric  $\langle x, y \rangle = x_1y_1 + \dots + x_ny_n - x_0y_0$ .

A vector  $x \in \mathbb{E}^{n,1}$  is called space-like if  $\langle x, x \rangle > 0$ , time-like if  $\langle x, x \rangle < 0$ , and light-like if  $x \neq 0$  and  $\langle x, x \rangle = 0$ . A surface in  $\mathbb{E}^{n,1}$  is called space-like (resp. time-like, light-like) if the induced metric on the tangent planes is a Riemannian (resp. Lorentzian, degenerate) metric.

Now, let  $\mathbb{E}^{2,1}$  be three dimensional Minkowski space with Lorentzian metric  $\langle \cdot, \cdot \rangle = x_1y_1 + x_2y_2 - x_3y_3$ . We identify  $\vec{x}$  and  $\vec{x}^t$  without further comment.



**Citation:** Güler, E. The Algebraic Surfaces of the Enneper Family of Maximal Surfaces in Three Dimensional Minkowski Space. *Axioms* **2022**, *11*, 4. <https://doi.org/10.3390/axioms11010004>

Academic Editor: Anna Maria Fino

Received: 3 December 2021

Accepted: 20 December 2021

Published: 22 December 2021

**Publisher's Note:** MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



**Copyright:** © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Let  $\mathcal{U}$  be an open subset of  $\mathbb{C}$ . A *maximal curve* is an analytic function  $\vartheta : \mathcal{U} \rightarrow \mathbb{C}^n$  such that  $\langle \vartheta'(\zeta), \vartheta'(\zeta) \rangle = 0$ , where  $\zeta \in \mathcal{U}$ , and  $\vartheta' := \frac{\partial \vartheta}{\partial \zeta}$ . In addition, if  $\langle \vartheta', \overline{\vartheta'} \rangle = |\vartheta'|^2 \neq 0$ , then  $\vartheta$  is a regular maximal curve. We then have maximal surfaces in the associated family of a maximal curve, given by the following Weierstrass representation theorem for ZMC (zero mean curvature) surfaces, or maximal surfaces.

Kobayashi [18] found a Weierstrass type representation for space-like conformal maximal surfaces in  $\mathbb{E}^{2,1}$ :

**Theorem 1.** *Let  $g(\omega)$  be a meromorphic function and let  $f(\omega)$  be a holomorphic function,  $fg^2$  is analytic, defined on a simply connected open subset  $U \subset \mathbb{C}$  such that  $f(\omega)$  does not vanish on  $U$  except at the poles of  $g(\omega)$ . Then,*

$$\mathbf{x}(u, v) = \operatorname{Re} \int^{\zeta} \left( f(1 + g^2), if(1 - g^2), -2fg \right) d\omega, \quad (\zeta = u + iv) \tag{1}$$

is a space-like conformal immersion with mean curvature identically 0 (i.e., space-like conformal maximal surface). Conversely, any spacelike conformal maximal surface can be described in this manner.

Next, we give some facts about Weierstrass data, and a maximal curve to construct some maximal surfaces.

**Definition 1.** *A pair of a meromorphic function  $g$  and a holomorphic function  $f$ ,  $(f, g)$  is called Weierstrass data for a maximal surface.*

**Lemma 1.** *The curve of Enneper of order  $m$ :*

$$\varepsilon_m(\zeta) = \left( \zeta + \frac{\zeta^{2m+1}}{2m+1}, i \left( \zeta - \frac{\zeta^{2m+1}}{2m+1} \right), -\frac{2\zeta^{m+1}}{m+1} \right) \tag{2}$$

is a maximal curve,  $\zeta \in \mathbb{C} - \{0\}$ ,  $i = \sqrt{-1}$ ,  $m \neq -1, -1/2$ .

Therefore, we have  $\langle \varepsilon_m, \varepsilon_m \rangle = 0$  by using (2). Hence, in  $\mathbb{E}^{2,1}$ , the Enneper maximal surface is given by

$$\mathcal{E}_m(u, v) = \operatorname{Re} \int \varepsilon_m(\zeta) d\zeta, \tag{3}$$


where  $\zeta = u + iv$ .  $\operatorname{Im} \int \varepsilon_m(\zeta) d\zeta$  gives the adjoint minimal surface  $\mathcal{E}_m^*(u, v)$  of the surface  $\mathcal{E}_m(u, v)$  in (3). Then, we get the following:

**Corollary 1.** *The Weierstrass data  $(1, \zeta^m)$  of (3) is a representation of the Enneper maximal surface, where integer  $m \geq 1$ .*

Considering the findings above with  $\zeta = re^{i\theta}$ , we get the following Enneper family of maximal surfaces:

$$\mathcal{E}_m(r, \theta) = \begin{pmatrix} r \cos(\theta) + \frac{1}{2m+1} r^{2m+1} \cos[(2m+1)\theta] \\ -r \sin(\theta) + \frac{1}{2m+1} r^{2m+1} \sin[(2m+1)\theta] \\ -\frac{2}{m+1} r^{m+1} \cos[(m+1)\theta] \end{pmatrix} \tag{4}$$

where  $m \neq -1, -1/2$ . See Figure 1 for Enneper maximal surfaces  $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$  in  $(r, \theta)$  coordinates.



**Figure 1.** Enneper maximal surfaces, and its top views (Left):  $\mathcal{E}_1(r, \theta)$ , (Middle):  $\mathcal{E}_2(r, \theta)$ , (Right):  $\mathcal{E}_3(r, \theta)$ .

Hence, using the binomial formula, we obtain more clear representation of  $\mathcal{E}_m(u, v)$  in (3):

$$\begin{aligned}
 x(u, v) &= \operatorname{Re} \left\{ u + iv + \frac{1}{2m+1} \sum_{k=0}^{2m+1} \binom{2m+1}{k} u^{2m+1-k} (iv)^k \right\}, \\
 y(u, v) &= \operatorname{Re} \left\{ iu - v + \frac{i}{2m+1} \sum_{k=0}^{2m+1} \binom{2m+1}{k} u^{2m+1-k} (iv)^k \right\}, \\
 z(u, v) &= \operatorname{Re} \left\{ -\frac{2}{m+1} \sum_{k=0}^{m+1} \binom{m+1}{k} u^{m+1-k} (iv)^k \right\}.
 \end{aligned} \tag{5}$$

We study surface  $\mathcal{E}_m(u, v)$  in  $(u, v)$  coordinates for  $m = 1, 2, \dots, 5$ , taking  $\zeta = u + iv$  at Cartesian coordinates  $x, y, z$ , and also in inhomogeneous tangential coordinates  $a, b, c$ , by using Weierstrass representation equation.

Next, we give a theorem about maximality of surface  $\mathcal{E}_1(u, v)$  (see Figure 1, Left):

**Theorem 2.** *The surface*

$$\mathcal{E}_1(u, v) = \begin{pmatrix} \frac{1}{3}u^3 - uv^2 + u \\ -\frac{1}{3}v^3 + u^2v - v \\ -u^2 + v^2 \end{pmatrix} = \begin{pmatrix} x(u, v) \\ y(u, v) \\ z(u, v) \end{pmatrix} \tag{6}$$

which has Weierstrass data  $(1, \zeta)$ , is an Enneper maximal surface in  $\mathbb{E}^{2,1}$ .

**Proof.** The coefficients of the first fundamental form of the surface  $\mathcal{E}_1(u, v)$  ( $\mathcal{E}_1$ , for short) are given by

$$\begin{aligned}
 E &= (\lambda - 1)^2 = G, \\
 F &= 0,
 \end{aligned}$$

where  $\lambda = u^2 + v^2$ . That is, conformality holds. Then, the Gauss map  $e_1(u, v)$  of  $\mathcal{E}_1$  is as follows

$$e_1 = \left( -\frac{2u}{\lambda - 1}, -\frac{2v}{\lambda - 1}, \frac{\lambda^2 + 1}{\lambda - 1} \right), \tag{7}$$

where  $\lambda \neq 1$ . The coefficients of the second fundamental form of  $\mathcal{E}_1$  are given by

$$L = -\frac{2(3\lambda + 1)}{\lambda - 1} = -N,$$

$$M = 0.$$

Then, we obtain the following mean curvature and the Gaussian curvature of the surface  $\mathcal{E}_1$ :

$$H = 0,$$

$$K = \frac{4(3\lambda + 1)^2}{(\lambda - 1)^6},$$

respectively. Here,  $H = \langle \sigma, \sigma \rangle \frac{EN+GL-2FM}{2(EG-F^2)}$ ,  $K = \langle \sigma, \sigma \rangle \frac{LN-M}{EG-F^2}$ , where  $\langle \sigma, \sigma \rangle = -1$ . Hence, the Enneper surface is maximal surface with positive Gaussian curvature.  $\square$

Therefore, we obtain the following parametric equations of the higher order maximal Enneper surfaces  $\mathcal{E}_m(u, v) = (x(u, v), y(u, v), z(u, v))$  (see Figure 2 Middle for  $\mathcal{E}_2$ , and Figure 2 Right for  $\mathcal{E}_3$ ):

$$\mathcal{E}_2(u, v) = \begin{pmatrix} \frac{1}{5}u^5 - 2u^3v^2 + uv^4 + u \\ \frac{1}{5}v^5 - 2u^2v^3 + u^4v - v \\ -\frac{2}{3}u^3 + 2uv^2 \end{pmatrix}, \tag{8}$$

$$\mathcal{E}_3(u, v) = \begin{pmatrix} \frac{1}{7}u^7 - 3u^5v^2 + 5u^3v^4 - uv^6 + u \\ -\frac{1}{7}v^7 + 3u^2v^5 - 5u^4v^3 + u^6v - v \\ -\frac{1}{2}u^4 + 3u^2v^2 - \frac{1}{2}v^4 \end{pmatrix}, \tag{9}$$

$$\mathcal{E}_4(u, v) = \begin{pmatrix} \frac{1}{9}u^9 - 4u^7v^2 + 14u^5v^4 - \frac{23}{3}u^3v^6 + uv^8 + u \\ \frac{1}{9}v^9 - 4u^2v^7 + 14u^4v^5 - \frac{23}{3}u^6v^3 + u^8v - v \\ -\frac{2}{5}u^5 + 4u^3v^2 - 2uv^4 \end{pmatrix}, \tag{10}$$

$$\mathcal{E}_5(u, v) = \begin{pmatrix} \frac{1}{11}u^{11} - 5u^9v^2 + 30u^7v^4 - 42u^5v^6 + 15u^3v^8 + uv^{10} + u \\ -\frac{1}{11}v^{11} + 5u^2v^9 - 30u^4v^7 + 42u^6v^5 - 15u^8v^3 + u^{10}v - v \\ -\frac{1}{3}u^6 + 5u^4v^2 - 5u^2v^4 + \frac{1}{3}v^6 \end{pmatrix}. \tag{11}$$




Figure 2. Enneper maximal surfaces (Left):  $\mathcal{E}_1(u, v)$ , (Middle):  $\mathcal{E}_2(u, v)$ , (Right):  $\mathcal{E}_3(u, v)$ .

### 3. Degree and Class of Enneper Maximal Surfaces

In this section, using some elimination techniques, we derive the irreducible algebraic surface equation, degree and class of Enneper maximal surfaces family  $\mathcal{E}_m(u, v)$  for integers  $1 \leq m \leq 5$  in three dimensional Minkowski space  $\mathbb{E}^{2,1}$ .

Let us see some basic notions of the surfaces.

**Definition 2.** The set of roots of a polynomial  $Q(x, y, z) = 0$  gives an algebraic surface equation. An algebraic surface  $\mathbf{s}$  is said to be of degree  $\mathbf{d}$  when  $\mathbf{d} = \text{deg}(\mathbf{s})$ .

**Definition 3.** At a point  $(u, v)$  on a surface  $\mathbf{s}(u, v) = (x(u, v), y(u, v), z(u, v))$ , the tangent plane is given by

$$Xx + Yy - Zz + P = 0, \tag{12}$$

where  $e = (X(u, v), Y(u, v), Z(u, v))$  is the Gauss map, and  $P = P(u, v)$ . Then, in inhomogeneous tangential coordinates  $a, b, c$ , we have the following surface:

$$\widehat{\mathbf{s}}(u, v) = (a, b, c) = (X/P, Y/P, Z/P). \tag{13}$$

Therefore, we can obtain an algebraic equation  $\hat{Q}(a, b, c) = 0$  of  $\widehat{\mathbf{s}}(u, v)$  in inhomogeneous tangential coordinates.

**Definition 4.** The maximum degree of the algebraic equation  $\hat{Q}(a, b, c) = 0$  of  $\widehat{\mathbf{s}}(u, v)$  in inhomogeneous tangential coordinates gives the class of  $\widehat{\mathbf{s}}(u, v)$ .

See [6], for details of a Euclidean case. Hence, we obtain the following findings for degrees and classes of Enneper maximal surfaces that we use:

#### 3.1. Degree

We compute the irreducible algebraic surface equation  $Q_1(x, y, z) = 0$  (see Figure 3, Left) of Enneper’s maximal surface  $\mathcal{E}_1(u, v)$  in (6) by using some elimination techniques. We find the following algebraic equation:

$$\begin{aligned} Q_1(x, y, z) = & 64z^9 + 432x^2z^6 - 432y^2z^6 - 1215x^4z^3 - 6318x^2y^2z^3 + 3888x^2z^5 \\ & - 1215y^4z^3 + 3888y^2z^5 - 1152z^7 + 729x^6 - 2187x^4y^2 - 4374x^4z^2 \tag{14} \\ & + 2187x^2y^4 + 6480x^2z^4 - 729y^6 + 4374y^4z^2 - 6480y^2z^4 + 729x^4z \\ & - 1458x^2y^2z - 3888x^2z^3 + 729y^4z - 3888y^2z^3 + 5184z^5. \end{aligned}$$


Then, its degree number is 9.

Next, we continue our computations to find  $Q_m(x, y, z) = 0$  for integers  $m = 2, 3$ . We compute the following irreducible algebraic surface equations  $Q_2(x, y, z) = 0$  (see Figure 3, Middle) and  $Q_3(x, y, z) = 0$  (see Figure 3, Right) of the surfaces  $\mathcal{E}_2(u, v)$  and  $\mathcal{E}_3(u, v)$ , respectively,

$$\begin{aligned} Q_2(x, y, z) = & 847\,288\,609\,443z^{25} - 4358\,480\,501\,250x^3z^{20} \\ & + 13\,075\,441\,503\,750xy^2z^{20} - 131\,157\,978\,046\,875x^6z^{15} \\ & - 474\,186\,536\,015\,625x^4y^2z^{15} + 107 \text{ other lower degree terms,} \end{aligned}$$

$$\begin{aligned}
 Q_3(x, y, z) = & 2475\,880\,078\,570\,760\,549\,798\,248\,448z^{49} \\
 & +5079\,604\,062\,565\,768\,134\,821\,675\,008x^4z^{42} \\
 & -30\,477\,624\,375\,394\,608\,808\,930\,050\,048x^2y^2z^{42} \\
 & +5079\,604\,062\,565\,768\,134\,821\,675\,008y^4z^{42} \\
 & -633\,850\,350\,654\,216\,217\,766\,624\,493\,568x^8z^{35} \\
 & +446 \text{ other lower degree terms.}
 \end{aligned}$$

Therefore,  $Q_m(x, y, z) = 0$  are the algebraic maximal surfaces of the surfaces  $\mathcal{E}_m(u, v)$ , where  $m = 2, 3$ , and they have degree numbers 25 and 49, respectively.



**Figure 3.** Enneper algebraic maximal surfaces (Left):  $Q_1(x, y, z) = 0$ , (Middle):  $Q_2(x, y, z) = 0$ , (Right):  $Q_3(x, y, z) = 0$ .

### 3.2. Class

Now, we introduce the class of the surfaces  $\mathcal{E}_m(u, v)$  for integers  $1 \leq m \leq 4$ . The case  $m = 5$ , marked with “\*” presented in tables of Section 4. Computing the irreducible algebraic surface equations  $\hat{Q}_m(a, b, c) = 0$ , we obtain the Gauss maps  $e_m(u, v)$  (see Figure 4 for  $e_1, e_2, e_3$ ) for integers  $1 \leq m \leq 5$  of the surfaces  $\mathcal{E}_m(u, v)$ , and we also generalize them as follows:

$$e_1 = \left( -2\frac{u}{\lambda - 1}, -2\frac{v}{\lambda - 1}, \frac{\lambda + 1}{\lambda - 1} \right),$$

$$e_2 = \left( -2\frac{u^2 - v^2}{\lambda^2 - 1}, -2\frac{2uv}{\lambda^2 - 1}, \frac{\lambda^2 + 1}{\lambda^2 - 1} \right), \tag{15}$$

$$e_3 = \left( -2\frac{u^3 - 3uv^2}{\lambda^3 - 1}, -2\frac{3u^2v - v^3}{\lambda^3 - 1}, \frac{\lambda^3 + 1}{\lambda^3 - 1} \right), \tag{16}$$

$$e_4 = \left( -2\frac{u^4 - 6u^2v^2 + v^4}{\lambda^4 - 1}, -2\frac{4u^3v - 4uv^3}{\lambda^4 - 1}, \frac{\lambda^4 + 1}{\lambda^4 - 1} \right), \tag{17}$$

$$e_5 = \left( -2\frac{u^5 - 10u^3v^2 + 5uv^4}{\lambda^5 - 1}, -2\frac{5u^4v - 10u^2v^3 + v^5}{\lambda^5 - 1}, \frac{\lambda^5 + 1}{\lambda^5 - 1} \right), \tag{18}$$

⋮

$$e_m = \left( -2\frac{\operatorname{Re}(\zeta^m)}{|\zeta|^m - 1}, -2\frac{\operatorname{Im}(\zeta^m)}{|\zeta|^m - 1}, \frac{|\zeta|^m + 1}{|\zeta|^m - 1} \right), \quad (\zeta = u + iv, |\zeta| = \lambda). \tag{19}$$


Using (6), (7), (12) and (13), with  $P_1(u, v) = \frac{(\lambda-3)(-u^2+v^2)}{3(\lambda-1)}$ , we get the following surface  $\hat{\mathcal{E}}_1(u, v)$  (see Figure 5, Left) in inhomogeneous tangential coordinates:

$$a = \frac{6u}{(-u^2 + v^2)(\lambda - 3)}, \quad b = \frac{6v}{(-u^2 + v^2)(\lambda - 3)}, \quad c = -\frac{3(\lambda + 1)}{(-u^2 + v^2)(\lambda - 3)}.$$

where  $\lambda = u^2 + v^2, \lambda \neq 3, u, v \neq 0$ . Therefore, we compute Enneper’s irreducible algebraic maximal surface equation  $\hat{Q}_1(a, b, c) = 0$  (see Figure 6, Left) of the surface  $\hat{\mathcal{E}}_1(u, v)$ :

$$\hat{Q}_1(a, b, c) = 4a^6 - 4a^4b^2 - 3a^4c^2 - 4a^2b^4 + 6a^2b^2c^2 + 4b^6 - 3b^4c^2 - 18a^4c + 12a^2c^3 + 18b^4c - 12b^2c^3 + 9a^4 + 18a^2b^2 + 9b^4.$$

So, Enneper’s maximal surface  $\mathcal{E}_1(u, v)$  in (6) has class number 6.



**Figure 4.** Top views of the Gauss maps of the surfaces  $\mathcal{E}_{m=1,2,3}(u, v)$  (Left):  $e_1(u, v)$ , (Middle):  $e_2(u, v)$ , (Right):  $e_3(u, v)$ .

Next, we continue our computations to find  $\hat{Q}_m$  for integers 2, 3, 4. To find the class of surface  $\mathcal{E}_2(u, v)$  (see Figure 5, Middle), we use (9), (12), (13) and (16). Calculating  $P_2(u, v) = -\frac{4(u^3 - 3uv^2)(\lambda^2 - 5)}{15(\lambda^2 - 1)}$ , we get the following surface  $\hat{\mathcal{E}}_2$  inhomogeneous tangential coordinates:

$$a = \frac{15(u^2 - v^2)}{2(u^3 - 3uv^2)(\lambda^2 - 5)}, \quad b = \frac{15uv}{(u^3 - 3uv^2)(\lambda^2 - 5)}, \quad c = -\frac{15(\lambda^2 + 1)}{4(u^3 - 3uv^2)(\lambda^2 - 5)},$$

where  $\lambda = u^2 + v^2, \lambda^2 \neq 5, u, v \neq 0$ . In the inhomogeneous tangential coordinates  $a, b, c$ , we find the following irreducible algebraic surface equation  $\hat{Q}_2(a, b, c) = 0$  (see Figure 6, Middle) of the surface  $\hat{\mathcal{E}}_2(u, v)$ :

$$\hat{Q}_2(a, b, c) = 2176782336a^{16}b^4 + 5804752896a^{14}b^6 - 4837294080a^{14}b^4c^2 + 2902376448a^{12}b^8 - 8062156800a^{12}b^6c^2 + 120 \text{ other lower degree terms.}$$

Hence,  $\hat{Q}_2(a, b, c) = 0$  is the algebraic surface of the surface  $\hat{\mathcal{E}}_2(u, v)$ , and Enneper’s maximal surface  $\mathcal{E}_2(u, v)$  in (8) has class number 20.


Using similar ways, we compute the irreducible algebraic surface equation  $\hat{Q}_3(a, b, c) = 0$  (see Figure 6, Right) of surface  $\hat{\mathcal{E}}_3(u, v)$  (see Figure 5, Right) as follows:

$$\hat{Q}_3(a, b, c) = 26623333280885243904a^{42} - 718829998583901585408a^{40}b^2 - 104829374793485647872a^{40}c^2 + 6868819986468392927232a^{38}b^4 + 2935222494217598140416a^{38}b^2c^2 + 774 \text{ other lower degree terms.}$$

$\hat{Q}_3(a, b, c) = 0$  is the algebraic surface of the surface  $\hat{E}_3(u, v)$ , and Enneper’s maximal surface  $E_3(u, v)$  in (9) has class number 42. We also compute the following irreducible algebraic surface equation  $\hat{Q}_4(a, b, c) = 0$  of the surface  $\hat{E}_4(u, v)$ :

$$\begin{aligned} \hat{Q}_4(a, b, c) = & 4294967296000a^{64}b^8 \\ & - 2473901162496000a^{62}b^8c^2 \\ & - 962072674304000a^{60}b^{12} \\ & + 247390116249600a^{60}b^{10}c^2 \\ & + 66795331387392000a^{60}b^8c^4 \\ & + 2604 \text{ other lower degree terms.} \end{aligned}$$

$\hat{Q}_4(a, b, c) = 0$  is an algebraic surface of  $\hat{E}_4(u, v)$ , and Enneper’s maximal surface  $E_4(u, v)$  in (10) has class number 72.



**Figure 5.** Enneper maximal surfaces in inhomogeneous tangential coordinates (Left):  $\hat{E}_1(u, v)$ , (Middle):  $\hat{E}_2(u, v)$ , (Right):  $\hat{E}_3(u, v)$ .

We obtain the following functions  $P_i(u, v)$ , where  $1 \leq i \leq 6$ ,

$$\begin{aligned} P_1(u, v) &= -\frac{(u^2 - v^2)(\lambda - 3)}{3(\lambda - 1)}, & P_2(u, v) &= -\frac{4(u^3 - 3uv^2)(\lambda^2 - 5)}{15(\lambda^2 - 1)}, \\ P_3(u, v) &= -\frac{3(u^4 - 6u^2v^2 + v^4)(\lambda^3 - 7)}{14(\lambda^3 - 1)}, & P_4(u, v) &= -\frac{8(u^5 - 10u^3v^2 + 5uv^4)(\lambda^4 - 9)}{45(\lambda^4 - 1)}, \\ P_5(u, v) &= -\frac{5(u^6 - 15u^4v^2 + 15u^2v^4 - v^6)(\lambda^5 - 11)}{33(\lambda^5 - 1)}, & P_6(u, v) &= -\frac{12(u^7 - 21u^5v^2 + 35u^3v^4 - 7uv^6)(\lambda^6 - 13)}{91(\lambda^6 - 1)}. \end{aligned}$$

We generalize the above functions, and give the following results:

**Corollary 2.** *The functions  $P_{m \geq 1}$  for integers  $m$ , are given by*

$$\begin{aligned} P_{2k-1} &= -\frac{2(2k - 1) [\lambda^{2k-1} - (2k + 1)]}{(k + 1)(2k + 1)(\lambda^{2k-1} - 1)} \operatorname{Re}(\zeta^{2k}), \\ P_{2k} &= -\frac{4k [\lambda^{2k} - (2k + 1)]}{(2k + 1)(4k + 1)(\lambda^{2k} - 1)} \operatorname{Re}(\zeta^{2k+1}), \end{aligned}$$



where integers  $k \geq 1$ ,  $\zeta = u + iv$  and  $|\zeta| = \lambda$ .

So far, we find surfaces  $\hat{\mathcal{E}}_1$  and  $\hat{\mathcal{E}}_2$ . By using  $\mathcal{E}_3 - \mathcal{E}_5$ ,  $e_3 - e_5$ , and also (12), (13), we obtain the following surfaces:  $\hat{\mathcal{E}}_m(u, v) = (a, b, c)$ :

$$\begin{aligned} \hat{\mathcal{E}}_1 &= -\frac{3}{(u^2 - v^2)(\lambda - 3)} \begin{pmatrix} -2u \\ -2v \\ \lambda + 1 \end{pmatrix}, \\ \hat{\mathcal{E}}_2 &= -\frac{15}{4(u^3 - 3uv^2)(\lambda^2 - 5)} \begin{pmatrix} -2(u^2 - v^2) \\ -4uv \\ \lambda^2 + 1 \end{pmatrix}, \\ \hat{\mathcal{E}}_3 &= -\frac{14}{3(u^4 - 6u^2v^2 + v^4)(\lambda^3 - 7)} \begin{pmatrix} -2(u^3 - 3uv^2) \\ -2(u^2v - v^3) \\ \lambda^3 + 1 \end{pmatrix}, \\ \hat{\mathcal{E}}_4 &= -\frac{45}{8(u^5 - 10u^3v^2 + 5uv^4)(\lambda^4 - 9)} \begin{pmatrix} -2(u^4 - 6u^2v^2 + v^4) \\ -2(4u^3v - 4uv^3) \\ \lambda^4 + 1 \end{pmatrix}, \\ \hat{\mathcal{E}}_5 &= -\frac{33}{5(u^6 - 15u^4v^2 + 15u^2v^4 - v^6)(\lambda^5 - 11)} \begin{pmatrix} -2(u^5 - 10u^3v^2 + 5uv^4) \\ -2(5u^4v - 10u^2v^3 + v^5) \\ \lambda^5 + 1 \end{pmatrix}. \end{aligned}$$

We also generalize the above functions, and find the following results:

**Corollary 3.** The surfaces  $\hat{\mathcal{S}}_{m \geq 1}(u, v)$  for integers  $m$ , are given by


$$\begin{aligned} \hat{\mathcal{E}}_{2k-1}(u, v) &= -\frac{k(4k-1)}{(2k-1)[\lambda^{2k-1} - (2k+1)]\text{Re}(\zeta^{2k})} \begin{pmatrix} -2\text{Re}(\zeta^{2k-1}) \\ -2\text{Im}(\zeta^{2k-1}) \\ |\zeta|^{2k-1} + 1 \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}, \\ \hat{\mathcal{E}}_{2k}(u, v) &= -\frac{(2k+1)(4k+1)}{4k[\lambda^{2k} - (4k+1)]\text{Re}(\zeta^{2k+1})} \begin{pmatrix} -2\text{Re}(\zeta^{2k}) \\ -2\text{Im}(\zeta^{2k}) \\ |\zeta|^{2k} + 1 \end{pmatrix} = \begin{pmatrix} a \\ b \\ c \end{pmatrix}, \end{aligned}$$

where integers  $k \geq 1$ ,  $\zeta = u + iv$  and  $|\zeta| = \lambda$ .

**Corollary 4.** In  $\mathbb{E}^{2,1}$ , the relations between the Enneper maximal surface  $\hat{\mathcal{E}}_{m \geq 1}(u, v)$  in the inhomogeneous tangential coordinates and the Gauss map  $e_{m \geq 1}(u, v)$  of the Enneper maximal surface  $\mathcal{E}_{m \geq 1}(u, v)$  in the cartesian coordinates are given by

$$\begin{aligned} \hat{\mathcal{E}}_{2k-1}(u, v) &= -\frac{k(4k-1)(\lambda^{2k-1} - 1)}{(2k-1)[\lambda^{2k-1} - (2k+1)]\text{Re}(\zeta^{2k})} e_{2k-1}(u, v), \\ \hat{\mathcal{E}}_{2k}(u, v) &= -\frac{(2k+1)(4k+1)(\lambda^{2k} - 1)}{4k[\lambda^{2k} - (2k+1)]\text{Re}(\zeta^{2k+1})} e_{2k}(u, v), \end{aligned}$$

where integers  $k \geq 1$ ,  $\zeta = u + iv$  and  $|\zeta| = \lambda$ .



**Figure 6.** Enneper’s algebraic maximal surfaces in inhomogeneous tangential coordinates (Left):  $\hat{Q}_1(a, b, c) = 0$ , (Middle):  $\hat{Q}_2(a, b, c) = 0$ , (Right):  $\hat{Q}_3(a, b, c) = 0$ .

**4. Conclusions**

To reveal the irreducible algebraic surface equations of the Enneper maximal surfaces  $\mathcal{E}_m(u, v)$  in  $\mathbb{E}^{2,1}$ , we have tried a series of standard techniques in elimination theory: only Sylvester by hand for  $Q_1(x, y, z) = 0$ , and then projective (Macaulay) and sparse multivariate resultants implemented in the Maple software [19] package multires for  $Q_m(x, y, z) = 0$  and  $\hat{Q}_m(a, b, c) = 0$ .

Maple’s native implicitization command Implicitize, and implicitization based on Maple’s native implementation of the Groebner Basis. For the latter, we implemented in Maple the method in [20] (Chapter 3, p. 128). Under reasonable time, we only succeed for  $m = 1, 2$  in all above methods.

For  $m = 3$ , the successful method we have tried was to compute the equation defining the elimination ideal using the Groebner Basis package FGb of Faugère in [21].

The time required to output the irreducible algebraic surface equations  $Q_m(x, y, z) = 0$  (resp.  $\hat{Q}_m(a, b, c) = 0$ ) for integers  $1 \leq m \leq 3$  and polynomials defining the elimination ideal was under reasonable seconds determined by the following Table 1 (resp. Table 2).

For the degree (resp. class) of the irreducible algebraic surface equation  $Q_4(x, y, z) = 0$  (resp.  $\hat{Q}_5(a, b, c) = 0$ ) of the surface  $\mathcal{E}_4(u, v)$  (resp.  $\hat{\mathcal{E}}_5(u, v)$ ), marked with “\*” in Table 1 (resp. Table 2), was rejected (i.e., “out of memory”) by Maple 17 on a laptop Pentium Core i5-4310M 2.00 GHz, 4 GB RAM, with the time given in CPU seconds.

Hence, we propose the following:

**Proposition 1.** For integers  $m \geq 1$ , degree number of the irreducible algebraic surfaces  $Q_m(x, y, z) = 0$  in the Cartesian coordinates is of  $(2m + 1)^2$ , and class number of irreducible algebraic surfaces  $\hat{Q}_m(a, b, c) = 0$  in inhomogeneous tangential coordinates is of  $2m(2m + 1)$  of the  $(1, \zeta^m)$ -type real Enneper maximal surfaces  $\mathcal{E}_m(u, v)$ .

*Open Problems*

Here, we give some problems that we could not find the answers in this paper:

**Problem 1.** Find the irreducible Enneper algebraic maximal surface eq.  $Q_{m \geq 4}(x, y, z) = 0$  in the cartesian coordinates by using the parametric equation of the Enneper maximal surface  $\mathcal{E}_{m \geq 4}(u, v)$ .

**Problem 2.** Find the irreducible Enneper algebraic maximal surface eq.  $\hat{Q}_{m \geq 5}(a, b, c) = 0$  in the inhomogeneous tangential coordinates by using the parametric equation of the Enneper maximal surface  $\hat{\mathcal{E}}_{m \geq 5}(u, v)$ .

Finally, we give all findings in Tables 1 and 2.

**Table 1.** Results for the Enneper algebraic maximal surfaces  $Q_m(x, y, z) = 0$ .

Algebraic Surface	Degree of Surface	Number of Terms	Gröbner Time (s)	FGb Time (s)
$Q_1$	9	23	0.266	0.041
$Q_2$	25	112	321.953	0.835
$Q_3$	49	451	*	266.854
$Q_4$	81	*	*	*
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$Q_m$	$(2m + 1)^2$	*	*	*

**Table 2.** Results for the Enneper algebraic maximal surfaces  $\hat{Q}_m(a, b, c) = 0$ .

Algebraic Surface	Class of Surface	Number of Terms	Gröbner Time (s)	FGb Time (s)
$\hat{Q}_1$	6	14	0.94	0.030
$\hat{Q}_2$	20	125	61.152	0.114
$\hat{Q}_3$	42	779	*	125.904
$\hat{Q}_4$	72	2609	*	1306.718
$\hat{Q}_5$	110	*	*	*
$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$
$\hat{Q}_m$	$2m(2m + 1)$	*	*	*

**Funding:** This research received no external funding.

**Institutional Review Board Statement:** Not applicable.

**Informed Consent Statement:** Not applicable.

**Data Availability Statement:** Not applicable.

**Acknowledgments:** The author is grateful to the referees for their suggestions, which improved the presentation of the paper.

**Conflicts of Interest:** The author declares no conflict of interest.

**References**

1. Darboux, G. *Leçons Sur la Théorie Générale des Surfaces, I, II, (French) [Lessons on the General Theory of Surfaces, I, II]*; Reprint of the Second (1914) Edition (I) and the Second (1915) Edition (II), Les Grands Classiques Gauthier-Villars [Gauthier-Villars Great Classics], Cours de Géométrie de la Faculté des Sciences [Course on Geometry of the Faculty of Science]; Éditions Jacques Gabay: Sceaux, France, 1993.
2. Darboux, G. *Leçons Sur la Théorie Générale des Surfaces, III, IV, (French) [Lessons on the General Theory of Surfaces, III, IV]*; Reprint of the 1894 Original (III) and the 1896 Original (IV), Les Grands Classiques Gauthier-Villars [Gauthier-Villars Great Classics], Cours de Géométrie de la Faculté des Sciences [Course on Geometry of the Faculty of Science]; Éditions Jacques Gabay: Sceaux, France, 1993.
3. Dierkes, U.; Hildebrandt, S. *Sauvigny F. Minimal Surfaces*, 2nd ed.; Springer: Berlin/Heidelberg, Germany, 2010.
4. Fomenko, A.T.; Tuzhilin, A.A. *Elements of the Geometry and Topology of Minimal Surfaces in Three-Dimensional Space, Translated from the Russian by E.J.F. Primrose*; Translations of Mathematical Monographs, 93; American Math. Soc.: Providence, RI, USA, 1991.
5. Gray, A.; Salamon, S.; Abbena, E. *Modern Differential Geometry of Curves and Surfaces with Mathematica*, 3rd ed.; Chapman & Hall/CRC Press: Boca Raton, FL, USA, 2006.
6. Nitsche, J.C.C. *Lectures on Minimal Surfaces. Volume 1. Introduction, Fundamentals, Geometry and Basic Boundary Value Problems*; Cambridge University Press: Cambridge, MA, USA, 1989.
7. Osserman, R. *A Survey of Minimal Surfaces*; Van Nostrand Reinhold Co.: New York, NY, USA, 1969.
8. Spivak, M. *A Comprehensive Introduction to Differential Geometry*, 3rd ed.; Publish or Perish, Inc.: Houston, TX, USA, 1999; Volume IV.
9. Lie, S. Beiträge zur Theorie der Minimalflächen. *Math. Ann.* **1878**, *14*, 331–416. [[CrossRef](#)]
10. Schwarz, H.A. Miscellen aus dem Gebiete der Minimalflächen. *J. für die reine und Angew. Math. (Crelle's J.)* **1875**, *80*, 280–300.
11. Small, A.J. Minimal surfaces in  $\mathbb{R}^3$  and algebraic curves. *Differ. Geom. Appl.* **1992**, *2*, 369–384. [[CrossRef](#)]

12. Small, A.J. Linear structures on the collections of minimal surfaces in  $\mathbb{R}^3$  and  $\mathbb{R}^4$ . *Ann. Glob. Anal. Geom.* **1994**, *12*, 97–101. [[CrossRef](#)]
13. Weierstrass, K. *Untersuchungen über die Flächen, deren Mittlere Krümmung überall Gleich Null ist*; Akademie der Wissenschaften zu Berlin: Berlin, Germany, 1866; pp. 612–625.
14. Weierstrass, K. *Über die analytische Darstellbarkeit Sogenannter Willkürlicher Functionen einer reellen Veränderlichen*; Akademie der Wissenschaften zu Berlin: Berlin, Germany, 1885; pp. 633–639, 789–805.
15. Enneper, A. Untersuchungen über einige Punkte aus der allgemeinen Theorie der Flächen. *Math. Ann.* **1870**, *2*, 58–623. [[CrossRef](#)]
16. Güler, E. Family of Enneper minimal surfaces. *Mathematics* **2018**, *6*, 281. [[CrossRef](#)]
17. Ribaucour, A. Etude des elassoides ou surfaces a courbure moyenne nulle. *Mem. Cour. et Mem. Sav. Etr. Acad. R. Sci. Belg. Bruxelles* **1882**, *44*, 236.
18. Kobayashi, O. Maximal surfaces in the 3-dimensional Minkowski space  $L^3$ . *Tokyo J. Math.* **1983**, *6*, 297–309. [[CrossRef](#)]
19. *Maple Software, (Version 17)*; Waterloo Maple Inc.: Waterloo, ON, Canada. Available online: <https://www.maplesoft.com/> (accessed on 10 October 2021).
20. Cox, D.; Little, J.; O’Shea, D. *Ideals, Varieties, and Algorithms. An Introduction to Computational Algebraic Geometry and Commutative Algebra*, 3rd ed.; Undergraduate Texts in Mathematics; Springer: New York, NY, USA, 2007.
21. Faugère, J.C. FGB: A library for computing Gröbner bases. In Proceedings of the Third International Congress Conference on Mathematical Software (ICMS’10), Kobe, Japan, 13–17 September 2010; Springer: Berlin/Heidelberg, Germany, 2010; pp. 84–87.

Reproduced with permission of copyright owner. Further reproduction prohibited without permission.