

# Plant Polyphenols Stimulate Adhesion to Intestinal Mucosa and Induce Proteome Changes in the Probiotic *Lactobacillus acidophilus* NCFM

Hasan Ufuk Celebioglu, Marta Delsoglio, Susanne Brix, Enrica Pessione, and Birte Svensson\*

**Scope:** Plant phenolics, known to exert beneficial effects on human health, were supplemented to cultures of the probiotic bacterium *Lactobacillus acidophilus* NCFM (NCFM) to assess their effect on its adhesive capacity and the abundance of individual proteins.

**Methods and results:** The presence of resveratrol and ferulic acid during bacterial growth stimulated adhesion of NCFM to mucin and human intestinal HT-29 cells, while tannic acid improved adhesion only to HT-29 cells and caffeic acid had very modest effect overall. Some dosage dependence was found for the four phenolics supplemented at 100, 250, and 500  $\mu\text{g mL}^{-1}$  to the cultures. Notably, 500  $\mu\text{g mL}^{-1}$  ferulic acid only stimulated adhesion to mucin. Analyses of differential whole-cell as well as surface proteomes revealed relative abundance changes for a total of 27 and 22 NCFM proteins, respectively. These changes include enzymes acting in metabolic pathways, such as glycolysis, nucleotide metabolism, and stress response, as well as known moonlighting or surface-associated proteins.

**Conclusion:** The five plant phenolics found in various foods stimulate the adhesive capacity of NCFM in diverse ways and elicit relative abundance changes of specific proteins, providing molecular level insight into the mechanism of the putative beneficial effects of the polyphenols.

pathogenic microorganisms and insects in plant defense.<sup>[2]</sup> We selected five common plant phenolics to explore their impact on the widely used probiotic bacterium *Lactobacillus acidophilus* NCFM (NCFM). Resveratrol (3,5,4'-trihydroxy-trans-stilbene) is found in grapes, wine, peanuts, pistachios, and berries and has been reported to have a large number of health-promoting effects as an antioxidant, anti-inflammatory, antitumor, anti-platelet aggregation, cardioprotective, and longevity agent.<sup>[3]</sup> Tannic acid belongs to tannins, widely distributed in plant-based human diet, and possesses radical scavenging, antimicrobial, and anti-carcinogenic properties.<sup>[4]</sup> Caffeic acid (3,4-dihydroxycinnamic) found in cereals, berries, herbs, and spices is produced in the shikimate pathway<sup>[5]</sup> and can be esterified by quinic acid to chlorogenic acid in fruits, vegetables, coffee, and tobacco.<sup>[6]</sup> The closely related ferulic acid (4-hydroxy-3-methoxycinnamic acid) together with dihydroferulic acid is

abundant in plant cell walls as a constituent of lignocellulose.<sup>[7]</sup> Ferulic acid is identified as monomer, dimer, free oligomer or in polymers esterifying polysaccharides, polyamines, and glycoproteins,<sup>[8]</sup> and high amounts are found in common foods such as bran, vegetables, fruits, and herbs.<sup>[9]</sup> Finally, salicin, a phenolic glycoside from willow bark that possesses analgesic and anti-rheumatic activities,<sup>[10]</sup> can be carbon source for NCFM.

NCFM is a Gram-positive, homofermentative, rod-shaped lactic acid bacterium (LAB) residing in the gastrointestinal tract<sup>[11]</sup> and used in dairy products and dietary supplements.<sup>[12]</sup> NCFM has excellent ability to adapt its metabolism in response to gut nutrients by negative transcriptional regulation enabling survival in nutrient-scarce competitive environments.<sup>[13]</sup> Its adaptation to the gastrointestinal tract (GIT) involves mucus-binding proteins belonging to the surface layer, as previously identified.<sup>[14,15]</sup> NCFM shows excellent stability in dairy and fermented products.<sup>[16]</sup>

Phenolics and probiotics are very important food components residing in the same environment (the GIT) and providing health benefits to humans. The present study describes effects of typical food phenolics on the adhesive capacity of NCFM toward mucin

## 1. Introduction

Plant phenolics are secondary metabolites abundant in foods and beverages where they confer bitterness, astringency, color, flavor, odor, and oxidative stability.<sup>[1]</sup> Polyphenols can interact with molecular targets in various organisms and compete with

Dr. H. U. Celebioglu, M. Delsoglio, Dr. S. Brix, Prof. B. Svensson  
Department of Biotechnology and Biomedicine  
Technical University of Denmark  
Kongens Lyngby, Denmark  
E-mail: bis@bio.dtu.dk

Dr. H. U. Celebioglu  
Department of Biotechnology  
Bartın University  
Bartın, Turkey

M. Delsoglio, Prof. E. Pessione  
Department of Life Sciences and Systems Biology  
University of Turin  
Turin, Italy

DOI: 10.1002/mnfr.201700638

and intestinal HT-29 cells and gains insights into the molecular mechanisms involved in adhesion by analysis of whole cell and surface-associated proteome changes.

## 2. Experimental Section

### 2.1. Growth Conditions

*L. acidophilus* NCFM ( $1.50 \times 10^{10}$  CFU g<sup>-1</sup>, DuPont) was grown aerobically without shaking at 37 °C in 50 mL batch culture on semisynthetic lactic acid bacteria medium (LABSEM)<sup>[17]</sup> containing 1% glucose as carbon source. Stock solutions of phenolics were prepared in ethanol (50 mg mL<sup>-1</sup>), except tannic acid (prepared in distilled water). Different cultures from the same bacterial stock were supplemented with final concentrations of 100 or 250 μg mL<sup>-1</sup> resveratrol (Veri-te resveratrol; kind gift of Evolva, Denmark) or tannic acid (Sigma); 100, 250, and 500 μg mL<sup>-1</sup> caffeic or ferulic acids (both Sigma) and were sub-cultured for three cycles. NCFM release glucose from salicin (1%) that was used as sole carbon source and compared to glucose-grown bacteria. Cells for proteome analyses were grown in the presence of 100 μg mL<sup>-1</sup> resveratrol or tannic acid and 500 μg mL<sup>-1</sup> caffeic or ferulic acids and were sub-cultured for three cycles. Cultures (glucose-grown) without phenolics served as control.

### 2.2. In Vitro Adhesion to Mucin and HT-29 Cells

Adhesion was measured as previously<sup>[18]</sup> described with some modification.<sup>[19,20]</sup> Briefly, freshly late-log phase grown NCFM (20 h, OD 0.5 for tannic acid; 24 h, OD 1.0–1.1 for control and resveratrol; 24 h, OD 1.4–1.5 for caffeic and ferulic acids) was labeled with 100 μM 5(6)-carboxyfluorescein diacetate (Sigma-Aldrich) in PBS (37 °C, 30 min), washed twice and resuspended in PBS to OD<sub>600</sub> 0.5 ± 0.05. A 96-well microtiter plate (Greiner Bioone) was coated with porcine mucin (1 mg mL<sup>-1</sup>, Sigma-Aldrich) in PBS (200 μL well<sup>-1</sup>; 4 °C, overnight). After decanting the mucin solution, wells were washed with PBS, added labeled NCFM (200 μL, OD<sub>600</sub> 0.5), incubated (2 h, 37 °C), followed by decanting the bacterial suspension and washing of wells thrice with PBS. Adhered bacteria were lysed by 1% (w/v) SDS in 0.1 M NaOH (200 μL; 1 h, 37 °C) and quantified by fluorescence measurements (Cytation5 Cell Imaging Multi-Mode Reader, BioTek) using 485 nm and 538 nm as excitation and emission wavelengths, respectively. Adhesion was expressed as the percentage of fluorescence recovered from the lysed bacteria that were bound versus the fluorescence of the total bacterial suspension added to the wells. Three independent experiments were conducted, each in quadruplicate, and data were subjected to one-way ANOVA using OriginPro v9. Human colonic HT-29 cell line (American Type Culture Collection, ATCC HTB38) was cultured and maintained according to the supplier's instructions. HT-29 cells in complete growth medium containing McCoy's 5a medium (ATCC) supplemented with 10% heat-inactivated fetal bovine serum (Sigma-Aldrich) and 1% penicillin-streptomycin (Lonza) were seeded in 24-well plates and cultivated until a confluent state, followed by removing the medium and washing with PBS to remove remain-

ing antibiotics. Labeled NCFM (500 μL, OD<sub>600</sub> 0.5) was added to the wells (2 h) and adhesion analyzed as in mucin adhesion assay.

### 2.3. Sample Preparation for Proteome Analyses

NCFM in late-log phase (OD<sub>600</sub> 0.5–1.6, Supporting Information Figure 1) was harvested by centrifugation (3200 × g, 10 min) and washed with 0.9% NaCl. Extracts were prepared by mechanical grinding (5 × 1-min vortex at maximum speed) with a small amount of acid-washed glass beads (<100 μm diameter; Sigma) in sample buffer (28 mM Tris-HCl, 22 mM Tris-base pH 8.5, 100 mM DTT) containing protease inhibitors (complete, MiniProtease Inhibitor Tablets, Roche). Following heating (100 °C, 2 min) and addition of rehydration buffer (7 M urea, 2 M thiourea, 2% CHAPS, 20 mM DTT), mixtures were vortexed, centrifuged (10 000 × g, 10 min), and supernatants were collected. Surface proteomes were prepared by incubating cell pellets with 5 M lithium chloride (Sigma-Aldrich) (30 min, R.T.) and collecting supernatants after centrifugation (15 000 × g, 15 min).<sup>[21]</sup> Proteins were precipitated by TCA/acetone,<sup>[22]</sup> washed with acetone, dissolved in rehydration buffer and concentrations were determined by using 2D Quant Kit (GE Life Sciences).

### 2.4. CyDye Labeling and Differential Gel Electrophoresis (DIGE)

Proteins in whole cell extracts were CyDye minimal-labeled for DIGE analysis using a dye-swapping approach.<sup>[17,23]</sup> Briefly, protein aliquots (50 μg) from four biological replicates were labeled interchangeably with 250 pM of either Cy5 or Cy3, vortexed, and left in the dark (30 min, 4 °C). For internal standard aliquots from both samples (25 μg protein of each) were combined and labeled with 250 pM Cy2. Labeling was quenched by 1 μL 10 mM lysine in the dark (10 min). Labeled internal standard and samples were mixed and adjusted to 450 μL with rehydration buffer (8 M urea, 2 M thiourea, 33 mM CHAPS, 195 mM DTT, 1% phosphate pH 4–7; GE Life Sciences). Separation in the first dimension (IEF) using IPG strips (pH 3–10; 18 cm Ettan IPGphor; GE Lifesciences) was performed after rehydration (20 °C, 12 h, 30 V) at a total of 65 kWh. Subsequently, the strips were equilibrated 2 × 15 min in 5 mL equilibration buffer (6 M urea, 30% glycerol, 50 mM Tris-HCl, pH 8.8, 2% SDS, 0.01% bromophenol blue) supplemented with 65 mM DTT and 135 mM iodoacetamide in first and second step, respectively. The second dimension (12.5% SDS-PAGE) was run overnight (EttanDALTSix Electrophoresis unit; GE Lifesciences) at 1 W/gel until the dye front reached the gel bottom. The gels were image-analyzed immediately after the second dimension, using excitation/emission wavelengths of Cy2 (488/520 nm), Cy3 (532/580 nm), and Cy5 (633/670 nm) (100 μm resolution; Typhoon 9410 Variable Mode Imager; GE Lifesciences).

### 2.5. 2-DE PAGE

First-dimension separation was performed using IPG strips (pH 3–10, 11, or 18 cm; GE Healthcare) on Ettan IPGphor (GE

Healthcare) to a total of 65 kVh and subsequently the strips were treated as in Section 2.4 first with DTT and then with iodoacetamide. Second dimension (SDS-PAGE) was run with 12.5% Tris-HCl gels for 45 min at 2 W/gel and 4 h at 12 W/gel. The gels were stained by colloidal CBB<sup>[24]</sup> and scanned (Microtek Scan maker 9800 XL; Microtek).

## 2.6. Image Analysis

Gel images were aligned by automated calculation of manually assigned landmark vectors (Progenesis SameSpots v3.3). Scanned gels were analyzed by intra-gel (difference in-gel) and inter-gel (biological variance) analysis. Spot volume ratio change of  $\geq 1.3$  fold and ANOVA  $p \leq 0.05$  were chosen as criteria for identification of proteins showing abundance differences.

## 2.7. In-Gel Digestion and Protein Identification by MS

Differentially abundant spots were excised manually from gels, subjected to in-gel digestion by trypsin and MS identification.<sup>[17]</sup> Briefly, gel pieces were washed in 40% ethanol until colorless, followed by 100% ACN, incubated with 5  $\mu\text{L}$  12.5 ng  $\text{mL}^{-1}$  trypsin (Promega) in 10 mM ammonium bicarbonate (45 min, on ice), added 10 mM ammonium bicarbonate for rehydration, and kept at 37 °C overnight. Supernatant (1  $\mu\text{L}$ ) was applied onto an Anchor Chip target (Bruker-Daltonics), added 1  $\mu\text{L}$  matrix (0.5 mg  $\text{mL}^{-1}$   $\alpha$ -cyano-4-hydroxycinnamic acid in 90% ACN, 0.1% TFA) and washed with 2  $\mu\text{L}$  0.02% TFA. MALDI-TOF MS spectra were obtained (Ultraflex II; Bruker-Daltonics) in auto-mode using Flex Control v3.0 and processed by Flex Analysis v3.0 (both Bruker-Daltonics). Spectra were externally calibrated using a trypsin digest of  $\beta$ -lactoglobulin (5 nM). MS spectra were searched against the NCBI database for bacteria using the MASCOT 2.0 software (<http://www.matrixscience.com>) integrated with BioTools v3.1 (Bruker-Daltonics). Protein identifications by PMF were confirmed with a MASCOT score of 80,  $p \leq 0.05$  and a minimum of six matched peptides.

## 3. Results

The effects of five plant phenolics on the probiotic *L. acidophilus* NCFM were monitored by determining adhesive capacity onto a mucin coating and an HT-29 cell layer as well as by differential whole cell and surface proteome analyses.

### 3.1. Effects of Phenolics on Adhesion

The adhesive ability of NCFM varied with the different phenolics and their concentrations supplemented during growth. Resveratrol (100  $\mu\text{g mL}^{-1}$ ) significantly increased ( $p < 0.05$ ) adhesion to mucin and HT-29 cells by +2.4 and +1.4 fold, respectively, and tannic acid (100  $\mu\text{g mL}^{-1}$ ) increased the adhesion to HT-29 cells by +5.1 fold, compared to a control grown on glucose alone (Figure 1). By contrast, ferulic acid (100  $\mu\text{g mL}^{-1}$ ) and tan-

nic acid (250  $\mu\text{g mL}^{-1}$ ) reduced ( $p < 0.05$ ) adhesion to mucin by  $-1.4$  and  $-3.2$  fold, respectively, whereas caffeic and ferulic acids (250  $\mu\text{g mL}^{-1}$ ) increased ( $p < 0.05$ ) adhesion to mucin by +1.3 and +2.0 folds, respectively. Notably, adhesion to HT-29 cells increased significantly after growth in the presence of 250  $\mu\text{g mL}^{-1}$  resveratrol (+2.3 fold) and ferulic acid (+1.6 fold). At the highest phenolic concentration (500  $\mu\text{g mL}^{-1}$ ) only ferulic acid stimulated adhesion and only to mucin (+1.3 fold). Although salicin was used previously as carbon source for selective enumeration of NCFM,<sup>[25]</sup> growth in 1% salicin did not modulate adhesion (Figure 2).

### 3.2. Effects of Phenolics on the Whole-Cell Proteome


Phenolic-treated bacteria and controls were analyzed by comparative proteomics. Growth at the presence of 100  $\mu\text{g mL}^{-1}$  resveratrol (Supporting Information Figure 1) altered relative abundancies of 12 protein spots in the whole-cell proteome (Figure 2, Table 1, Supporting Information Table 1, Supporting Information File 1a), 11 showing +1.3 to +2.0 fold increase (pre-protein translocase subunit SecA, multiple sugar-binding ABC-transporter ATPase, tRNA N6-adenosine(37)-threonylcarbamoyl transferase complex transferase subunit TsaD (two spots), oxalyl-CoA decarboxylase, ribokinase, molecular chaperone Hsp33, and alanyl-tRNA synthetase), whereas a spot containing one of several glyceraldehyde-3-p dehydrogenase (GAPDH) forms present in the proteome showed  $-1.5$  fold lower relative abundance compared to the control.

Tannic acid (100  $\mu\text{g mL}^{-1}$ , Supporting Information Figure 1) similarly altered abundance for 11 protein spots (Figure 2, Table 1, Supporting Information Table 2, Supporting Information File 1b). Ten increased +1.4 to +2.1 fold (DNA-binding response regulator, prolyl-tRNA synthetase, pyruvate kinase, GAPDH (two spots), glutamyl-tRNA synthetase, oligoendopeptidase F, and adenylosuccinate lyase), whereas tRNA (guanine-N(7)-methyltransferase) was  $-3.0$  fold less abundant.


Growth in the presence of caffeic acid (500  $\mu\text{g mL}^{-1}$ , Supporting Information Figure 1) slightly increased abundance for D-lactate dehydrogenase (+1.4 fold) and elongation factor Ts (+1.3 fold), while it decreased abundance for GAPDH ( $-2.3$  fold), 50S ribosomal protein L1 ( $-1.5$  fold), and heat shock protein Hsp33 ( $-1.3$  fold) (Figure 3, Table 1, Supporting Information Table 3, Supporting Information File 1c). Ferulic acid (500  $\mu\text{g mL}^{-1}$ , Supporting Information Figure 1) increased transcriptional regulator LBA0733 (+1.9 fold), while purine *trans*-deoxyribosylase, 50S ribosomal protein L1, fructokinase, ribose-p-pyrophosphokinase, and phosphomethylpyrimidine kinase decreased in abundance ( $-1.4$  to  $-1.7$  fold) (Figure 3, Table 1, Supporting Information Table 4, Supporting Information File 1d).

### 3.3. Effects of Phenolics on the Surface Proteome

Differential abundancies in the surface proteome were previously reported for NCFM to accompany increased adhesion when using plant-derived oligosaccharides as carbon sources.<sup>[17,23]</sup> Addition of resveratrol (100  $\mu\text{g mL}^{-1}$ ) during growth altered nine



**Figure 1.** In vitro adhesion of *L. acidophilus* NCFM. a) Adhesion to mucin coating by bacteria grown with glucose as carbon source in the presence of resveratrol, tannic acid, caffeic acid, ferulic acid, or with salicin as carbon source. b) Adhesion to HT-29 cells by bacteria exposed during growth to the phenolics as in mucin adhesion assay. Asterisks indicate statistically significant differences (\*,  $p \leq 0.05$ ; \*\*,  $p \leq 0.01$ ; \*\*\*,  $p \leq 0.001$ ) as compared to control.



**Figure 2.** Representative whole cell 2D proteome map of *L. acidophilus* NCFM grown with glucose as carbon source in the presence of resveratrol or tannic acid. Numbers indicate differentially abundant spots (ANOVA  $p \leq 0.05$ ), compared to control (non-treated), identified by in-gel digestion and mass spectrometry. NCFM was grown in the presence of  $100 \mu\text{g mL}^{-1}$  resveratrol (Res-treated) and  $100 \mu\text{g mL}^{-1}$  tannic acid (TA-treated). Selected spots are shown for relative abundance.

spots in the surface proteome. Five increased in abundance by +1.4 to +2.1 fold (pyruvate kinase, 50S ribosomal protein L7/L12, elongation factor P, 50S ribosomal protein L22 and hypothetical protein LBA1769) and four decreased  $-2.0$  to  $-1.4$  fold (GAPDH (two spots), adenylosuccinate synthetase, and 6-phosphofructokinase) (Figure 4A, Table 2, Supporting Information Table 5, Supporting Information File 2a). Tannic acid ( $100 \mu\text{g mL}^{-1}$ ) slightly increased relative abundance of aminopeptidase and glycoprotein endopeptidase by +1.4 and +1.3 fold, while elongation factor G and manganese-dependent inorganic pyrophosphatase decreased by  $-2.0$  and  $-1.5$  fold, respectively (Figure 4A, Table 2, Supporting Information Table 6, Supporting Information File 2b). Caffeic acid ( $500 \mu\text{g mL}^{-1}$ ) affected nine surface protein spots, of which Glutamate tRNA ligase increased +1.5 fold, whereas eight decreased in abundance from  $-1.5$  to  $-2.2$  fold (*D*-lactate dehydrogenase, elongation factor Tu, triosephosphate isomerase, 30S ribosomal protein S1, adenylosuccinate synthetase, lysine tRNA ligase, elongation fac-

tor P, aspartate tRNA ligase (Figure 4B, Table 2, Supporting Information Table 7, Supporting Information File 2c). Ferulic acid ( $500 \mu\text{g mL}^{-1}$ ) only caused seven protein spots to reduce in abundance including L-lactate dehydrogenase ( $-1.6$  fold), oligoribonuclease ( $-1.8$  fold), and pyruvate kinase ( $-1.6$  fold) (Figure 4B, Table 2, Supporting Information Table 8, Supporting Information File 2d).

#### 4. Discussion

The daily intake of polyphenols varies a lot, typically ranging from  $<100$  mg to  $>2$  g. Over 95% of the phenolics in the diet supposedly reach the colon and become metabolized by the gut microbiota.<sup>[26]</sup> Although phenolics are generally known to exert positive human health effects, their efficacy depends on the bioavailability of different forms—esters, glycosides, polymers—which must be hydrolyzed by intestinal enzymes or the

**Table 1.** Protein identifications of differentially abundant spots (ANOVA  $p \leq 0.05$ ) of whole-cell proteins of *L. acidophilus* NCFM grown in the presence of resveratrol ( $100 \mu\text{g mL}^{-1}$ ), tannic acid ( $100 \mu\text{g mL}^{-1}$ ), caffeic acid ( $500 \mu\text{g mL}^{-1}$ ), and ferulic acid ( $500 \mu\text{g mL}^{-1}$ ) compared to non-treated control. Protein identifications were confirmed with MASCOT score of 80 for peptide mass fingerprint, ANOVA  $p \leq 0.05$ , and a minimum of 6 matched peptides.

Protein name	Gene	Fold change in resveratrol (spot no.)	Fold change in tannic acid (spot no.)	Fold change in caffeic acid (spot no.)	Fold change in ferulic acid (spot no.)
L-lactate dehydrogenase	<i>lba0271</i>	+2.0 (R2)			
Preprotein translocase subunit SecA	<i>lba0673</i>	+1.7 (R3)			
Ribonucleoside triphosphate reductase	<i>lba0041</i>	+1.6 (R4)			
Multiple sugar-binding ABC-transporter ATPase	<i>lba1645</i>	+1.5 (R5)			
GAPDH	<i>lba0698</i>	-1.5 (R6)	+1.5 (T6-T7)	-2.3 (C1)	
TsaD	<i>lba0390</i>	+1.4 (R7-R8)			
Oxalyl-CoA decarboxylase	<i>lba0396</i>	+1.4 (R9)			
Ribokinase	<i>lba0587</i>	+1.4 (R10)			
Molecular chaperone Hsp33	<i>lba0279</i>	+1.4 (R11)		-1.3 (C5)	
Alanyl-tRNA synthetase	<i>lba0417</i>	+1.3 (R12)			
tRNA (guanine-N(7)-)-methyltransferase	<i>lba1582</i>		-3.0 (T1)		
DNA-binding response regulator	<i>lba1820</i>		+2.1 (T2)		
tRNA-specific 2-thiouridylase MnmA	<i>lba0822</i>		+1.7 (T3)		
Prolyl-tRNA synthetase	<i>lba1262</i>		+1.6 (T4)		
Pyruvate kinase	<i>lba0957</i>		+1.6 (T5)		
Glutamyl-tRNA synthetase	<i>lba0347</i>		+1.5 (T8)		
Oligoendopeptidase F	<i>lba1763</i>		+1.5 (T9)		
Adenylosuccinate lyase	<i>lba1891</i>		+1.4 (T10)		
Phosphoglycerate kinase	<i>lba0699</i>		+1.4 (T11)		
50S ribosomal protein L1	<i>lba0360</i>			-1.5 (C2)	-1.7 (F2)
D-Lactate dehydrogenase	<i>lba0055</i>			+1.4 (C3)	
Elongation factor Ts	<i>lba1269</i>			+1.3 (C4)	
Purine <i>trans</i> -deoxyribosylase	<i>lba0145</i>				-1.4 (F1)
Transcriptional regulator LBA0733	<i>lba0733</i>				+1.9 (F3)
Fructokinase	<i>lba0016</i>				-1.4 (F4)
Ribose-p-pyrophosphokinase	<i>lba0224</i>				-1.6 (F5)
Phosphomethylpyrimidine kinase	<i>lba1879</i>				-1.4 (F6)


microbiota prior to absorption.<sup>[27]</sup> Phenolics and probiotic bacteria have been shown to interact with each other when coexisting in food products, dietary supplements, or in the gastrointestinal tract.<sup>[28]</sup>

The presence of plant phenolics did not inhibit the growth of NCFM cells (Supporting Information Figure 1). However, tannic acid had some growth-reducing effects on NCFM. In general, to survive in tannic acid, organisms express tannase but this is not the case of NCFM that did not display tannase activity. On the other hand, NCFM cells can protect themselves by producing exopolysaccharides, which can help the cells growing in the presence of tannic acid.<sup>[29,30]</sup> This is probably the reason why the growth is decreased but not fully inhibited.

Adhesion of microorganisms to the intestinal mucosa is important for GIT residence time and correlated to the ability of the strains to beneficially influence host health including immune modulation and competitive exclusion of pathogens.<sup>[27,28,31,32]</sup> Favorable effects of polyphenols were reported for the probiotic *Lactobacillus rhamnosus* 299 enhancing proliferation and adhesion simultaneously with inhibition of growth and adhesion of

pathogens.<sup>[3]</sup> The impact largely depended on the phenolic structure and its dosage,<sup>[34]</sup> as also found in our study to represent important factors in conferring beneficial effects.

To gain deeper insight into modulation of epithelial adhesion and changes in relative protein abundancies in the bacterial whole cell and surface-associated proteomes, four phenolics were individually added to NCFM growing with glucose as carbon source (Supporting Information Figure 1), while salicin was used as carbon source. The mechanism of bacterial adhesion to the gastrointestinal mucosa is complex and includes non-specific electrostatic and hydrophobic interactions along with specific phenomena sustained at the molecular level by bacterial adhesins and mucosal receptors.<sup>[35,36]</sup> Indeed, the most important bacterial determinants for mucosa adherence are cell-wall components and adhesins. Both specific and nonspecific mechanisms apply to the interaction of NCFM with the intestinal mucosa involving molecules of different nature, including proteins, lipids, and carbohydrates.<sup>[15]</sup> Whole-cell proteome comparison indicated that plant polyphenols can alter abundance of NCFM proteins involved in energy metabolism, general and



**Figure 3.** Representative whole-cell 2D proteome map of *L. acidophilus* NCFM grown with glucose as carbon source in the presence of caffeic acid or ferulic acid. Numbers indicate differentially abundant spots (ANOVA  $p \leq 0.05$ ), compared to control (non-treated), identified by in-gel digestion and mass spectrometry. NCFM was grown in the presence of  $500 \mu\text{g mL}^{-1}$  caffeic acid (CA-treated) and  $500 \mu\text{g mL}^{-1}$  ferulic acid (FA-treated). Selected spots are shown for relative abundance.


oxidative stress responses, transcription and translation processes, as also observed in previous proteomics studies on phenolics and LAB.<sup>[37–40]</sup>

Preprotein translocase subunit SecA is coupled with the SecYEG channel to transport polypeptides from the inside to the outside of the cell playing a central role in bacterial protein secretion.<sup>[41]</sup> The mechanism by which SecA cooperates in this function includes two phases: i) SecA (alone or together with chaperones) aids the targeting of extracellular-located proteins from the ribosome to the membrane and then ii) uses ATP for translocation of the preproteins through the SecYEG channel.<sup>[41]</sup> Increased abundance of SecA induced by resveratrol may be a first step in the adjustment of NCFM to environmental changes as extracellular proteins are crucial for this adaptation.<sup>[42]</sup>

tRNA N6-adenosine(37)-threonylcarbamoyl transferase complex transferase subunit TsaD (TsaD), also annotated as endopeptidase<sup>[43]</sup> and DNA-binding/iron metalloprotein/AP endonuclease (NCBI Reference Sequence: YP\_193312.1) is univer-

sally occurring. Its exact function is unknown, but it is thought to participate in the modification of adenosine in tRNAs reading codons beginning with adenine,<sup>[44]</sup> as well as in the modification of cell-wall peptidoglycan connected with cell division.<sup>[45]</sup> Resveratrol is known to reduce negative effects of oxidative stress on DNA and RNA and also interacts with tRNA synthetase<sup>[46]</sup> and tRNAs<sup>[47]</sup> and the increased abundance of TsaD may protect RNA against oxidative stress. Secondly, cell-wall stability may be maintained during environmental changes through increased abundance of TsaD.

Oxalyl-CoA decarboxylase together with formyl-CoA transferase is responsible for catabolism of oxalate, a toxic compound in normal human diet.<sup>[48]</sup> Oxalate is a strong chelator of cations, especially  $\text{Ca}^{2+}$ , and can lead to severe pathologies such as hyperoxaluria, urolithiasis, and renal failure. Oxalate is primarily absorbed in the colon and the gut microbiota has evolved to degrade it.<sup>[48]</sup> Even though the main organism for this degradation is *Oxalobacter formigenes*, also probiotic bacteria belonging to *Bifidobacteria* and *Lactobacilli* genera, encode genes responsible for



**Figure 4.** Surface proteomes of *L. acidophilus* NCFM grown with glucose as carbon source in the presence of phenolics. Numbers indicate differentially abundant spots (ANOVA  $p \leq 0.05$ ), compared to control (non-treated), which were identified by in-gel digestion and mass spectrometry. NCFM was grown in the presence of a)  $100 \mu\text{g mL}^{-1}$  resveratrol (Res-treated) and tannic acid (TA-treated) or b)  $500 \mu\text{g mL}^{-1}$  ferulic acid (FA-treated) and caffeic acid (CA-treated).

removal of oxalate.<sup>[49,50]</sup> NCFM is known for this activity<sup>[48]</sup> and the increase in oxalyl-CoA decarboxylase concurs with the health-promoting potential of resveratrol.

DNA-binding response regulator is a component of bacterial two-component signal transduction pathway involved in sensing and responding to environmental changes.<sup>[51]</sup> It is coupled with histidine kinase, a transmembrane protein responsible for recognition of signals from the extracellular environment by autophosphorylation. This phosphoryl group is transferred to the response regulator for activation, which once phosphory-

lated initiates differential gene transcription to trigger metabolic reactions.<sup>[51]</sup> NCFM responds to tannic acid by increasing abundance of this protein.

The glycolytic enzymes pyruvate kinase (PK) and GAPDH both increased in tannic acid-treated NCFM, indicating elevated ATP production necessary for dealing with environmental stress, as observed in other phenolic-LAB interactions.<sup>[38,40,52]</sup> By contrast, abundance of GAPDH was lower when resveratrol or caffeic acid were present during the NCFM culture. As it has been hypothesized that multiple GAPDH forms have different functions,<sup>[53]</sup>



**Table 2.** Protein identifications of differentially abundant spots (ANOVA  $p \leq 0.05$ ) of surface proteins of *L. acidophilus* NCFM treated with resveratrol (100  $\mu\text{g mL}^{-1}$ ), tannic acid (100  $\mu\text{g mL}^{-1}$ ), caffeic acid (500  $\mu\text{g mL}^{-1}$ ), or ferulic acid (500  $\mu\text{g mL}^{-1}$ ), compared to non-treated control. Protein identifications were confirmed with MASCOT score of 80 for peptide mass fingerprint, ANOVA  $p \leq 0.05$ , and a minimum of 6 matched peptides.

Protein name	Gene	Fold change in resveratrol (spot no.)	Fold change in tannic acid (spot no.)	Fold change in caffeic acid (spot no.)	Fold change in ferulic acid (spot no.)
Pyruvate kinase	<i>lba0957</i>	+2.1 (RS1)			-1.6 (FS4)
50S ribosomal protein L7/L12	<i>lba0370</i>	+2.0 (RS2)			
GAPDH	<i>lba0698</i>	-2.0 (RS3), -1.6 (RS6)			
Elongation factor P	<i>lba1668</i>	+1.7 (RS4)		-1.7 (CS8)	
50S ribosomal protein L22	<i>lba0296</i>	+1.6 (RS5)			
Adenylosuccinate synthetase	<i>lba1892</i>	-1.5 (RS7)		-1.5 (CS6)	
Hypothetical protein LBA1769	<i>lba1769</i>	+1.4 (RS8)			
6-phosphofructokinase	<i>lba0956</i>	-1.4 (RS9)			
Elongation factor G	<i>lba0289</i>		-2.0 (TS1)		
Mn-dependent inorganic pyrophosphatase	<i>lba1125</i>		-1.5 (TS2)		-1.5 (FS2)
Aminopeptidase	<i>lba1849</i>		+1.4 (TS3)		
Glycoprotein endopeptidase	<i>lba0388</i>		+1.3 (TS4)		
Glutamyl-tRNA synthetase	<i>lba0347</i>			+1.5 (CS1)	
D-Lactate dehydrogenase	<i>lba0055</i>			-1.5 (CS2)	
Elongation factor Tu	<i>lba0845</i>			-1.8 (CS3)	
Triosephosphate isomerase	<i>lba0700</i>			-2.0 (CS4)	-1.7 (FS5)
30S Ribosomal protein S1	<i>lba0968</i>			-1.8 (CS5)	-1.7 (FS6)
Lysine tRNA ligase	<i>lba0281</i>			-1.9 (CS7)	
Aspartate tRNA ligase	<i>lba0936</i>			-2.2 (CS9)	
L-Lactate dehydrogenase	<i>lba0271</i>				-1.6 (FS1)
Oligoribonuclease	<i>lba0415</i>				-1.8 (FS3)
Trigger factor	<i>lba0846</i>				-1.7 (FS7)

these results suggest that the individual plant phenolics influence occurrence of distinct GAPDH forms and hence different cellular functions. Different GAPDH forms of altered abundance were derived from the same gene and appeared in several spots probably reflecting various posttranslational modifications.

Elongation factor Ts during translation escorts aminoacyl tRNAs to the ribosome as it proceeds along the mRNA. Studies on LAB showed that acid stress and also tannic acid induce elongation factor proteins<sup>[37,38,54]</sup> and the increased abundance of elongation factor Ts by caffeic acid may be a defensive response. Furthermore, 50S ribosomal protein L1 plays a role in the structure and activity of the ribosome and can participate in the mechanism of stress adaptation as shown in *L. plantarum*.<sup>[55]</sup>

The chaperone Hsp33, which was +1.4 fold increased in resveratrol- and -1.3 fold decreased in caffeic acid-treated bacteria, deals with misfolded proteins and provides an immediate response to oxidative stress. Under oxidative conditions, Hsp33 is activated by disulfide bond formation, while under non-stressed conditions, it is deactivated by elimination of disulfide bonds with reversal of conformational changes.<sup>[56]</sup>

Remarkably, proteins changing in abundance differ for resveratrol and tannic acid, suggesting that these polyphenols have different roles in molecular reactions eliciting beneficial effects and thus act in a cooperative manner.

Regarding the surface protein profiles, some interesting observations emerge. Actually, several moonlighting or putative moon-

lighting proteins known to play a role in adhesion<sup>[57]</sup> undergo abundance changes in phenolics-stimulated NCFM. Surface proteomes after resveratrol treatment revealed increase in PK that may be responsible for adhesion, being identified as a moonlighting protein in adhesion to mucin, HT-29 cells, and yeast mannan.<sup>[58,59]</sup> Other higher-abundant proteins, such as elongation factor P (EF-P) and ribosomal proteins are putative moonlighting proteins previously found on bacterial surfaces.<sup>[59]</sup> Elongation factors besides participating in protein synthesis are often described as moonlighting, especially elongation factor Tu (EF-Tu) that promoted adhesion of *L. johnsonii* to human tissues,<sup>[60]</sup> thus contributing to its health-promoting effect. Similarly, EF-Ts and EF-G, seen in the exoproteome of *Bacillus anthracis*<sup>[61]</sup> were referred to as signal peptide-lacking exoproteins in *Staphylococcus aureus*.<sup>[62]</sup> The very interesting protein trigger factor behaves in *Lactobacillus reuteri* NCIB11951 as a collagen I binding protein<sup>[63]</sup> and cooperates in *Streptococcus mutans* with surface adhesin P1.<sup>[64]</sup> Notably NCFM EF-P is more abundant after resveratrol treatment, stimulating adhesion to both mucin and HT-29 cells, and less abundant in caffeic acid-treated NCFM not showing improved adhesion. It is tempting to hypothesize that different plant phenolics influence synthesis of EF-P in NCFM in different ways or regulate secretion of EF-P to the outside of the cell, thus controlling adhesion to host and tissue components such as mucin and collagen. The lack of effect of caffeic acid on adhesion at 500  $\mu\text{g mL}^{-1}$  agrees with low abundance of EF-Tu.<sup>[60]</sup>

Remarkably, improved adhesion by resveratrol treatment does not fit with the well-known moonlighting adhesive protein GAPDH being less abundant.<sup>[60]</sup> Probably several proteins including EF-P and PK support the adhesive effect.

Apart from the EF-P and EF-Tu discussed above connected with the good (resveratrol) or poor (caffeic acid) adhesive capabilities of stimulated NCFM, it can be underlined that other elongation (EF-G) and trigger factors are of low abundance in the surface proteome of tannic and ferulic acid-stimulated bacteria, respectively. In the case of ferulic acid (500  $\mu\text{g mL}^{-1}$ ), reduced abundance of PK is also consistent with poor adhesion to HT-29 at this concentration. By contrast, improved binding to HT-29 resulting for tannic acid, probably is due to other components, maybe exopolysaccharides often reported as involved in cell adhesion.<sup>[65]</sup> This hypothesis is supported by the fact that cells stimulated with tannic acid did not adhere to mucin and possibly factors stimulating adhesion to HT-29 are unfavorable for binding to mucin.

Finally, it is worth considering that both ferulic acid and resveratrol seem to preferentially influence L-lactate dehydrogenase (L-LDH) synthesis, whereas caffeic acid seems to specifically control D-lactate dehydrogenase (D-LDH) abundance. Curiously, caffeic acid causes enhancement of intracellular D-LDH and decreases abundance of surface D-LDH, suggesting that it directs the cellular location of this enzyme rather than stimulating the synthesis.

In conclusion, the present study has brought new knowledge to elucidate complex interactions occurring in the human gut between health-promoting bacteria and diet components. It has been demonstrated that some plant phenolics (but not all) can improve the adhesive capabilities to mucin and HT-29 cells of NCFM probably by inducing biosynthesis or secretion of moonlighting proteins engaged in adhesion. Among these compounds, resveratrol proved most effective.

## Abbreviations

5(6)-CFDA, 5(6)-carboxyfluorescein diacetate; GAPDH, glyceraldehyde-3-p dehydrogenase; GIT, gastrointestinal tract; LABSEM, semisynthetic lactic acid bacteria medium; NCFM, *Lactobacillus acidophilus* NCFM

## Supporting Information

Supporting Information is available from the Wiley Online Library or from the author.

## Acknowledgements

Karina Jensen, Anne Blicher, and Lisbeth Buus Rosholm are thanked for technical assistance. The authors are grateful to Prof. Ayse Mine Gencler-Ozkan, Ankara University, for very stimulating discussions about plants and plant phenolics. This work was supported by the Danish Strategic Research Council's Program Committee on Health, Food and Welfare (FøSu), the Danish Council for Independent Research | Natural Sciences and the Danish Center for Advanced Food Studies (LMC). H.U.C. is grateful to the Republic of Turkey, Ministry of National Education for a Ph.D. scholarship. M.D. benefitted from an Erasmus agreement to perform her master project at DTU.H.U.C., S.B., E.P., and B.S. conceived and designed the work. H.U.C. and M.D. performed experimental work, data analysis. and

interpretation and drafted the paper. S.B., E.P., and B.S. helped writing and revising the paper. All authors read and approved the final version.

## Conflict of Interest

The authors have declared no conflict of interest.

## Keywords

ferulic acid, probiotics, resveratrol, surface proteome, whole-cell proteome

Received: July 24, 2017  
Revised: November 20, 2017

- [1] K. B. Pandey, S. I. Rizvi, *Oxid. Med. Cell. Longev.* **2009**, 2, 270.
- [2] C. H. Beckman, *Physiol. Mol. Plant Pathol.* **2000**, 57, 101.
- [3] J. A. Baur, D. A. Sinclair, *Nat. Rev. Drug Discov.* **2006**, 5, 493.
- [4] J. Serrano, R. Puupponen-Pimia, A. Dauer, A. M. Aura, F. Saura-Calixto, *Mol. Nutr. Food Res.* **2009**, 53, 310.
- [5] T. Silva, C. Oliveira, F. Borges, *Expert Opin. Ther. Pat.* **2014**, 24, 1257.
- [6] R. Upadhyay, L. J. Mohan Rao, *Crit. Rev. Food Sci. Nutr.* **2013**, 53, 968.
- [7] N. Kumar, V. Pruthi, *Biotechnol. Rep.* **2014**, 4, 86.
- [8] L. B. de Paiva, R. Goldbeck, W. D. dos Santos, F. M. Squina, *Brazilian J. Pharm. Sci.* **2013**, 49, 395.
- [9] Z. Zhao, M. H. Moghadasian, *Food Chem.* **2008**, 109, 691.
- [10] B. Schmid, I. Kötter, L. Heide, L. Heide, *Eur. J. Clin. Pharmacol.* **2001**, 57, 387.
- [11] M. E. Sanders, T. R. Klaenhammer, *J. Dairy Sci.* **2001**, 84, 319.
- [12] S. E. Gilliland, M. L. Speck, C. G. Morgan, *Appl. Microbiol.* **1975**, 30, 541.
- [13] R. Barrangou, M. A. Azcarate-Peril, T. Duong, S. B. Connors, R. M. Kelly, T. R. Klaenhammer, *Proc. Natl. Acad. Sci. U. S. A.* **2006**, 103, 3816.
- [14] B. Johnson, K. Selle, S. O'Flaherty, Y. J. Goh, T. Klaenhammer, *Microbiology* **2013**, 159, 2269.
- [15] B. Buck, E. Altermann, T. Svingerud, T. R. Klaenhammer, *Appl. Environ. Microbiol.* **2005**, 71, 8344.
- [16] K. J. Heller, *Am. J. Clin. Nutr.* **2001**, 73, 374S.
- [17] A. Majumder, A. Sultan, R. R. Jersie-Christensen, M. Ejby, B. G. Schmidt, S. J. Lahtinen, S. Jacobsen, B. Svensson, *Proteomics* **2011**, 11, 3470.
- [18] E. Izquierdo, P. Horvatovich, E. Marchioni, D. Aoude-Werner, Y. Sanz, S. Ennahar, *Electrophoresis* **2009**, 30, 949.
- [19] H. U. Celebioglu, M. Ejby, A. Majumder, C. Købler, Y. J. Goh, K. Thorsen, B. Schmidt, S. O'Flaherty, M. Abou Hachem, S. J. Lahtinen, S. Jacobsen, T. R. Klaenhammer, S. Brix, K. Mølhave, B. Svensson, *Proteomics* **2016**, 16, 1361.
- [20] H. U. Celebioglu, S. V. Olesen, K. Prehn, S. J. Lahtinen, S. Brix, M. Abou Hachem, B. Svensson, *J. Proteomics* **2017**, 163, 102.
- [21] S. Lortal, J. V. Heijenoort, K. Gruber, U. B. Sleytr, *J. Gen. Microbiol.* **1992**, 138, 611.
- [22] F. Genovese, J. D. Coisson, A. Majumder, A. Pessione, B. Svensson, S. Jacobsen, E. Pessione, *Food Res. Int.* **2013**, 54, 1072.
- [23] N. A. Karp, J. L. Griffin, K. S. Lilley, *Proteomics* **2005**, 5, 81.
- [24] G. Candiano, M. Bruschi, L. Musante, L. Santucci, G. M. Ghiggeri, B. Carnemolla, P. Orecchia, L. Zardi, P. G. Righetti, *Electrophoresis* **2004**, 25, 1327.
- [25] R. I. Dave, N. P. Shah, *J. Dairy Sci.* **1996**, 79, 1529.
- [26] M. N. Clifford, *Planta Med.* **2004**, 70, 1103.
- [27] L. Marín, E. M. Miguélez, C. J. Villar, F. Lombó, *Biomed Res. Int.* **2015**, 2015, 18.

- [28] A. Duda-Chodak, T. Tarko, M. Statek, *Acta Sci. Pol., Technol. Aliment.* **2008**, *7*, 39.
- [29] A. H. Smith, E. Zoetendal, R. I. Mackie, *Microb. Ecol.* **2005**, *50*, 197.
- [30] L. O'Donovan, J. D. Brooker, *Microbiology* **2001**, *147*, 1025.
- [31] I. González-Rodríguez, L. Ruiz, M. Gueimonde, A. Margolles, B. Sánchez, *FEMS Microbiol. Lett.* **2013**, *340*, 1.
- [32] M. L. Van Tassel, M. J. Miller, *Nutrients* **2011**, *3*, 613.
- [33] S. G. Parkar, D. E. Stevenson, M. A. Skinner, *Int. J. Food Microbiol.* **2008**, *124*, 295.
- [34] I. Bustos, T. García-Cayuela, B. Hernández-Ledesma, C. Peláez, T. Requena, M. C. Martínez-Cuesta, *J. Agric. Food Chem.* **2012**, *60*, 9082.
- [35] C. N. Jacobsen, V. R. Nielsen, A. E. Hayford, P. L. Moller, K. F. Michaelsen, A. Pærregaard, B. Sandström, M. Tvede, M. Jakobsen, *Appl. Environ. Microbiol.* **1999**, *65*, 4949.
- [36] M. Gueimonde, A. Margolles, C. G. de los Reyes-Gavilán, S. Salminen, *Int. J. Food Microbiol.* **2007**, *113*, 228.
- [37] J. A. Curiel, H. Rodríguez, B. de Las Rivas, P. Anglade, F. Baraige, M. Zagorec, M. Champomier-Vergès, R. Muñoz, F. López de Felipe, *Mol. Nutr. Food Res.* **2011**, *55*, 1454.
- [38] D. Cecconi, M. Cristofolletti, A. Milli, P. Antonioli, S. Rinalducci, L. Zolla, G. Zapparoli, *Electrophoresis* **2009**, *30*, 957.
- [39] A. Bossi, S. Rinalducci, L. Zolla, P. Antonioli, P. G. Righetti, G. Zapparoli, *J. Appl. Microbiol.* **2007**, *102*, 787.
- [40] M. F. Mazzeo, R. Lippolis, A. Sorrentino, S. Liberti, F. Fragnito, R. A. Siciliano, *PLoS One* **2015**, *10*, e0142376.
- [41] K. E. Chatzi, M. F. Sardis, A. Economou, S. Karamanou, *Biochim. Biophys. Acta. Mol. Cell Res.* **2014**, *1843*, 1466.
- [42] M. Zhou, D. Theunissen, M. Wels, R. J. Siezen, *BMC Genomics* **2010**, *11*, 651.
- [43] Ma. Azcarate-Peril, R. Tallon, T. R. Klaenhammer, *J. Dairy Sci.* **2009**, *92*, 870.
- [44] L. Perrochia, E. Crozat, A. Hecker, W. Zhang, J. Bareille, B. Collinet, H. van Tilbeurgh, P. Forterre, T. Basta, *Nucleic Acids Res.* **2013**, *41*, 1953.
- [45] L. Zheng, C. Yu, K. Bayles, I. Lasa, Y. Ji, *J. Bacteriol.* **2007**, *189*, 2734.
- [46] M. Sajish, P. Schimmel, *Nature* **2015**, *519*, 370.
- [47] C. N. N'soukpoé-Kossi, P. Bourassa, J. S. Mandeville, L. Bekale, J. Bariyanga, H. A. Tajmir-Riahi, *Int. J. Biol. Macromol.* **2015**, *80*, 41.
- [48] S. Turroni, B. Vitali, C. Bendazzoli, M. Candela, R. Gotti, F. Federici, F. Pirovano, P. Brigidi, *J. Appl. Microbiol.* **2007**, *103*, 1600.
- [49] F. Federici, B. Vitali, R. Gotti, R. Pasca, S. Gobbi, A. B. Peck, P. Brigidi, *Appl. Environ. Microbiol.* **2004**, *70*, 5066.
- [50] E. Altermann, W. M. Russell, M. A. Azcarate-Peril, R. Barrangou, B. L. Buck, O. McAuliffe, N. Souther, A. Dobson, T. Duong, M. Callanan, S. Lick, A. Hamrick, R. Cano, T. R. Klaenhammer, *Proc. Natl. Acad. Sci. U. S. A.* **2005**, *102*, 3906.
- [51] J. M. Skerker, M. S. Prasol, B. S. Perchuk, E. G. Biondi, M. T. Laub, *PLoS Biol.* **2005**, *3*, e334.
- [52] I. Reveron, H. Rodriguez, G. Campos, J. A. Curiel, C. Ascaso, A. V. Carrascosa, A. Prieto, B. de las Rivas, R. Munoz, F. L. de Felipe, *PLoS One* **2013**, *8*, e66473.
- [53] C. Tristan, N. Shahani, T. W. Sedlak, A. Sawa, *Cell. Signal.* **2011**, *23*, 317.
- [54] K. Lee, H. G. Lee, K. Pi, Y. J. Choi, *Proteomics* **2008**, *8*, 1624.
- [55] M. De Angelis, R. Di Cagno, C. Huet, P. F. Fox, P. F. Fox, M. Gobbetti, *Appl. Environ. Microbiol.* **2004**, *70*, 1336.
- [56] Y. O. Ali, B. M. Kitay, R. G. Zhai, *Molecules* **2010**, *15*, 6859.
- [57] G. Bergonzelli, D. Granato, *Infect. Immun.* **2006**, *74*, 425.
- [58] Y. Katakura, R. Sano, T. Hashimoto, K. Ninomiya, S. Shioya, *Appl. Microbiol. Biotechnol.* **2010**, *86*, 319.
- [59] W. Wang, C. J. Jeffery, *Mol. Biosyst.* **2016**, *12*, 1420.
- [60] D. Granato, G. E. Bergonzelli, R. D. Pridmore, L. Marvin, M. Rouvet, I. E. Corthésy-Theulaz, *Infect. Immun.* **2004**, *72*, 2160.
- [61] M. Gohar, N. Gilois, R. Graveline, C. Garreau, V. Sanchis, D. Lereclus, *Proteomics* **2005**, *5*, 3696.
- [62] M. Sibbald, A. K. Ziebandt, S. Engelmann, M. Hecker, A. de Jong, H. J. Harmsen, G. C. Raangs, I. Stokroos, J. P. Arends, J. Y. Dubois, J. M. van Dijk, *Microbiol. Mol. Biol. Rev.* **2006**, *70*, 755.
- [63] P. Aleljung, W. Shen, B. Rozalska, U. Hellman, A. Ljungh, T. Wadström, *Curr. Microbiol.* **1994**, *28*, 231.
- [64] P. J. Crowley, T. B. Seifert, R. Isoda, M. van Tilburg, M. W. Oli, R. A. Robinette, W. P. McArthur, A. S. Bleiweis, L. J. Brady, *Infect. Immun.* **2008**, *76*, 2456.
- [65] E. Vera-Pingitore, A. Pessione, C. Fontana, R. Mazzoli, E. Pessione, *Int. J. Food Micro.* **2016**, *238*, 96.