

*IFSCOM 2016*

## **IFSCOM 2016**

**29 August - 01 September 2016**

**MERSİN UNIVERSITY  
MERSİN  
TURKEY**

**Proceeding Book**

**Website: <http://iifsc.com>**

Thanks to Mersin University for support.

*IFSCOM 2016*

ISBN: 978-975-6900-54-3

## Preface

We are very pleased to introduce the abstracts of the 3rd International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference (IFSCOM2016).

As for previous conferences, the theme was the link between the Mathematics by many valued logics and its applications.

In this context, there is a need to discuss the relationships and interactions between many valued logics and contemporary mathematics.

Finally, in the previous conference, it made successful activities to communicate with scientists working in similar fields and relations between the different disciplines.

This conference has papers in different areas; multi-valued logic, geometry, algebra, applied mathematics, theory of fuzzy sets, intuitionistic fuzzy set theory, mathematical physics, mathematics applications, etc

Thank you to the scientists offering the most significant contribution to this conference.

Thank you to the Scientific Committee Members, Referee Committee Members, Local Committee Members, University Administrators and Mersin University Mathematics Department Scientists.

Organizing Committee  
Asist. Prof. GÖKHAN ÇUVALCIOĞLU  
Co-chairman

Organizing Committee  
Asist. Prof. SAÏD MELLIANI  
Co-chairman

## **Invited Speakers**

**ÖMER AKIN**

TOBB Economy and Technology University  
TR

**OSCAR CASTILLO**

Tijuana Institute of Technology  
MX

**ABDULLAH ÇAVUŞ**

Karadeniz Technical University  
TR

## Scientific Committee

- OSMAN ABUL, Turkey  
ÇİĞDEM GÜNDÜZ ARAS, Turkey  
ISMAT BEG, Pakistan  
HUMBERTO BUSTINCE, Spain  
OSCAR CASTILLO, Mexico  
AYNUR ÇÖL, Turkey  
VİLDAN ÇETKİN, Turkey  
MEHMET ÇİTİL, Turkey  
BIJAN DAVVAZ, Iran  
UĞUR DEĞER, Turkey  
ARINDAM GARAI, India  
ALİ KEMAL HAVARE, Turkey  
JANUSZ KACPRZYK, Poland  
ERDENER KAYA, Turkey  
TAEKYUN KIM, Korea  
TUFAN KUMBASAR, Turkey  
POOM KUMAM, Thailand  
HAMZA MENKEN, Turkey  
JUAN MARTÍNEZ MORENO, Spain  
PIOTR NOWAK, Poland  
MADHUMANGAL PAL, India  
THEMISTOCLES M. RASSIAS, Greece  
SALIM REZVANI, Iran  
EKREM SAVAŞ, Turkey  
MAURİCE SALLES, France  
CALOGERO VETRO, Italy  
ZESHUİ XU, China  
EROL YAŞAR, Turkey  
CALOGERO VETRO, Italy  
HAKAN YAZICI, Turkey
- ÖMER AKIN, Turkey  
VASSIA K. ATANASSOVA, Bulgaria  
ADRIAN BAN, Romania  
GALİP CANSEVER, Turkey  
LALLA SAADIA CHADLI, Morocco  
ORKUN COŞKUNTUNCEL, Turkey  
ALİ ÇEVİK, Turkey  
İLHAN DAĞADUR, Turkey  
ÜMİT DENİZ, Turkey  
ESFANDIAR ESLAMI, Iran  
HENRI GWÉT, Cameroon  
BIPAN HAZARIKA, India  
EKREM KADIOĞLU, Turkey  
NAZIM KERİMOV, Turkey  
ŞÜKRAN KONCA, Turkey  
MEHMET KÜÇÜKASLAN, Turkey  
PATRICIA MELIN, Mexico  
ÖZGÜR MIZRAK, Turkey  
EFENDİ NASİBOĞLU, Turkey  
TAHA YASİN ÖZTÜRK, Turkey  
DAOWU PEI, China  
HANLAR REŞİDOĞLU, Turkey  
REZA SAADATI, Iran  
RIDVAN ŞAHİN, Turkey  
SOTIR SOTIROV, Bulgaria  
YEJUN XU, China  
RONALD R. YAGER, USA  
MURAT İBRAHİM YAZAR, Turkey  
ADNAN YAZICI, Turkey

## Referees

AZAMAT AKHTYAMOV, Russia

ÖMER AKIN, Turkey

KRASSIMIR T. ATANASSOV, Bulgaria

OSCAR CASTILLO, Mexico

GÖKHAN ÇUVALCIOĞLU, Turkey

JANUSZ KACPRZYK, Poland

EKREM KADIOĞLU, Turkey

MEHMET KÜÇÜKASLAN, Turkey

SAID MELLIANI, Morocco

HANLAR REŞİDOLU, Turkey

EKREM SAVAŞ, Turkey

SOTIR SOTIROV, Bulgaria

## Local Organizing Committee

HANLAR REŞİDOĞLU  
MEHMET KÜÇÜKASLAN  
NAZIM KERİMOV  
FAHREDDİN ABDULLAYEV  
İLHAN DAĞADUR  
HAMZA MENKEN  
EROL YAŞAR  
ALİ ÇEVİK  
GÖKHAN ÇUVALCIOĞLU  
UĞUR DEĞER  
ÖZGÜR MIZRAK  
TUNCAY TUNÇ  
İHSAN SEFA ÇETİN  
FATMA AYÇA ÇETİNKAYA  
VOLKAN ALA  
MAYA ALTINOK<sup>1</sup>  
ÖZGE ÇOLAKOĞLU HAVARE  
BURÇİN DOĞAN  
SERTAÇ GÖKTAŞ  
ŞEYDA SEZGEK  
SİBEL YASEMİN GÖLBOL  
İLKNUR YEŞİLCE  
SİNEM YILMAZ  
HALİL İBRAHİM YOLDAŞ  
ARİF BAL  
FATİH KUTLU

---

<sup>1</sup>Proceeding book designed by Maya ALTINOK

## CONTENTS

PREFACE .....	i
INVITED SPEAKERS .....	ii
SCIENTIFIC COMMITTEE .....	iii
REFEREES .....	iv
LOCAL ORGANIZING COMMITTEE .....	v
SECTION I (Fulltexts) .....	x
Seda Kılınc, Sümeyye Ermeydan, Hüseyin Yıldırım	
New Integral Inequalities Involving Beta Function Via $P_\varphi$ -Preinvex Convexity .....	1-9
Ferit Gurbuz	
Boundedness Of The Sublinear Operators With Rough Kernel Generated By Fractional Integrals And Their Commutators On Generalized Vanishing Morrey Spaces II .....	10-36
Adem Başkaya, Orkun Coşkuntuncel	
Teachers' Opinions About Mathematic Program Revised With 4+4+4 Education System.....	37-41
Khanlar R. Mamedov, Döne Karahan	
The Direct And Inverse Spectral Problem For Sturm-Liouville Operator With Discontinuous Coefficient.....	42-53
Hamiyet Merkezci, Necati Olgun, Ela Aydın	
On Second Order Symmetric Derivations of Kähler Modules and It's Projective Dimension for Hypersurfaces .....	54-57
Olgun Cabri, Khanlar R. Mamedov	
On A Nonlocal Boundary Value Problem.....	58-64
Mehmet Cital, Muhammed Gezeran	
Fuzzy Right Fractional Ostrowski Inequalities .....	65-72
Gökhan Çuvalcıoğlu	
New Intuitionistic Fuzzy Level Sets .....	73-77
Özgür Mızrak, Kenan Sogut	
Energy Spectrum of Spinless Particles in Electromagnetic Fields.....	78-83
Sinem Tarsuslu(Yılmaz), Gökhan Çuvalcıoğlu, Arif Bal	
Some Intuitionistic Fuzzy Modal Operators Over Intuitionistic Fuzzy Ideals and Groups .....	84-90
Mehmet Emre Erdogan, Kemal Uslu	
Stability Analysis on Effect of System Restore on Epidemic Model for Computer Viruses .....	91-104
Ferit Gurbuz	
Boundedness of the Sublinear Operators with Rough Kernel Generated by Calderón-Zygmund Operators and Their Commutators on Generalized Vanishing Morrey Spaces.....	105-129
Fatih Kutlu, Özkan Atan, Tunay Bilgin	
Distance Measure, Similarity Measure, Entropy and Inclusion Measure for Temporal Intuitionistic Fuzzy Sets .....	130-148
Olgun Cabri, Khanlar R. Mamedov	
On A Social Economic Model.....	149-154
Gökhan Çuvalcıoğlu, Krassimir T. Atanassov, Sinem Tarsuslu(Yılmaz)	
Some Results on $S_{\alpha,\beta}$ and $T_{\alpha,\beta}$ Intuitionistic Fuzzy Modal Operators .....	155-161
Sümeyye Ermeydan, Hüseyin Yıldırım	
On Hermite-Hadamard Inequalities for Geometric-Arithmetically $\varphi$ -s-Convex Functions Via Fractional Integrals .....	162-174
F. Ayca Cetinkaya, Khanlar R. Mamedov, Gizem Cerci	
On A Boundary Value Problem With Retarded Argument In The Differential Equation.....	175-178
Mehmet Cital, Sinem Tarsuslu(Yılmaz)	
$\Omega$ -Algebras on co-Heyting Valued Sets.....	179-184
Şeyda Sezgek, İlhan Dağadur	
Some Clasifications of an X FDK-Spaces .....	185-191
Maya Altınok, Mehmet Küçükaslan	
A Note on Porosity Cluster Points .....	192-195



SECTION II (Abstracts) .....	196
Mustafa Saltı, Oktay Aydoğdu	
Fracral Reconstruction of Dilaton Field .....	197
A. El Allaoui, S. Melliani	
The Cauchy Problem for Complex Intuitionistic Fuzzy Differential Equations .....	198
B. Ben Amma, L. S. Chadli	
Numerical Solution of Intuitionistic Fuzzy Differential Equations	
by Runge-Kutta Method of Order Four .....	199
R. Ettoussi, L. S. Chadli	
Fractional Differential Equations with Intuitionistic Fuzzy Data .....	200
Ömer Akın, Selami Bayeg	
Solving Second Order Intuitionistic Fuzzy Initial Value Problems with Heaviside Function .....	201
Sukran Konca, Mochammad İdris	
Equivalence Among Three 2-Norms on the Space of $\mathbf{p}$ –Summable Sequences .....	202
Murat İbrahim Yazar, Cigdem Gunduz (Aras)	
Some Properties of Soft Mappings on Soft Metric Spaces .....	203
Cigdem Gunduz (Aras), Murat İbrahim Yazar	
Soft Totally Bounded Spaces in Soft Metric Spaces .....	204
Nihal Taş, Nihal Yılmaz Özgür	
Some Generalized Fixed Point Type Theorems on an $\mathbf{S}$ –Metric Space .....	205
Nihal Taş, Nihal Yılmaz Özgür	
A New Generalization of Soft Metric Spaces .....	206
Vildan Çetkin, Halis Aygün	
Some Separation Axioms in Fuzzy Soft Topological Spaces .....	207
Gizem Temelcan, Hale Gonce Kocken, İnci Albayrak	
Fuzzy Equilibrium Analysis of a Transportation Network Problem .....	208
Nesrin Çalışkan, Ayşe Funda Yaliniz	
A Special Type of Sasakian Finsler Structures on Vector Bundles .....	209
Abdullah Çavuş	
On Some Properties and Applications of the Quasi-Resolvent Operators	
of the Infinitesimal Operator of a Strongly Continuous	
Linear Representation of the Unit Circle Group in a Complex Banach Space .....	210
B. Abdellaoui, K. Biroud, J. Davila, F. Mahmoudi	
Existence and Nonexistence for Nonlinear Problems with Singular Potential .....	211
I. Bakhadach, S. Melliani, L. S. Chadli	
Intuitionistic Fuzzy Soft Generalized Superconnected .....	211
İzzettin Demir	
Vietoris Topology in the Context of Soft Set .....	212
Burçin Doğan, Ali Yakar, Erol Yaşar	
On Totally Umbilical and Minimal Cauchy Riemannian	
Lightlike Submanifolds of an Indefinite Kaehler Manifold .....	213
Murat Bekar, Yusuf Yaylı	
Algebraic Properties of Dual Quasi-Quaternions .....	214
Ferhan Sola Erduran	
Fixed Intuitionistic Fuzzy Point Theorem in Hausdorff Intuitionistic Fuzzy Metric Spaces .....	215
Erdener Kaya	
Orthonormal Systems in Spaces of Number Theoretical Functions .....	216
Gülnur Çelik Kızıllan, Kemal Aydın	
A Step Size Strategy for Numerical Integration of the	
Hurwitz Stable Differential Equation Systems .....	217
Ahmet Duman, Ali Osman Cıbıkdiken, Gülnur Çelik Kızıllan, Kemal Aydın	
Schur Stabilitiy in Floating Point Arithmetic: Systems with Constant Coefficients .....	218

Gülnur Çelik Kızılkın, Ahmet Duman, Kemal Aydın	
The Numerical Solution of Some SIR Epidemic Models with Variable Step Size Strategy .....	220
Faruk Polat	
When is an Archimedean $\mathbf{f}$ -Algebra Finite Dimensional? .....	221
Müzeyyen Özhavzalı	
By Calculating for Some Linear Positive	
Operators to Compare of the Errors in the Approximations .....	222
Banu Pazar Varol, Abdulkadir Aygünoğlu, Halis Aygün	
Intuitionistic Fuzzy Soft Neighborhoods .....	223
Melek Erdoğan	
On Nonlightlike Off set Curves in Minkowski 3-Space .....	224
Erhan Güler, Ömer Kişi, Semra Saraçoğlu Çelik	
Weierstrass Representation, Degree and Classes of the Surfaces	
in the Four Dimensional Euclidean Space .....	225
Merve Özdemir, Nihat Akgüneş	
A New Type Graph and Their Parameters .....	226
Büşra Cağan, Nihat Akgüneş	
The Dot Product Graph of Monogenic Semigroup .....	226
Seda Öztürk	
Some Number Theoretical Results Related to the Suborbital Graphs	
for the Congruence Subgroup $\Gamma_0\left(\frac{n}{h}\right)$ .....	227
Ali Osman Çıbıkdiken, Ahmet Duman, Kemal Aydın	
Schur Stability in Floating Point Arithmetic: Systems with Periodic Coefficients .....	227
Ömer Kişi, Fatih Nuray	
I-Limit Inferior and I-Limit Superior of Sequences of Sets .....	230
Filiz Yıldız	
A Ditopological Fuzzy Structural View of Inverse Systems and Inverse Limits .....	230
Emel Aşıcı	
On Nullnorms on Bounded Lattices .....	231
Sukran Konca, Ergin Genç, Selman Ekin	
Ideal Version of Weighted Lacunary Statistical Convergence of Sequences of Order .....	233
Mahmut Karakuş	
AK (S) and AB (S) Properties of a K-Space .....	234
Melek Yağcı	
On the Second Homology of the Schützenberger Product of Monoids 39 .....	235
Ömer Kişi, Semra Saraçoğlu Çelik, Erhan Güler	
New Sequence Spaces with Respect to a Sequence of Modulus Functions.....	236
Erhan Güler, Ömer Kişi, Semra Saraçoğlu Çelik	
TF-Type Hypersurfaces in 4-Space .....	236
Özge Çolakoğlu Havare, Hamza Menken	
A Note on $\mathbf{q}$ -Binomial Coefficients .....	237
Mehmet Cihan Bozdağ, Hamza Menken	
On Statistical Convergence of Sequences of $\mathbf{p}$ -Adic Numbers .....	238
Halil İbrahim Yoldaş, Erol Yaşar	
Lightlike Hypersurface of an Indefinite Kaehler Manifold with a	
Complex Semi-Symmetric Metric Connection .....	239
M. Elomari, S. Melliani, L. S. Chadli	
Intuitionistic Fuzzy Fractional Evolution Problem .....	239
Philippe Balbiani, Çiğdem Gencer, Zafer Özdemir	
A General Tableaux Method for Contact Logics Interpreted over Intervals.....	241
Suvankar Biswas, Sanhita Banerjee, Tapan Kumar Roy	
Solving Intuitionistic Fuzzy Differential Equations with Linear	
Differential Operator by Adomain Decomposition Method .....	242
Murat Bekar	
Involution Matrices of $\frac{1}{4}$ -Quaternions .....	245
Anıl Özdemir	

Topological Full Groups of Cantor Minimal Systems .....	246
Ramazan Yazgan, Cemil Tunç	
Global Stability to Nonlinear Neutral Differential Equations of First Order .....	247
Nihat Akgüneş, Ahmet Sinan Çevik	
A Study on the Cartesian Product of a Special Graphs .....	247
Murat Bekar, Yusuf Yaylı	
Unit Dual Lorentzian Sphere and Tangent Bundle of Lorentzian Unit 2-Sphere .....	248
Volkan Ala , Khanlar R. Mamedov	
On the Inverse Problem for a Sturm-Liouville Equation with Discontinuous Coefficient .....	249
Ali Kemal Havare, Özge Çolakoğlu Havare	
Calculation and Analysis of Electronic Parameters of Electroluminescent Device Cells Through I-V Based Modeling .....	249
Mahmut Karakuş	
On Some Properties of Sum Spaces .....	250
Mahmut Karakuş	
On $q^\lambda$ and $q_0^\lambda$ Invariant Sequence Spaces .....	251
Nihat Akgüneş	
Some New Results on a Graph of Monogenic Semigroup .....	252
Fuat Usta	
Computational Solution of Katugampola Conformable Fractional Differential Equations Via RBF Collocation Method .....	253
Fuat Usta, Mehmet Zeki Sarıkaya	
Some Integral Inequalities Via Conformable Calculus .....	254
Hüseyin Budak, Fuat Usta, Mehmet Zeki Sarıkaya	
Weighted Ostrowski, Chebyshev and Gruss Type Inequalities for Conformable Fractional Integrals .....	254
Gökçe Sucu	
Bifurcation and Stability Analysis of a Discrete-Time Predator- Prey Model .....	255
Gabil Adilov, İlknur Yeşilce	
Applications of Hermite-Hadamard Inequalities for $\mathbb{B}$ -Convex Functions and $\mathbb{B}^{-1}$ -Convex Functions .....	256
Arindam Garal, Palash Mandal, Tapan Kumar Roy	
Operations and Extension Principle under T-Intuitionistic Fuzzy Environment .....	257
Ekrem Savaş	
On Generalized Double Statistical Convergence of Order in Intuitionistic Fuzzy N- Normed Spaces .....	258
Abdullah Akkurt, M. Esra Yıldırım, Hüseyin Yıldırım	
On Feng Qi-Type Integral Inequalities for Conformable Fractional Integrals .....	259

SECTION I  
FULLTEXTS

**NEW INTEGRAL INEQUALITIES INVOLVING BETA  
 FUNCTION VIA  $P_\varphi$ -PREINVEX CONVEXITY**

SEDA KILINÇ, SÜMEYYE ERMEYDAN, AND HÜSEYİN YILDIRIM

ABSTRACT. In this note, we establish some inequalities, involving the Euler Beta function, of the integral  $\int_a^b (x-a)^p (b-x)^q f(x) dx$  for functions when a power of the absolute value is  $P_\varphi$ -preinvex.

Received: 26–August–2016

Accepted: 29–August–2016

1. INTRODUCTION

[27], Wenjun Lio introduced new inequalities for  $P$ -convexity. We establish new Hermite-Hadamard inequalities for quasi-preinvex and  $P_\varphi$ -preinvex functions.

Let  $I$  be an interval in  $\mathbb{R}$ . Then  $f : I \rightarrow \mathbb{R}$  is said to be preinvex convex if

$$(1.1) \quad f(x + (1-t)e^{i\varphi}\eta(y,x)) \leq tf(x) + (1-t)f(y)$$

holds for all  $x, y \in I$  and  $t \in [0, 1]$ .

The notion of quasi-preinvex functions generalizes the notion  $P_\varphi$ -preinvex functions. More precisely, a function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be quasi-preinvex on  $[a, b]$  if,

$$(1.2) \quad f(x + (1-t)e^{i\varphi}\eta(y,x)) \leq \max\{f(x), f(y)\}$$

holds for any  $x, y \in [a, b]$  and  $t \in [0, 1]$ . Clearly, any preinvex function is a quasi-preinvex function. Furthermore, there exist quasi-preinvex functions which are not preinvex.

The generalized quadrature formula of Gauss-Jacobi type has the form

$$\int_a^b (x-a)^p (b-x)^q f(x) dx = \sum_{k=0}^m B_{m,k} f(\gamma_k) + \mathfrak{R}_m[f]$$

for certain  $B_{m,k,\gamma_k}$  and rest term  $\mathfrak{R}_m[f]$  (see [22]).

Let  $\mathbb{R}^n$  be Euclidian space and  $K$  is said to a nonempty closed in  $\mathbb{R}^n$ . Let  $f : K \rightarrow \mathbb{R}$ ,  $\varphi : K \rightarrow \mathbb{R}$  and  $\eta : K \times K \rightarrow \mathbb{R}$  be a continuous functions.

**Definition 1.1.** ([13]) Let  $u \in K$ . The set  $K$  is said to be  $\varphi$ -invex at  $u$  with respect to  $\eta$  and  $\varphi$  if

$$(1.3) \quad u + te^{i\varphi}\eta(v, u) \in K$$

for all  $u, v \in K$  and  $t \in [0, 1]$ .

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
 2010 *Mathematics Subject Classification.* 26A33, 26D15, 41A55.

*Key words and phrases.* Fractional Hermite-Hadamard ineaualities,  $\varphi$ -preinvex functions, Riemann-Liouville Fractional Integral.

*Remark 1.1.* Some special cases of Definition 2 are as follows.

- (1) If  $\varphi = 0$ , then  $K$  is called an invex set.
- (2) If  $\eta(v, u) = v - u$ , then  $K$  is called a  $\varphi$ -convex set.
- (3) If  $\varphi = 0$  and  $\eta(v, u) = v - u$ , then  $K$  is called a convex set.

**Definition 1.2.** (see [13]) The set  $K_{\varphi\eta}$  in  $\mathbb{R}^n$  is said to be  $\varphi$ -invex at  $u$  with respect to  $\varphi(\cdot)$ , if there exists a bifunction  $\eta(\cdot, \cdot) : K_{\varphi\eta} \times K_{\varphi\eta} \rightarrow \mathbb{R}$ , such that

$$(1.4) \quad u + te^{i\varphi}\eta(v, u) \in K_{\varphi\eta}, \quad \forall u, v \in K_{\varphi\eta}, t \in [0, 1].$$

Where  $\mathbb{R}^n$  Euclidian space. The  $\varphi$ -invex set  $K_{\varphi\eta}$  is also called  $\varphi\eta$ -connected set. Note that the convex set with  $\varphi = 0$  and  $\eta(v, u) = v - u$  is a  $\varphi$ -invex set, but the converse is not true.

**Definition 1.3.** Let  $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$  be a nonnegative function. A function  $f$  on the set  $K_{\varphi\eta}$  is said to be  $P_{\varphi}$ -preinvex function according to  $\varphi$  and bifunction  $\eta$ . Let  $\forall u, v \in I$ ,  $\eta(v, u) > 0$  and  $t \in (0, 1)$ , then

$$(1.5) \quad f(u + te^{i\varphi}\eta(v, u)) \leq f(u) + f(v).$$

## 2. MAIN RESULTS

In this section, we will give lemma which we use later in this work.

**Lemma 2.1.** Let  $f : [a, b] \subset [0, \infty) \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ . Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$ ,  $p, q > 0$  and  $a < b$ . Then the following equality holds,

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ &= [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 (1-t)^p t^q f(a+(1-t)e^{i\varphi}\eta(b,a)) dt. \end{aligned}$$

*Proof.* By using Definition 3, if left-hand side of equality use  $x = a+(1-t)e^{i\varphi}\eta(b,a)$ , we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ &= \int_0^1 ((1-t)e^{i\varphi}\eta(b,a))^p (te^{i\varphi}\eta(b,a))^q f(a+(1-t)e^{i\varphi}\eta(b,a)) e^{i\varphi}\eta(b,a) dt \\ &= [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 (1-t)^p t^q f(a+(1-t)e^{i\varphi}\eta(b,a)) dt, \end{aligned}$$

the proof is done. □

*Remark 2.1.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Lemma 1, we obtain Lemma 1 in [27],

$$\begin{aligned} & \int_a^b (x-a)^p (b-x)^q f(x) dx \\ &= [b-a]^{p+q+1} \int_0^1 (1-t)^p t^q f(at+(1-t)b) dt. \end{aligned}$$

**Theorem 2.1.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ . Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$  and  $0 \leq a < b < \infty$ . If  $f$  is quasi-preinvek on  $[a, b]$ , then for some fixed  $p, q > 0$ , we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) \max\{f(a), f(b)\}, \end{aligned}$$

here  $\beta(x, y)$  is the Euler Beta function.

*Proof.* By using inequality in (1.2), if left-hand side of equality use  $x = a + (1-t)e^{i\varphi}\eta(b, a)$ , we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & = \int_0^1 ((1-t)e^{i\varphi}\eta(b,a))^p (te^{i\varphi}\eta(b,a))^q f(a+(1-t)e^{i\varphi}\eta(b,a)) e^{i\varphi}\eta(b,a) dt \\ & = [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 (1-t)^p t^q f(a+(1-t)e^{i\varphi}\eta(b,a)) dt \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) \max\{f(a), f(b)\}, \end{aligned}$$

the proof is done.  $\square$

*Remark 2.2.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 1, we obtain Theorem 1 in [27]

$$\begin{aligned} & \int_a^b (x-a)^p (b-x)^q f(x) dx \\ & \leq (b-a)^{p+q+1} \beta(p+1, q+1) \max\{f(a), f(b)\}. \end{aligned}$$

**Theorem 2.2.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ . Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$ ,  $p, q > 0$  and  $0 \leq a < b < \infty$ . If  $|f|$  is  $P_\varphi$ -preinvek on  $[a, b]$ , then following inequality, we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) (|f(a)| + |f(b)|), \end{aligned}$$

*Proof.* By using Definition 3, if left-hand side of equality use  $x = a + (1-t)e^{i\varphi}\eta(b, a)$ , we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))| dt \\ & \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 [(1-t)^p t^q] (|f(a)| + |f(b)|) dt \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) (|f(a)| + |f(b)|), \end{aligned}$$

the proof is done.  $\square$

*Remark 2.3.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 2, we obtain Theorem 4 in [27],

$$\int_a^b (x-a)^p (b-x)^q f(x) dx \leq (b-a)^{p+q+1} \beta(p+1, q+1) (|f(a)| + |f(b)|).$$

**Theorem 2.3.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ .

Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$ ,  $p, q > 0$  and  $0 \leq a < b < \infty$ . If  $|f|^{\frac{k}{k-1}}$  is quasi-preinvex on  $[a, b]$  and  $k > 1$ , then following inequality, we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( \max \left\{ |f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}} \right\} \right)^{\frac{k-1}{k}}. \end{aligned}$$

*Proof.* By using lemma 1, quasi-preinvex of  $|f|^{\frac{k}{k-1}}$  and Hölder's inequality, we obtain

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))| dt \\ & \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 |(1-t)^p t^q|^k dt \right)^{\frac{1}{k}} \left( \int_0^1 |f(a+(1-t)e^{i\varphi}\eta(b,a))|^{\frac{k}{k-1}} dt \right)^{\frac{k-1}{k}} \\ & \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 (1-t)^{kp} t^{kq} dt \right)^{\frac{1}{k}} \left( \int_0^1 \max \left( |f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}} \right) dt \right)^{\frac{k-1}{k}} \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta^{\frac{1}{k}}(kp+1, kq+1) \left[ \max \left( |f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}} \right) \right]^{\frac{k-1}{k}}, \end{aligned}$$

the proof is done.  $\square$

*Remark 2.4.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 3, we obtain Theorem 2 in [27],

$$\int_a^b (x-a)^p (b-x)^q f(x) dx \leq (b-a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( \max \left\{ |f(a)|^{\frac{k}{k-1}}, |f(b)|^{\frac{k}{k-1}} \right\} \right)^{\frac{k-1}{k}}.$$

**Theorem 2.4.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ .

Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$ ,  $p, q > 0$  and  $0 \leq a < b < \infty$ . If  $|f|^{\frac{k}{k-1}}$  is  $P_\varphi$ -preinvex on  $[a, b]$  and  $k > 1$ , then following inequality, we have

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( |f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right)^{\frac{k-1}{k}}. \end{aligned}$$



*Proof.* By using lemma 1,  $P_\varphi$ -preinvek of  $|f|^{\frac{k}{k-1}}$  and Hölder's inequality, we obtain

$$\begin{aligned}
& \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))| dt \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 |(1-t)^p t^q|^k dt \right)^{\frac{1}{k}} \left( \int_0^1 |f(a+(1-t)e^{i\varphi}\eta(b,a))|^{\frac{k}{k-1}} dt \right)^{\frac{k-1}{k}} \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 (1-t)^{kp} t^{kq} dt \right)^{\frac{1}{k}} \left( \int_0^1 (|f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}}) dt \right)^{\frac{k-1}{k}} \\
& \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta^{\frac{1}{k}} (kp+1, kq+1) \left[ |f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right]^{\frac{k-1}{k}},
\end{aligned}$$

The proof is done.  $\square$

*Remark 2.5.* If we consider  $\eta(b,a) = b-a$  and  $\varphi = 0$  in Theorem 4, we obtain Theorem 5 in [27],

$$\begin{aligned}
& \int_a^b (x-a)^p (b-x)^q f(x) dx \\
& \leq (b-a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( |f(a)|^{\frac{k}{k-1}} + |f(b)|^{\frac{k}{k-1}} \right)^{\frac{k-1}{k}}.
\end{aligned}$$

**Theorem 2.5.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ .

Where is  $f \in L([a, a+e^{i\varphi}\eta(b,a)])$ ,  $p, q > 0$  and  $0 \leq a < b < \infty$ . If  $|f|^l$  is quasi-preinvek on  $[a, b]$  and  $l \geq 1$ , then following inequality, we have

$$\begin{aligned}
& \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\
& \leq (e^{i\varphi}\eta(b,a))^{p+q+1} [\beta(p+1, q+1)] \left( \max \left\{ |f(a)|^l, |f(b)|^l \right\} \right)^{\frac{1}{l}}.
\end{aligned}$$

*Proof.* By using lemma 1, quasi-preinvek of  $|f|^l$  and Power Mean inequality, we obtain

$$\begin{aligned}
& \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))| dt \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 |(1-t)^p t^q| dt \right)^{1-\frac{1}{l}} \left( \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))|^l dt \right)^{\frac{1}{l}} \\
& \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} (\beta(p+1, q+1))^{1-\frac{1}{l}} \left( \int_0^1 |(1-t)^p t^q| \max \left( |f(a)|^l, |f(b)|^l \right) dt \right)^{\frac{1}{l}} \\
& \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) \left[ \max \left( |f(a)|^l, |f(b)|^l \right) \right]^{\frac{1}{l}},
\end{aligned}$$

the proof is done.  $\square$

*Remark 2.6.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 5, we obtain Theorem 3 in [27],

$$\begin{aligned} \int_a^b (x-a)^p (b-x)^q f(x) dx \\ \leq (b-a)^{p+q+1} [\beta(p+1, q+1)] \left( \max \left\{ |f(a)|^l, |f(b)|^l \right\} \right)^{\frac{1}{l}}. \end{aligned}$$

**Theorem 2.6.** Let  $f : [a, b] \rightarrow \mathbb{R}$  be continuous on  $[a, b]$ .

Where is  $f \in L([a, a + e^{i\varphi}\eta(b, a)])$ ,  $p, q > 0$  and  $0 \leq a < b < \infty$ . If  $|f|^l$  is quasi-preinvex on  $[a, b]$  and  $l \geq 1$ , then following inequality, we have

$$\begin{aligned} \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ \leq (e^{i\varphi}\eta(b,a))^{p+q+1} [\beta(p+1, q+1)] \left( |f(a)|^l + |f(b)|^l \right)^{\frac{1}{l}}. \end{aligned}$$

*Proof.* By using lemma 1,  $P_\varphi$ -preinvex of  $|f|^l$  and Power Mean inequality, we obtain

$$\begin{aligned} \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x) dx \\ \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))| dt \\ \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} \left( \int_0^1 |(1-t)^p t^q| dt \right)^{1-\frac{1}{l}} \left( \int_0^1 |(1-t)^p t^q| |f(a+(1-t)e^{i\varphi}\eta(b,a))|^l dt \right)^{\frac{1}{l}} \\ \leq [e^{i\varphi}\eta(b,a)]^{p+q+1} (\beta(p+1, q+1))^{1-\frac{1}{l}} \left( \int_0^1 |(1-t)^p t^q| \left( |f(a)|^l + |f(b)|^l \right) dt \right)^{\frac{1}{l}} \\ \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) \left[ |f(a)|^l + |f(b)|^l \right]^{\frac{1}{l}}, \end{aligned}$$

the proof is done.

In this section some new integral inequalities for functions of several variables on preinvex subsets of  $\mathbb{R}^n$  will be given. First we recall the notion of  $P_\varphi$ -preinvex convexity for functions on a preinvex subset  $U$  of  $\mathbb{R}^n$ .  $\square$

*Remark 2.7.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 6, we obtain Theorem 6 in [27],

$$\begin{aligned} \int_a^b (x-a)^p (b-x)^q f(x) dx \\ \leq (b-a)^{p+q+1} [\beta(p+1, q+1)] \left( |f(a)|^l + |f(b)|^l \right)^{\frac{1}{l}}. \end{aligned}$$

**Definition 2.1.** The functions  $f : U \rightarrow \mathbb{R}$  is said to be  $P_\varphi$ -preinvex convexity on  $U$  if it is nonnegative and, for all  $x, y \in U$  and  $\lambda \in [0, 1]$ , satisfies the inequality

$$f(x + (1-\lambda)e^{i\varphi}\eta(y, x)) \leq f(x) + f(y).$$

The following proposition will be used throughout this section.

**Proposition 2.1.** Let  $U \subseteq \mathbb{R}$  be a preinvex subset of  $\mathbb{R}$  and  $f : U \rightarrow \mathbb{R}$  be a function. Then  $f$  is  $P_\varphi$ -preinvex on  $U$  if and only if, for every  $x, y \in U$ , the function  $\varphi : [0, 1] \rightarrow \mathbb{R}$ , defined by

$$\varphi(t) := f(x + te^{i\varphi}\eta(y, x)),$$

is  $P_\varphi$ -convex on  $I$  with  $I = [0, 1]$ .

**Theorem 2.7.** *Let  $U \subseteq \mathbb{R}$  be a preinvex subset of  $\mathbb{R}$ . Assume that  $f : U \rightarrow \mathbb{R}^+$  is a  $P_\varphi$ -preinvex function on  $U$ . Then, for every  $x, y \in U$  and every  $[a, b] \in [0, 1]$  with  $a < b$ , the following inequality holds:*

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (t-a)^p (a+e^{i\varphi}\eta(b,a)-t)^q f(x+te^{i\varphi}\eta(y,x)) dt \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) [f(x+ae^{i\varphi}\eta(y,x)) + f(x+be^{i\varphi}\eta(y,x))]. \end{aligned}$$

*Proof.* Let  $x, y \in U$  and every  $[a, b] \in [0, 1]$  with  $a < b$ . Since  $f : U \rightarrow \mathbb{R}^+$  is a  $P_\varphi$ -preinvex function, by Proposition 1 the function  $\varphi : [0, 1] \rightarrow \mathbb{R}^+$  defined by

$$\varphi(t) := f(x+te^{i\varphi}\eta(y,x)),$$

is  $P_\varphi$ -preinvex on  $I$  with  $I = [0, 1]$ . Applying Theorem 4 to the function  $\varphi$  implies that

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (t-a)^p (a+e^{i\varphi}\eta(b,a)-t)^q \varphi(t) dt \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) [|\varphi(a)| + |\varphi(b)|] \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) [f(x+ae^{i\varphi}\eta(y,x)) + f(x+be^{i\varphi}\eta(y,x))], \end{aligned}$$

the proof is done.  $\square$

*Remark 2.8.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 7, we obtain Theorem 7 in [27],

$$\begin{aligned} & \int_a^b (t-a)^p (b-t)^q f((1-t)x+ty) dt \\ & (b-a)^{p+q+1} \beta(p+1, q+1) [f((1-a)x+ay) + f((1-b)x+by)]. \end{aligned}$$

**Theorem 2.8.** *Let  $U \subseteq \mathbb{R}$  be a preinvex subset of  $\mathbb{R}$  and let  $k > 1$ . Assume that  $f^{\frac{k}{k-1}} : U \rightarrow \mathbb{R}^+$  is a  $P_\varphi$ -preinvex function on  $U$ . Then, for every  $x, y \in U$  and every  $[a, b] \in [0, 1]$  with  $a < b$ , the following inequality holds:*

$$\begin{aligned} & \int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x+te^{i\varphi}\eta(y,x)) dx \\ & \leq (e^{i\varphi}\eta(b,a))^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( f^{\frac{k}{k-1}}(x+ae^{i\varphi}\eta(y,x)) + f^{\frac{k}{k-1}}(x+be^{i\varphi}\eta(y,x)) \right)^{\frac{k-1}{k}}. \end{aligned}$$

*Remark 2.9.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 8, we obtain Theorem 8 in [27],

$$\begin{aligned} & \int_a^b (t-a)^p (b-t)^q f((1-t)x+ty) dt \\ & \leq (b-a)^{p+q+1} [\beta(kp+1, kq+1)]^{\frac{1}{k}} \left( f^{\frac{k}{k-1}}((1-a)x+ay) + f^{\frac{k}{k-1}}((1-b)x+by) \right)^{\frac{k-1}{k}}. \end{aligned}$$

**Theorem 2.9.** *Let  $U \subseteq \mathbb{R}$  be a preinvex subset of  $\mathbb{R}$  and let  $k > 1$ . Assume that  $f^l : U \rightarrow \mathbb{R}^+$  is a  $P_\varphi$ -preinvex function on  $U$ . Then, for every  $x, y \in U$  and every*

$[a, b] \in [0, 1]$  with  $a < b$ , the following inequality holds:

$$\int_a^{a+e^{i\varphi}\eta(b,a)} (x-a)^p (a+e^{i\varphi}\eta(b,a)-x)^q f(x+te^{i\varphi}\eta(y,x)) dx \\ \leq (e^{i\varphi}\eta(b,a))^{p+q+1} \beta(p+1, q+1) \left( f^l(x+ae^{i\varphi}\eta(y,x)) + f^l(x+be^{i\varphi}\eta(y,x)) \right)^{\frac{1}{l}}.$$

*Remark 2.10.* If we consider  $\eta(b, a) = b - a$  and  $\varphi = 0$  in Theorem 9, we obtain Theorem 9 in [27],

$$\int_a^b (t-a)^p (b-t)^q f((1-t)x+ty) dt \\ \leq (b-a)^{p+q+1} \beta(p+1, q+1) (f^l((1-a)x+ay) + f^l((1-b)x+by))^{\frac{1}{l}}.$$

#### REFERENCES

- [1] A. O. Akdemir and M. E. Özdemir, "Some Hadamard- type inequalities for coordinated P-convex functions and Godunova-Levin functions,' AIP Conf. Proc., vol. 1309, pp. 7-15,2010.
- [2] M. Alomari, M. Darus, and S.S. Dragomir, 'Inequalities of Hermite-Hadamard's type for functions whose derivates absolute values are quasi-convex. " RGMIA Res. Rep. Coll., vol. 12, p.11, Article 14,2009.
- [3] A. Barani and S. Barani, "Hermite-Hadamard type inequalities for functions when a power of the absolute value of the first derivative is P-convex, " Bull. Aust. Math. Soc., vol. 86, no. 1, pp. 126-134, 2012.
- [4] S. S. Dragomir, "On some new inequalities of Hermite-Hadamard type for m-convex functions. "Tamkang J. Math., vol. 33, no. 1, pp. 55-65, 2012
- [5] S. S. Dagomir and C. E. M. Pearce, Selected toics on Hermite-Hadamard inequalities and applications, ser. RGMIA Monographs. Victoria University, 2000.
- [6] S. S. Dragomir, J. E. Pečarić, and L. Persson, "Some inequalities of Hadamard type" Soochow J. Math., vol. 21, no. 3, pp. 335-341, 1995.
- [7] S. S. Dragomir and C. Pearce, "Quasi-convex functions and Hadamar's inequality" Bull. Aust. Math. Soc., vol.57 no. 3, pp. 377-385, 1998.
- [8] A. Hāzy, "Bernstein- Doetsch type results for h-convex functions,"Math. Inequal., vol. 14, no. 3, pp. 499-508,2011.
- [9] A. Hāzy, "Bernstein- Doetsch type results for (k, h)-convex functions,"Miskolc Math. Notes, vol. 13, no. 2, pp. 325-336, 2012.
- [10] V.N.Huy and Nguyen Thanh Chung, "Some generalizations of the Fejër and Hermide-Hadamard inequalities in Hölder spaces ," J. Appl. Math. Inform., vol. 29, no. 3-4, pp. 859-868. 2011.
- [11] D.A. Ion, "Some estimates on the Hermite- Hadamard inequality through quasi-convex functions,"An. Univ. Craiova, Ser. Mat. Inf., vol. 34, pp. 83-88, 2007.
- [12] W. Liu, "New integral inequalities via  $(\alpha, m)$ -convexity and quasi-convexity, "Hacet. J. Math. Stat., vol. 42, no. 3, pp. 289-297, 2013.
- [13] W. Liu, "Some Ostrowski type inequalities via Riemann-Liouville fractional intagrals for  $h$ -convex functions," J. Comput. Anal. Appl., vol. 16, no. 5, pp. 998-1004, 2014.
- [14] W. Liu, "Some Simpson type inequalities for  $h$ -convex and  $(\alpha, m)$ -convex functions," J. Comput. Anal. Appl., vol. 16, no. 5, pp. 1005-1012, 2014
- [15] B. Micherda and T. Rajba, " On some Hermite-Hadamard-Fejër inequalities for  $(k, h)$ -convex functions," Math. Inequal. Appl. vol. 15, no. 4, pp. 931-940, 2012.
- [16] M. E. Özdemir and C. Yildiz, " New inequalities for Hermite- Hadamard and simpson type and applications, " arXiv, no. 1103.1965v1 [math. CA].
- [17] M. E. Özdemir, E. Set, and M.Alomari, "Integral inequalities via several kinds of convexity,"Creat. Math. Inform., vol. 20, no. 1, pp. 62-73, 2011.
- [18] M. Z. Sarikaya, A. Saglam, and H. Yildirim, " On some Hadamard-type inequalities for  $h$ -convex functions," J. Math. Inequal., vol.2, no.3, pp. 335-341,2008.
- [19] M. Z. Sarikaya, E. Set, and M. E. Özdemir, " On some new inequalities of Hadamard type involving  $h$ -convex functions," Acta Math. Univ. Comen., New Ser., vol. 79, no. 2, pp. 265-272, 2010.

- [20] E. Set, M. Sardari, M.E.Özdemir, and J. Roojin, "On generalizations of the Hadamard inequality for  $(\alpha, m)$ -convex functions," RGMIA Res. Rep. Coll., vol. 12,p. 10, Article 4, 2009
- [21] E. Set, M. E. Özdemir, and S. S. Dragomir," On Hadamard type inequalities involving several kinds of convexity, " J. Inequal. Appl., vol. 2010, p. 12,2010.
- [22] D. D. Stancu , G. Coman, and P. Blaga, Analiză numerică și teoria aproximării Vol. II. Cluj-Napoca: Presa Universitară Clujeană, 2012.
- [23] K. L. Tseng, G. S. Yang, and S. S. Dragamir, "On quasiconvex functions and Hadamard's inequality," RGMIA Res. Rep. Coll., vol. 6, NO. 3,p. 12, Article 1, 2003.
- [24] M. Tunç, "On some nequalities via  $h$ -convexity," Miskolc Math. Notes, vol. 14, no. 3, pp. 1041-1057, 2013.
- [25] M. Tunç," Ostrowski type inequalities via  $h$ -convex functions with applications to special means, " J.Inequal. Appl., vol. 2013, p. 10, 2013.
- [26] S. Varošanec, " On  $h$ -convexity, " J. Math. Appl., vol. 326, no. 1, pp. 303-311,2007.
- [27] W. Liu, "New Integral Inequalities Involving Beta Function via P-Convexity," Miskolc Matematical Notes, Vol. 15 No. 2, pp. 585-591, 2014.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, UNIVERSITY OF KAHRAMANMARAŞ  
SÜTÇÜ İMAM, 46000, KAHRAMANMARAŞ, TURKEY

*E-mail address:* `sedaaa.kilinc@hotmail.com`

*E-mail address:* `sumeyye_ermeydan@hotmail.com`

*E-mail address:* `hyildir@ksu.edu.tr`

**BOUNDEDNESS OF THE SUBLINEAR OPERATORS WITH  
ROUGH KERNEL GENERATED BY FRACTIONAL INTEGRALS  
AND THEIR COMMUTATORS ON GENERALIZED VANISHING  
MORREY SPACES II**

FERIT GURBUZ

ABSTRACT. In this paper, we consider the norm inequalities for sublinear operators with rough kernel generated by fractional integrals and their commutators on generalized Morrey spaces and on generalized vanishing Morrey spaces including their weak versions under generic size conditions which are satisfied by most of the operators in harmonic analysis, respectively. In all the cases the conditions for the boundedness of sublinear operators with rough kernel and their commutators are given in terms of Zygmund-type integral inequalities on  $(\varphi_1, \varphi_2)$ , where there is no assumption on monotonicity of  $\varphi_1, \varphi_2$  in  $r$ . As an example to the conditions of these theorems are satisfied, we can consider the Marcinkiewicz operator.

Received: 27–July–2016

Accepted: 29–August–2016

1. INTRODUCTION

The classical Morrey spaces  $M_{p,\lambda}$  have been introduced by Morrey in [32] to study the local behavior of solutions of second order elliptic partial differential equations(PDEs). In recent years there has been an explosion of interest in the study of the boundedness of operators on Morrey-type spaces. It has been obtained that many properties of solutions to PDEs are concerned with the boundedness of some operators on Morrey-type spaces. In fact, better inclusion between Morrey and Hölder spaces allows to obtain higher regularity of the solutions to different elliptic and parabolic boundary problems (see [14, 36, 41, 43] for details).

Let  $B = B(x_0, r_B)$  denote the ball with the center  $x_0$  and radius  $r_B$ . For a given measurable set  $E$ , we also denote the Lebesgue measure of  $E$  by  $|E|$ . For any given  $\Omega_0 \subseteq \mathbb{R}^n$  and  $0 < p < \infty$ , denote by  $L_p(\Omega_0)$  the spaces of all functions  $f$  satisfying

$$\|f\|_{L_p(\Omega_0)} = \left( \int_{\Omega_0} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty.$$

We recall the definition of classical Morrey spaces  $M_{p,\lambda}$  as

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 42B20, 42B25, 42B35.

*Key words and phrases.* Sublinear operator; fractional integral operator; rough kernel; generalized vanishing Morrey space; commutator; *BMO*.

$$M_{p,\lambda}(\mathbb{R}^n) = \left\{ f : \|f\|_{M_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))} < \infty \right\},$$

where  $f \in L_p^{loc}(\mathbb{R}^n)$ ,  $0 \leq \lambda \leq n$  and  $1 \leq p < \infty$ .

Note that  $M_{p,0} = L_p(\mathbb{R}^n)$  and  $M_{p,n} = L_\infty(\mathbb{R}^n)$ . If  $\lambda < 0$  or  $\lambda > n$ , then  $M_{p,\lambda} = \Theta$ , where  $\Theta$  is the set of all functions equivalent to 0 on  $\mathbb{R}^n$ .

We also denote by  $WM_{p,\lambda} \equiv WM_{p,\lambda}(\mathbb{R}^n)$  the weak Morrey space of all functions  $f \in WL_p^{loc}(\mathbb{R}^n)$  for which

$$\|f\|_{WM_{p,\lambda}} \equiv \|f\|_{WM_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{WL_p(B(x,r))} < \infty,$$

where  $WL_p(B(x,r))$  denotes the weak  $L_p$ -space of measurable functions  $f$  for which

$$\begin{aligned} \|f\|_{WL_p(B(x,r))} &\equiv \|f\chi_{B(x,r)}\|_{WL_p(\mathbb{R}^n)} \\ &= \sup_{t > 0} t |\{y \in B(x,r) : |f(y)| > t\}|^{1/p} \\ &= \sup_{0 < t \leq |B(x,r)|} t^{1/p} (f\chi_{B(x,r)})^*(t) < \infty, \end{aligned}$$

where  $g^*$  denotes the non-increasing rearrangement of a function  $g$ .

Throughout the paper we assume that  $x \in \mathbb{R}^n$  and  $r > 0$  and also let  $B(x,r)$  denotes the open ball centered at  $x$  of radius  $r$ ,  $B^C(x,r)$  denotes its complement and  $|B(x,r)|$  is the Lebesgue measure of the ball  $B(x,r)$  and  $|B(x,r)| = v_n r^n$ , where  $v_n = |B(0,1)|$ . It is known that  $M_{p,\lambda}(\mathbb{R}^n)$  is an extension (a generalization) of  $L_p(\mathbb{R}^n)$  in the sense that  $M_{p,0} = L_p(\mathbb{R}^n)$ .

Morrey has stated that many properties of solutions to PDEs can be attributed to the boundedness of some operators on Morrey spaces. For the boundedness of the Hardy–Littlewood maximal operator, the fractional integral operator and the Calderón–Zygmund singular integral operator on these spaces, we refer the readers to [1, 6, 38]. For the properties and applications of classical Morrey spaces, see [7, 8, 14, 36, 41, 43] and references therein. The generalized Morrey spaces  $M_{p,\varphi}$  are obtained by replacing  $r^\lambda$  with a function  $\varphi(r)$  in the definition of the Morrey space. During the last decades various classical operators, such as maximal, singular and potential operators have been widely investigated in classical and generalized Morrey spaces.

The study of the operators of harmonic analysis in vanishing Morrey space, in fact has been almost not touched. A version of the classical Morrey space  $M_{p,\lambda}(\mathbb{R}^n)$  where it is possible to approximate by "nice" functions is the so called vanishing Morrey space  $VM_{p,\lambda}(\mathbb{R}^n)$  has been introduced by Vitanza in [50] and has been applied there to obtain a regularity result for elliptic PDEs. This is a subspace of functions in  $M_{p,\lambda}(\mathbb{R}^n)$ , which satisfies the condition

$$\lim_{r \rightarrow 0} \sup_{\substack{x \in \mathbb{R}^n \\ 0 < t < r}} t^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,t))} = 0.$$

Later in [51] Vitanza has proved an existence theorem for a Dirichlet problem, under weaker assumptions than in [30] and a  $W^{3,2}$  regularity result assuming that the partial derivatives of the coefficients of the highest and lower order terms belong to vanishing Morrey spaces depending on the dimension. Also Ragusa has proved a sufficient condition for commutators of fractional integral operators to belong to

vanishing Morrey spaces  $VM_{p,\lambda}(\mathbb{R}^n)$  (see [39, 40]). For the properties and applications of vanishing Morrey spaces, see also [3]. It is known that, there is no research regarding boundedness of the sublinear operators with rough kernel on vanishing Morrey spaces.

Maximal functions and singular integrals play a key role in harmonic analysis since maximal functions could control crucial quantitative information concerning the given functions, despite their larger size, while singular integrals, Hilbert transform as it's prototype, recently intimately connected with PDEs, operator theory and other fields.

Let  $f \in L^{loc}(\mathbb{R}^n)$ . The Hardy-Littlewood(H-L) maximal operator  $M$  is defined by

$$Mf(x) = \sup_{t>0} |B(x,t)|^{-1} \int_{B(x,t)} |f(y)|dy.$$

Let  $\bar{T}$  be a standard Calderón-Zygmund(C-Z) singular integral operator, briefly a C-Z operator, i.e., a linear operator bounded from  $L_2(\mathbb{R}^n)$  to  $L_2(\mathbb{R}^n)$  taking all infinitely continuously differentiable functions  $f$  with compact support to the functions  $f \in L_1^{loc}(\mathbb{R}^n)$  represented by

$$\bar{T}f(x) = p.v. \int_{\mathbb{R}^n} k(x-y)f(y)dy \quad x \notin \text{supp}f.$$

Such operators have been introduced in [11]. Here  $k$  is a C-Z kernel [16]. Chiarenza and Frasca [6] have obtained the boundedness of H-L maximal operator  $M$  and C-Z operator  $\bar{T}$  on  $M_{p,\lambda}(\mathbb{R}^n)$ . It is also well known that H-L maximal operator  $M$  and C-Z operator  $\bar{T}$  play an important role in harmonic analysis (see [15, 29, 46, 47, 48]). Also, the theory of the C-Z operator is one of the important achievements of classical analysis in the last century, which has many important applications in Fourier analysis, complex analysis, operator theory and so on.

Let  $f \in L^{loc}(\mathbb{R}^n)$ . The fractional maximal operator  $M_\alpha$  and the fractional integral operator (also known as the Riesz potential)  $\bar{T}_\alpha$  are defined by

$$M_\alpha f(x) = \sup_{t>0} |B(x,t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |f(y)|dy \quad 0 \leq \alpha < n$$

$$\bar{T}_\alpha f(x) = \int_{\mathbb{R}^n} \frac{f(y)}{|x-y|^{n-\alpha}} dy \quad 0 < \alpha < n.$$

It is well known that  $M_\alpha$  and  $\bar{T}_\alpha$  play an important role in harmonic analysis (see [47, 48]).

An early impetus to the study of fractional integrals originated from the problem of fractional derivation, see e.g. [35]. Besides its contributions to harmonic analysis, fractional integrals also play an essential role in many other fields. The H-L Sobolev inequality about fractional integral is still an indispensable tool to establish time-space estimates for the heat semigroup of nonlinear evolution equations, for some of this work, see e.g. [24]. In recent times, the applications to Chaos and Fractal have become another motivation to study fractional integrals, see e.g. [26]. It is well known that  $\bar{T}_\alpha$  is bounded from  $L_p$  to  $L_q$ , where  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$  and  $1 < p < \frac{n}{\alpha}$ .



Spanne (published by Peetre [38]) and Adams [1] have studied boundedness of the fractional integral operator  $\bar{T}_\alpha$  on  $M_{p,\lambda}(\mathbb{R}^n)$ . Their results, can be summarized as follows.

**Theorem 1.1.** (*Spanne, but published by Peetre [38]*) *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $0 < \lambda < n - \alpha p$ . Moreover, let  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$  and  $\frac{\lambda}{p} = \frac{\mu}{q}$ . Then for  $p > 1$  the operator  $\bar{T}_\alpha$  is bounded from  $M_{p,\lambda}$  to  $M_{q,\lambda}$  and for  $p = 1$  the operator  $\bar{T}_\alpha$  is bounded from  $M_{1,\lambda}$  to  $WM_{q,\lambda}$ .*

**Theorem 1.2.** (*Adams [1]*) *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $0 < \lambda < n - \alpha p$  and  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n-\lambda}$ . Then for  $p > 1$  the operator  $\bar{T}_\alpha$  is bounded from  $M_{p,\lambda}$  to  $M_{q,\lambda}$  and for  $p = 1$  the operator  $\bar{T}_\alpha$  is bounded from  $M_{1,\lambda}$  to  $WM_{q,\lambda}$ .*

Recall that, for  $0 < \alpha < n$ ,

$$M_\alpha f(x) \leq \nu_n^{\frac{\alpha}{n}-1} \bar{T}_\alpha(|f|)(x)$$

holds (see [25], Remark 2.1). Hence Theorems 1.1 and 1.2 also imply boundedness of the fractional maximal operator  $M_\alpha$ , where  $\nu_n$  is the volume of the unit ball on  $\mathbb{R}^n$ .

Suppose that  $S^{n-1}$  is the unit sphere in  $\mathbb{R}^n$  ( $n \geq 2$ ) equipped with the normalized Lebesgue measure  $d\sigma$ . Let  $\Omega \in L_s(S^{n-1})$  with  $1 < s \leq \infty$  be homogeneous of degree zero. We define  $s' = \frac{s}{s-1}$  for any  $s > 1$ . Suppose that  $T_{\Omega,\alpha}$ ,  $\alpha \in (0, n)$  represents a linear or a sublinear operator, which satisfies that for any  $f \in L_1(\mathbb{R}^n)$  with compact support and  $x \notin \text{supp} f$

$$(1.1) \quad |T_{\Omega,\alpha} f(x)| \leq c_0 \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |f(y)| dy,$$

where  $c_0$  is independent of  $f$  and  $x$ .

For a locally integrable function  $b$  on  $\mathbb{R}^n$ , suppose that the commutator operator  $T_{\Omega,b,\alpha}$ ,  $\alpha \in (0, n)$  represents a linear or a sublinear operator, which satisfies that for any  $f \in L_1(\mathbb{R}^n)$  with compact support and  $x \notin \text{supp} f$

$$(1.2) \quad |T_{\Omega,b,\alpha} f(x)| \leq c_0 \int_{\mathbb{R}^n} |b(x) - b(y)| \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |f(y)| dy,$$

where  $c_0$  is independent of  $f$  and  $x$ .

We point out that the condition (1.1) in the case of  $\Omega \equiv 1$ ,  $\alpha = 0$  has been introduced by Soria and Weiss in [44]. The conditions (1.1) and (1.2) are satisfied by many interesting operators in harmonic analysis, such as fractional Marcinkiewicz operator, fractional maximal operator, fractional integral operator (Riesz potential) and so on (see [27], [44] for details).

In 1971, Muckenhoupt and Wheeden [34] defined the fractional integral operator with rough kernel  $\bar{T}_{\Omega,\alpha}$  by

$$\bar{T}_{\Omega,\alpha} f(x) = \int_{\mathbb{R}^n} \frac{\Omega(x-y)}{|x-y|^{n-\alpha}} f(y) dy \quad 0 < \alpha < n$$

and a related fractional maximal operator with rough kernel  $M_{\Omega,\alpha}$  is given by

$$M_{\Omega,\alpha}f(x) = \sup_{t>0} |B(x,t)|^{-1+\frac{\alpha}{n}} \int_{B(x,t)} |\Omega(x-y)| |f(y)| dy \quad 0 \leq \alpha < n,$$

where  $\Omega \in L_s(S^{n-1})$  with  $1 < s \leq \infty$  is homogeneous of degree zero on  $\mathbb{R}^n$  and  $\bar{T}_{\Omega,\alpha}$  satisfies the condition (1.1).

If  $\alpha = 0$ , then  $M_{\Omega,0} \equiv M_{\Omega}$  H-L maximal operator with rough kernel. It is obvious that when  $\Omega \equiv 1$ ,  $M_{1,\alpha} \equiv M_{\alpha}$  and  $\bar{T}_{1,\alpha} \equiv \bar{T}_{\alpha}$  are the fractional maximal operator and the fractional integral operator, respectively.

In recent years, the mapping properties of  $\bar{T}_{\Omega,\alpha}$  on some kinds of function spaces have been studied in many papers (see [5], [12], [13], [34] for details). In particular, the boundedness of  $\bar{T}_{\Omega,\alpha}$  in Lebesgue spaces has been obtained.

**Lemma 1.1.** [5, 12, 33] *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . If  $\Omega \in L_s(S^{n-1})$ ,  $s > \frac{n}{n-\alpha}$ , then we have*

$$\|\bar{T}_{\Omega,\alpha}f\|_{L_q} \leq C \|f\|_{L_p}.$$

**Corollary 1.1.** *Under the assumptions of Lemma 1.1, the operator  $M_{\Omega,\alpha}$  is bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ . Moreover, we have*

$$\|M_{\Omega,\alpha}f\|_{L_q} \leq C \|f\|_{L_p}.$$

*Proof.* Set

$$\tilde{T}_{|\Omega|,\alpha}(|f|)(x) = \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |f(y)| dy \quad 0 < \alpha < n,$$

where  $\Omega \in L_s(S^{n-1})$  ( $s > 1$ ) is homogeneous of degree zero on  $\mathbb{R}^n$ . It is easy to see that, for  $\tilde{T}_{|\Omega|,\alpha}$ , Lemma 1.1 is also hold. On the other hand, for any  $t > 0$ , we have

$$\begin{aligned} \tilde{T}_{|\Omega|,\alpha}(|f|)(x) &\geq \int_{B(x,t)} \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |f(y)| dy \\ &\geq \frac{1}{t^{n-\alpha}} \int_{B(x,t)} |\Omega(x-y)| |f(y)| dy. \end{aligned}$$

Taking the supremum for  $t > 0$  on the inequality above, we get

$$M_{\Omega,\alpha}f(x) \leq C_{n,\alpha}^{-1} \tilde{T}_{|\Omega|,\alpha}(|f|)(x) \quad C_{n,\alpha} = |B(0,1)|^{\frac{n-\alpha}{n}}.$$

□

In 1976, Coifman, Rocherberg and Weiss [9] introduced the commutator generated by  $\bar{T}_{\Omega}$  and a local integrable function  $b$ :

$$(1.3) \quad [b, \bar{T}_{\Omega}]f(x) \equiv b(x)\bar{T}_{\Omega}f(x) - \bar{T}_{\Omega}(bf)(x) = p.v. \int_{\mathbb{R}^n} [b(x) - b(y)] \frac{\Omega(x-y)}{|x-y|^n} f(y) dy.$$

Sometimes, the commutator defined by (1.3) is also called the commutator in Coifman-Rocherberg-Weiss's sense, which has its root in the complex analysis and harmonic analysis (see [9]).

Let  $b$  be a locally integrable function on  $\mathbb{R}^n$ , then for  $0 < \alpha < n$  and  $f$  is a suitable function, we define the commutators generated by fractional integral and maximal operators with rough kernel and  $b$  as follows, respectively:

$$[b, \bar{T}_{\Omega, \alpha}]f(x) \equiv b(x)\bar{T}_{\Omega, \alpha}f(x) - \bar{T}_{\Omega, \alpha}(bf)(x) = \int_{\mathbb{R}^n} [b(x) - b(y)] \frac{\Omega(x-y)}{|x-y|^{n-\alpha}} f(y) dy,$$

$$M_{\Omega, b, \alpha}(f)(x) = \sup_{t>0} |B(x, t)|^{-1+\frac{\alpha}{n}} \int_{B(x, t)} |b(x) - b(y)| |\Omega(x-y)| |f(y)| dy$$

satisfy condition (1.2). The proof of boundedness of  $[b, \bar{T}_{\Omega, \alpha}]$  in Lebesgue spaces can be found in [12] (by taking  $w = 1$  there).

**Theorem 1.3.** [12] *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero and has mean value zero on  $S^{n-1}$ . Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$ , and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ . If  $s' < p$  or  $q < s$ , then the operator  $[b, \bar{T}_{\Omega, \alpha}]$  is bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ .*

*Remark 1.1.* Using the method in the proof of Corollary 1.1 we have that

$$(1.4) \quad M_{\Omega, b, \alpha} f(x) \leq C_{n, \alpha}^{-1} [b, \bar{T}_{|\Omega|, \alpha}] (|f|)(x) \quad C_{n, \alpha} = |B(0, 1)|^{\frac{n-\alpha}{n}}.$$

By (1.4) we see that under the conditions of Theorem 1.3, the consequences of  $(L_p, L_q)$ -boundedness still hold for  $M_{\Omega, b, \alpha}$ .

*Remark 1.2.* [41, 42] When  $\Omega$  satisfies the specified size conditions, the kernel of the operator  $\bar{T}_{\Omega, \alpha}$  has no regularity, so the operator  $\bar{T}_{\Omega, \alpha}$  is called a rough fractional integral operator. In recent years, a variety of operators related to the fractional integrals, but lacking the smoothness required in the classical theory, have been studied. These include the operator  $[b, \bar{T}_{\Omega, \alpha}]$ . For more results, we refer the reader to [2, 4, 12, 13, 18, 19, 20, 28].

Finally, we present a relationship between essential supremum and essential infimum.

**Lemma 1.2.** (see [52] page 143) *Let  $f$  be a real-valued nonnegative function and measurable on  $E$ . Then*

$$(1.5) \quad \left( \operatorname{ess\,inf}_{x \in E} f(x) \right)^{-1} = \operatorname{ess\,sup}_{x \in E} \frac{1}{f(x)}.$$

Throughout the paper we use the letter  $C$  for a positive constant, independent of appropriate parameters and not necessarily the same at each occurrence. By  $A \lesssim B$  we mean that  $A \leq CB$  with some positive constant  $C$  independent of appropriate quantities. If  $A \lesssim B$  and  $B \lesssim A$ , we write  $A \approx B$  and say that  $A$  and  $B$  are equivalent.

## 2. GENERALIZED VANISHING MORREY SPACES

After studying Morrey spaces in detail, researchers have passed to generalized Morrey spaces. Mizuhara [31] has given generalized Morrey spaces  $M_{p, \varphi}$  considering  $\varphi = \varphi(r)$  instead of  $r^\lambda$  in the above definition of the Morrey space. Later, Guliyev [17] has defined the generalized Morrey spaces  $M_{p, \varphi}$  with normalized norm as follows:

**Definition 2.1.** [17] (**generalized Morrey space**) Let  $\varphi(x, r)$  be a positive measurable function on  $\mathbb{R}^n \times (0, \infty)$  and  $1 \leq p < \infty$ . We denote by  $M_{p,\varphi} \equiv M_{p,\varphi}(\mathbb{R}^n)$  the generalized Morrey space, the space of all functions  $f \in L_p^{loc}(\mathbb{R}^n)$  with finite quasinorm

$$\|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x,r))}.$$

Also by  $WM_{p,\varphi} \equiv WM_{p,\varphi}(\mathbb{R}^n)$  we denote the weak generalized Morrey space of all functions  $f \in WL_p^{loc}(\mathbb{R}^n)$  for which

$$\|f\|_{WM_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x,r))} < \infty.$$

According to this definition, we recover the Morrey space  $M_{p,\lambda}$  and weak Morrey space  $WM_{p,\lambda}$  under the choice  $\varphi(x, r) = r^{\frac{\lambda-n}{p}}$ :

$$M_{p,\lambda} = M_{p,\varphi} \Big|_{\varphi(x,r)=r^{\frac{\lambda-n}{p}}}, \quad WM_{p,\lambda} = WM_{p,\varphi} \Big|_{\varphi(x,r)=r^{\frac{\lambda-n}{p}}}.$$

For brevity, in the sequel we use the notations

$$\mathfrak{M}_{p,\varphi}(f; x, r) := \frac{|B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x,r))}}{\varphi(x, r)}$$

and

$$\mathfrak{M}_{p,\varphi}^W(f; x, r) := \frac{|B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x,r))}}{\varphi(x, r)}.$$

In this paper, extending the definition of vanishing Morrey spaces [50], we introduce the generalized vanishing Morrey spaces  $VM_{p,\varphi}(\mathbb{R}^n)$ , including their weak versions and studies the boundedness of the sublinear operators with rough kernel generated by fractional integrals and their commutators in these spaces. Indeed, we find it convenient to define generalized vanishing Morrey spaces in the form as follows.

**Definition 2.2.** (**generalized vanishing Morrey space**) The generalized vanishing Morrey space  $VM_{p,\varphi}(\mathbb{R}^n)$  is defined as the spaces of functions  $f \in M_{p,\varphi}(\mathbb{R}^n)$  such that

$$(2.1) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi}(f; x, r) = 0.$$

**Definition 2.3.** (**weak generalized vanishing Morrey space**) The weak generalized vanishing Morrey space  $WVM_{p,\varphi}(\mathbb{R}^n)$  is defined as the spaces of functions  $f \in WM_{p,\varphi}(\mathbb{R}^n)$  such that

$$(2.2) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi}^W(f; x, r) = 0.$$

Everywhere in the sequel we assume that

$$(2.3) \quad \lim_{r \rightarrow 0} \frac{1}{\inf_{x \in \mathbb{R}^n} \varphi(x, r)} = 0,$$

and

$$(2.4) \quad \sup_{0 < r < \infty} \frac{1}{\inf_{x \in \mathbb{R}^n} \varphi(x, r)} < \infty,$$

which make the spaces  $VM_{p,\varphi}(\mathbb{R}^n)$  and  $WVM_{p,\varphi}(\mathbb{R}^n)$  non-trivial, because bounded functions with compact support belong to this space. The spaces  $VM_{p,\varphi}(\mathbb{R}^n)$  and  $WVM_{p,\varphi}(\mathbb{R}^n)$  are Banach spaces with respect to the norm

$$(2.5) \quad \|f\|_{VM_{p,\varphi}} \equiv \|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \mathfrak{M}_{p,\varphi}(f; x, r),$$

$$(2.6) \quad \|f\|_{WVM_{p,\varphi}} = \|f\|_{WM_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \mathfrak{M}_{p,\varphi}^W(f; x, r),$$

respectively.

### 3. SUBLINEAR OPERATORS WITH ROUGH KERNEL $T_{\Omega,\alpha}$ ON THE SPACES $M_{p,\varphi}$ AND $VM_{p,\varphi}$

In this section, we will first prove the boundedness of the operator  $T_{\Omega,\alpha}$  satisfying (1.1) on the generalized Morrey spaces  $M_{p,\varphi}(\mathbb{R}^n)$  by using Lemma 1.2 and the following Lemma 3.1. Then, we will also give the boundedness of  $T_{\Omega,\alpha}$  satisfying (1.1) on generalized vanishing Morrey spaces  $VM_{p,\varphi}(\mathbb{R}^n)$ .

We first prove the following lemma (our main lemma).

**Lemma 3.1.** *(Our main lemma) Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let  $T_{\Omega,\alpha}$  be a sublinear operator satisfying condition (1.1), bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_q(\mathbb{R}^n)$  for  $p = 1$ .*

*If  $p > 1$  and  $s' \leq p$ , then the inequality*

$$(3.1) \quad \|T_{\Omega,\alpha}f\|_{L_q(B(x_0,r))} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))} dt$$

*holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .*

*If  $p > 1$  and  $q < s$ , then the inequality*

$$\|T_{\Omega,\alpha}f\|_{L_q(B(x_0,r))} \lesssim r^{\frac{n}{q}-\frac{n}{s}} \int_{2r}^{\infty} t^{\frac{n}{s}-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))} dt$$

*holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .*

*Moreover, for  $p = 1 < q < s$  the inequality*

$$(3.2) \quad \|T_{\Omega,\alpha}f\|_{WL_q(B(x_0,r))} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} t^{-\frac{n}{q}-1} \|f\|_{L_1(B(x_0,t))} dt$$

*holds for any ball  $B(x_0, r)$  and for all  $f \in L_1^{loc}(\mathbb{R}^n)$ .*

*Proof.* Let  $0 < \alpha < n$ ,  $1 \leq s' < p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Set  $B = B(x_0, r)$  for the ball centered at  $x_0$  and of radius  $r$  and  $2B = B(x_0, 2r)$ . We represent  $f$  as

$$(3.3) \quad f = f_1 + f_2, \quad f_1(y) = f(y) \chi_{2B}(y), \quad f_2(y) = f(y) \chi_{(2B)^c}(y), \quad r > 0$$

and have

$$\|T_{\Omega,\alpha}f\|_{L_q(B)} \leq \|T_{\Omega,\alpha}f_1\|_{L_q(B)} + \|T_{\Omega,\alpha}f_2\|_{L_q(B)}.$$

Since  $f_1 \in L_p(\mathbb{R}^n)$ ,  $T_{\Omega,\alpha}f_1 \in L_q(\mathbb{R}^n)$  and from the boundedness of  $T_{\Omega,\alpha}$  from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$  (see Lemma 1.1) it follows that:

$$\|T_{\Omega,\alpha}f_1\|_{L_q(B)} \leq \|T_{\Omega,\alpha}f_1\|_{L_q(\mathbb{R}^n)} \leq C \|f_1\|_{L_p(\mathbb{R}^n)} = C \|f\|_{L_p(2B)},$$

where constant  $C > 0$  is independent of  $f$ .

It is clear that  $x \in B$ ,  $y \in (2B)^C$  implies  $\frac{1}{2}|x_0 - y| \leq |x - y| \leq \frac{3}{2}|x_0 - y|$ . We get

$$|T_{\Omega,\alpha}f_2(x)| \leq 2^{n-\alpha}c_1 \int_{(2B)^C} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^{n-\alpha}} dy.$$

By the Fubini's theorem, we have

$$\begin{aligned} \int_{(2B)^C} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^{n-\alpha}} dy &\approx \int_{(2B)^C} |f(y)| |\Omega(x-y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1-\alpha}} dy \\ &\approx \int_{2r}^{\infty} \int_{2r \leq |x_0-y| \leq t} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}} \\ &\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}}. \end{aligned}$$

Applying the Hölder's inequality, we get

$$\begin{aligned} &\int_{(2B)^C} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^{n-\alpha}} dy \\ (3.4) \quad &\lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \|\Omega(x-\cdot)\|_{L_s(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{s}} \frac{dt}{t^{n+1-\alpha}}. \end{aligned}$$

For  $x \in B(x_0, t)$ , notice that  $\Omega$  is homogenous of degree zero and  $\Omega \in L_s(S^{n-1})$ ,  $s > 1$ . Then, we obtain

$$\begin{aligned}
\left( \int_{B(x_0, t)} |\Omega(x-y)|^s dy \right)^{\frac{1}{s}} &= \left( \int_{B(x-x_0, t)} |\Omega(z)|^s dz \right)^{\frac{1}{s}} \\
&\leq \left( \int_{B(0, t+|x-x_0|)} |\Omega(z)|^s dz \right)^{\frac{1}{s}} \\
&\leq \left( \int_{B(0, 2t)} |\Omega(z)|^s dz \right)^{\frac{1}{s}} \\
&= \left( \int_{S^{n-1}} \int_0^{2t} |\Omega(z')|^s d\sigma(z') r^{n-1} dr \right)^{\frac{1}{s}} \\
(3.5) \quad &= C \|\Omega\|_{L_s(S^{n-1})} |B(x_0, 2t)|^{\frac{1}{s}}.
\end{aligned}$$

Thus, by (3.5), it follows that:

$$|T_{\Omega, \alpha} f_2(x)| \lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

Moreover, for all  $p \in [1, \infty)$  the inequality

$$(3.6) \quad \|T_{\Omega, \alpha} f_2\|_{L_q(B)} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}$$

is valid. Thus, we obtain

$$\|T_{\Omega, \alpha} f\|_{L_q(B)} \lesssim \|f\|_{L_p(2B)} + r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

On the other hand, we have

$$\begin{aligned}
\|f\|_{L_p(2B)} &\approx r^{\frac{n}{q}} \|f\|_{L_p(2B)} \int_{2r}^{\infty} \frac{dt}{t^{\frac{n}{q}+1}} \\
(3.7) \quad &\leq r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}.
\end{aligned}$$

By combining the above inequalities, we obtain

$$\|T_{\Omega, \alpha} f\|_{L_q(B)} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

Let  $1 < q < s$ . Similarly to (3.5), when  $y \in B(x_0, t)$ , it is true that

$$(3.8) \quad \left( \int_{B(x_0, r)} |\Omega(x-y)|^s dy \right)^{\frac{1}{s}} \leq C \|\Omega\|_{L_s(S^{n-1})} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{s}}.$$

By the Fubini's theorem, the Minkowski inequality and (3.8), we get

$$\begin{aligned} \|T_{\Omega, \alpha} f_2\|_{L_q(B)} &\leq \left( \int_B \left| \int_{2r}^{\infty} \int_{B(x_0, t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}} \right|^q dx \right)^{\frac{1}{q}} \\ &\leq \int_{2r}^{\infty} \int_{B(x_0, t)} |f(y)| \|\Omega(\cdot - y)\|_{L_q(B)} dy \frac{dt}{t^{n+1-\alpha}} \\ &\leq |B(x_0, r)|^{\frac{1}{q} - \frac{1}{s}} \int_{2r}^{\infty} \int_{B(x_0, t)} |f(y)| \|\Omega(\cdot - y)\|_{L_s(B)} dy \frac{dt}{t^{n+1-\alpha}} \\ &\lesssim r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \|f\|_{L_1(B(x_0, t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{s}} \frac{dt}{t^{n+1-\alpha}} \\ &\lesssim r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} t^{\frac{n}{s} - \frac{n}{q} - 1} dt. \end{aligned}$$

Let  $p = 1 < q < s \leq \infty$ . From the weak  $(1, q)$  boundedness of  $T_{\Omega, \alpha}$  and (3.7) it follows that:

$$(3.9) \quad \begin{aligned} \|T_{\Omega, \alpha} f_1\|_{WL_q(B)} &\leq \|T_{\Omega, \alpha} f_1\|_{WL_q(\mathbb{R}^n)} \lesssim \|f_1\|_{L_1(\mathbb{R}^n)} \\ &= \|f\|_{L_1(2B)} \lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_1(B(x_0, t))} \frac{dt}{t^{\frac{n}{q}+1}}. \end{aligned}$$

Then from (3.6) and (3.9) we get the inequality (3.2), which completes the proof.  $\square$

In the following theorem (our main result), we get the boundedness of the operator  $T_{\Omega, \alpha}$  on the generalized Morrey spaces  $M_{p, \varphi}$ .

**Theorem 3.1.** *(Our main result) Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let  $T_{\Omega, \alpha}$  be a sublinear operator satisfying condition (1.1), bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_q(\mathbb{R}^n)$  for  $p = 1$ . Let also, for  $s' \leq p < q$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies the condition*

$$(3.10) \quad \int_r^{\infty} \frac{\text{essinf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q}+1}} dt \leq C \varphi_2(x, r),$$



and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(3.11) \quad \int_r^\infty \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt \leq C \varphi_2(x, r) r^{\frac{n}{s}},$$

where  $C$  does not depend on  $x$  and  $r$ .

Then the operator  $T_{\Omega, \alpha}$  is bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$  for  $p > 1$  and from  $M_{1, \varphi_1}$  to  $WM_{q, \varphi_2}$  for  $p = 1$ . Moreover, we have for  $p > 1$

$$\|T_{\Omega, \alpha} f\|_{M_{q, \varphi_2}} \lesssim \|f\|_{M_{p, \varphi_1}},$$

and for  $p = 1$

$$\|T_{\Omega, \alpha} f\|_{WM_{q, \varphi_2}} \lesssim \|f\|_{M_{1, \varphi_1}}.$$

*Proof.* Since  $f \in M_{p, \varphi_1}$ , by (2.6) and the non-decreasing, with respect to  $t$ , of the norm  $\|f\|_{L_p(B(x_0, t))}$ , we get

$$\begin{aligned} & \frac{\|f\|_{L_p(B(x_0, t))}}{\operatorname{ess\,inf}_{0 < t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \operatorname{ess\,sup}_{0 < t < \tau < \infty} \frac{\|f\|_{L_p(B(x_0, t))}}{\varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \operatorname{ess\,sup}_{0 < \tau < \infty} \frac{\|f\|_{L_p(B(x_0, \tau))}}{\varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \|f\|_{M_{p, \varphi_1}}. \end{aligned}$$

For  $s' \leq p < \infty$ , since  $(\varphi_1, \varphi_2)$  satisfies (3.10), we have

$$\begin{aligned} & \int_r^\infty \|f\|_{L_p(B(x_0, t))} t^{-\frac{n}{q}} \frac{dt}{t} \\ & \leq \int_r^\infty \frac{\|f\|_{L_p(B(x_0, t))}}{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q}}} \frac{dt}{t} \\ & \leq C \|f\|_{M_{p, \varphi_1}} \int_r^\infty \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q}}} \frac{dt}{t} \\ & \leq C \|f\|_{M_{p, \varphi_1}} \varphi_2(x_0, r). \end{aligned}$$

Then by (3.1), we get

$$\begin{aligned} \|T_{\Omega, \alpha} f\|_{M_{q, \varphi_2}} &= \sup_{x_0 \in \mathbb{R}^n, r > 0} \varphi_2(x_0, r)^{-1} |B(x_0, r)|^{-\frac{1}{q}} \|T_{\Omega, \alpha} f\|_{L_q(B(x_0, r))} \\ &\leq C \sup_{x_0 \in \mathbb{R}^n, r > 0} \varphi_2(x_0, r)^{-1} \int_r^\infty \|f\|_{L_p(B(x_0, t))} t^{-\frac{n}{q}} \frac{dt}{t} \\ &\leq C \|f\|_{M_{p, \varphi_1}}. \end{aligned}$$

For the case of  $p = 1 < q < s$ , we can also use the same method, so we omit the details. This completes the proof of Theorem 3.1.  $\square$

In the case of  $q = \infty$  by Theorem 3.1, we get

**Corollary 3.1.** *Let  $1 \leq p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10). Then the operators  $M_\alpha$  and  $\bar{T}_\alpha$  are bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$  for  $p > 1$  and from  $M_{1, \varphi_1}$  to  $WM_{q, \varphi_2}$  for  $p = 1$ .*

**Corollary 3.2.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let also for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.11). Then the operators  $M_{\Omega, \alpha}$  and  $\bar{T}_{\Omega, \alpha}$  are bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$  for  $p > 1$  and from  $M_{1, \varphi_1}$  to  $WM_{q, \varphi_2}$  for  $p = 1$ .*

Now using above results, we get the boundedness of the operator  $T_{\Omega, \alpha}$  on the generalized vanishing Morrey spaces  $VM_{p, \varphi}$ .

**Theorem 3.2.** *(Our main result) Let  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , be homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let  $T_{\Omega, \alpha}$  be a sublinear operator satisfying condition (1.1), bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$ . Let for  $s' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and*

$$(3.12) \quad c_\delta := \int_{\delta}^{\infty} \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q}+1}} dt < \infty$$

for every  $\delta > 0$ , and

$$(3.13) \quad \int_r^{\infty} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q}+1}} dt \leq C_0 \varphi_2(x, r),$$

and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and also

$$(3.14) \quad c_{\delta'} := \int_{\delta'}^{\infty} \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt < \infty$$

for every  $\delta' > 0$ , and

$$(3.15) \quad \int_r^{\infty} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt \leq C_0 \varphi_2(x, r) r^{\frac{n}{s}},$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$ .

Then the operator  $T_{\Omega, \alpha}$  is bounded from  $VM_{p, \varphi_1}$  to  $VM_{q, \varphi_2}$  for  $p > 1$  and from  $M_{1, \varphi_1}$  to  $WVM_{q, \varphi_2}$  for  $p = 1$ . Moreover, we have for  $p > 1$

$$(3.16) \quad \|T_{\Omega, \alpha} f\|_{VM_{q, \varphi_2}} \lesssim \|f\|_{VM_{p, \varphi_1}},$$

and for  $p = 1$

$$(3.17) \quad \|T_{\Omega, \alpha} f\|_{WVM_{q, \varphi_2}} \lesssim \|f\|_{VM_{1, \varphi_1}}.$$

*Proof.* The norm inequalities follow from Theorem 3.1. Thus we only have to prove that

$$(3.18) \quad \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p, \varphi_1}(f; x, r) = 0 \text{ implies } \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{q, \varphi_2}(T_{\Omega, \alpha} f; x, r) = 0$$

and

$$(3.19) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p, \varphi_1}(f; x, r) = 0 \text{ implies } \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{q, \varphi_2}^W(T_{\Omega, \alpha} f; x, r) = 0.$$

To show that  $\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{q}} \|T_{\Omega, \alpha} f\|_{L_q(B(x, r))}}{\varphi_2(x, r)} < \epsilon$  for small  $r$ , we split the right-hand side of (3.1):

$$(3.20) \quad \frac{r^{-\frac{n}{q}} \|T_{\Omega, \alpha} f\|_{L_q(B(x, r))}}{\varphi_2(x, r)} \leq C [I_{\delta_0}(x, r) + J_{\delta_0}(x, r)],$$

where  $\delta_0 > 0$  (we may take  $\delta_0 < 1$ ), and

$$I_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_r^{\delta_0} t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x, t))} dt,$$

and

$$J_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_{\delta_0}^{\infty} t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x, t))} dt,$$

and  $r < \delta_0$ . Now we use the fact that  $f \in VM_{p, \varphi_1}$  and we choose any fixed  $\delta_0 > 0$  such that

$$\sup_{x \in \mathbb{R}^n} \frac{t^{-\frac{n}{p}} \|f\|_{L_p(B(x, t))}}{\varphi_1(x, t)} < \frac{\epsilon}{2CC_0}, \quad t \leq \delta_0,$$

where  $C$  and  $C_0$  are constants from (3.13) and (3.20). This allows to estimate the first term uniformly in  $r \in (0, \delta_0)$ :

$$\sup_{x \in \mathbb{R}^n} CI_{\delta_0}(x, r) < \frac{\epsilon}{2}, \quad 0 < r < \delta_0.$$

The estimation of the second term may be obtained by choosing  $r$  sufficiently small. Indeed, we have

$$J_{\delta_0}(x, r) \leq c_{\delta_0} \frac{\|f\|_{M_{p, \varphi_1}}}{\varphi_2(x, r)},$$

where  $c_{\delta_0}$  is the constant from (3.12) with  $\delta = \delta_0$ . Then, by (2.3) it suffices to choose  $r$  small enough such that

$$\sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_2(x, r)} \leq \frac{\epsilon}{2c_{\delta_0} \|f\|_{M_{p, \varphi_1}}},$$

which completes the proof of (3.18).

The proof of (3.19) is similar to the proof of (3.18). For the case of  $q < s$ , we can also use the same method, so we omit the details.  $\square$

*Remark 3.1.* Conditions (3.12) and (3.14) are not needed in the case when  $\varphi(x, r)$  does not depend on  $x$ , since (3.12) follows from (3.13) and similarly, (3.14) follows from (3.15) in this case.

**Corollary 3.3.** *Let  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , be homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let also for  $s' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.12)-(3.13) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.14)-(3.15). Then the operators*

$M_{\Omega,\alpha}$  and  $\bar{T}_{\Omega,\alpha}$  are bounded from  $VM_{p,\varphi_1}$  to  $VM_{q,\varphi_2}$  for  $p > 1$  and from  $VM_{1,\varphi_1}$  to  $WVM_{q,\varphi_2}$  for  $p = 1$ .

In the case of  $q = \infty$  by Theorem 3.2, we get

**Corollary 3.4.** *Let  $1 \leq p < \infty$  and the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.12)-(3.13). Then the operators  $M_\alpha$  and  $\bar{T}_\alpha$  are bounded from  $VM_{p,\varphi_1}$  to  $VM_{q,\varphi_2}$  for  $p > 1$  and from  $VM_{1,\varphi_1}$  to  $WVM_{q,\varphi_2}$  for  $p = 1$ .*

#### 4. COMMUTATORS OF THE SUBLINEAR OPERATORS WITH ROUGH KERNEL $T_{\Omega,\alpha}$ ON THE SPACES $M_{p,\varphi}$ AND $VM_{p,\varphi}$

In this section, we will first prove the boundedness of the operator  $T_{\Omega,b,\alpha}$  satisfying (1.2) with  $b \in BMO(\mathbb{R}^n)$  on the generalized Morrey spaces  $M_{p,\varphi}$  by using Lemma 1.2 and the following Lemma 4.1. Then, we will also obtain the boundedness of  $T_{\Omega,b,\alpha}$  satisfying (1.2) with  $b \in BMO(\mathbb{R}^n)$  on generalized vanishing Morrey spaces  $VM_{p,\varphi}$ .

Let  $T$  be a linear operator. For a locally integrable function  $b$  on  $\mathbb{R}^n$ , we define the commutator  $[b, T]$  by

$$[b, T]f(x) = b(x)Tf(x) - T(bf)(x)$$

for any suitable function  $f$ . Let  $\bar{T}$  be a C-Z operator. A well known result of Coifman et al. [9] states that when  $K(x) = \frac{\Omega(x')}{|x|^n}$  and  $\Omega$  is smooth, the commutator  $[b, \bar{T}]f = b\bar{T}f - \bar{T}(bf)$  is bounded on  $L_p(\mathbb{R}^n)$ ,  $1 < p < \infty$ , if and only if  $b \in BMO(\mathbb{R}^n)$ . The commutator of C-Z operators plays an important role in studying the regularity of solutions of elliptic partial differential equations of second order (see, for example, [7, 8, ?]). The boundedness of the commutator has been generalized to other contexts and important applications to some non-linear PDEs have been given by Coifman et al. [10]. On the other hand, For  $b \in L_1^{loc}(\mathbb{R}^n)$ , the commutator  $[b, \bar{T}_\alpha]$  of fractional integral operator (also known as the Riesz potential) is defined by

$$[b, \bar{T}_\alpha]f(x) = b(x)\bar{T}_\alpha f(x) - \bar{T}_\alpha(bf)(x) = \int_{\mathbb{R}^n} \frac{b(x) - b(y)}{|x - y|^{n-\alpha}} f(y) dy \quad 0 < \alpha < n$$

for any suitable function  $f$ .

The function  $b$  is also called the symbol function of  $[b, \bar{T}_\alpha]$ . The characterization of  $(L_p, L_q)$ -boundedness of the commutator  $[b, \bar{T}_\alpha]$  of fractional integral operator has been given by Chanillo [4]. A well known result of Chanillo [4] states that the commutator  $[b, \bar{T}_\alpha]$  is bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ ,  $1 < p < q < \infty$ ,  $\frac{1}{p} - \frac{1}{q} = \frac{\alpha}{n}$  if and only if  $b \in BMO(\mathbb{R}^n)$ . There are two major reasons for considering the problem of commutators. The first one is that the boundedness of commutators can produce some characterizations of function spaces (see [2, 4, 18, 19, 20, 21, 37, 42]). The other one is that the theory of commutators plays an important role in the study of the regularity of solutions to elliptic and parabolic PDEs of the second order (see [7, 8, 14, 41, 43]).

Let us recall the definition of the space of  $BMO(\mathbb{R}^n)$ .

**Definition 4.1.** Suppose that  $b \in L_1^{loc}(\mathbb{R}^n)$ , let

$$\|b\|_* = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}| dy < \infty,$$

where

$$b_{B(x,r)} = \frac{1}{|B(x,r)|} \int_{B(x,r)} b(y) dy.$$

Define

$$BMO(\mathbb{R}^n) = \{b \in L_1^{loc}(\mathbb{R}^n) : \|b\|_* < \infty\}.$$

If one regards two functions whose difference is a constant as one, then the space  $BMO(\mathbb{R}^n)$  is a Banach space with respect to norm  $\|\cdot\|_*$ .

*Remark 4.1.* [23] (1) The John-Nirenberg inequality [22]: there are constants  $C_1, C_2 > 0$ , such that for all  $b \in BMO(\mathbb{R}^n)$  and  $\beta > 0$

$$|\{x \in B : |b(x) - b_B| > \beta\}| \leq C_1 |B| e^{-C_2 \beta / \|b\|_*}, \quad \forall B \subset \mathbb{R}^n.$$

(2) The John-Nirenberg inequality implies that

$$(4.1) \quad \|b\|_* \approx \sup_{x \in \mathbb{R}^n, r > 0} \left( \frac{1}{|B(x,r)|} \int_{B(x,r)} |b(y) - b_{B(x,r)}|^p dy \right)^{\frac{1}{p}}$$

for  $1 < p < \infty$ .

(3) Let  $b \in BMO(\mathbb{R}^n)$ . Then there is a constant  $C > 0$  such that

$$(4.2) \quad |b_{B(x,r)} - b_{B(x,t)}| \leq C \|b\|_* \ln \frac{t}{r} \text{ for } 0 < 2r < t,$$

where  $C$  is independent of  $b, x, r$  and  $t$ .

As in the proof of Theorem 3.1, it suffices to prove the following Lemma (our main lemma).

**Lemma 4.1.** (Our main lemma) Let  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , be homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ ,  $b \in BMO(\mathbb{R}^n)$ , and  $T_{\Omega,b,\alpha}$  is a sublinear operator satisfying condition (1.2) and bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ . Then, for  $s' \leq p$  the inequality

$$(4.3) \quad \|T_{\Omega,b,\alpha} f\|_{L_q(B(x_0,r))} \lesssim \|b\|_* r^{\frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x_0,t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .

Also, for  $q < s$  the inequality

$$\|T_{\Omega,b,\alpha} f\|_{L_q(B(x_0,r))} \lesssim \|b\|_* r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{s} - \frac{n}{q} - 1} \|f\|_{L_p(B(x_0,t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .

*Proof.* Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . As in the proof of Lemma 3.1, we represent  $f$  in form (3.3) and have

$$\|T_{\Omega,b,\alpha} f\|_{L_q(B)} \leq \|T_{\Omega,b,\alpha} f_1\|_{L_q(B)} + \|T_{\Omega,b,\alpha} f_2\|_{L_q(B)}.$$

From the boundedness of  $T_{\Omega,b,\alpha}$  from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$  (see Theorem 1.3) it follows that:

$$\begin{aligned} \|T_{\Omega,b,\alpha}f_1\|_{L_q(B)} &\leq \|T_{\Omega,b,\alpha}f_1\|_{L_q(\mathbb{R}^n)} \\ &\lesssim \|b\|_* \|f_1\|_{L_p(\mathbb{R}^n)} = \|b\|_* \|f\|_{L_p(2B)}. \end{aligned}$$

It is known that  $x \in B$ ,  $y \in (2B)^C$ , which implies  $\frac{1}{2}|x_0 - y| \leq |x - y| \leq \frac{3}{2}|x_0 - y|$ . Then for  $x \in B$ , we have

$$\begin{aligned} |T_{\Omega,b,\alpha}f_2(x)| &\lesssim \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |b(y) - b(x)| |f(y)| dy \\ &\approx \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |b(y) - b(x)| |f(y)| dy. \end{aligned}$$

Hence we get

$$\begin{aligned} \|T_{\Omega,b,\alpha}f_2\|_{L_q(B)} &\lesssim \left( \int_B \left( \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |b(y) - b(x)| |f(y)| dy \right)^q dx \right)^{\frac{1}{q}} \\ &\lesssim \left( \int_B \left( \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |b(y) - b_B| |f(y)| dy \right)^q dx \right)^{\frac{1}{q}} \\ &\quad + \left( \int_B \left( \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |b(x) - b_B| |f(y)| dy \right)^q dx \right)^{\frac{1}{q}} \\ &= J_1 + J_2. \end{aligned}$$

We have the following estimation of  $J_1$ . When  $s' \leq p$  and  $\frac{1}{\mu} + \frac{1}{p} + \frac{1}{s} = 1$ , by the Fubini's theorem

$$\begin{aligned} J_1 &\approx r^{\frac{n}{q}} \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |b(y) - b_B| |f(y)| dy \\ &\approx r^{\frac{n}{q}} \int_{(2B)^C} |\Omega(x-y)| |b(y) - b_B| |f(y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1-\alpha}} dy \\ &\approx r^{\frac{n}{q}} \int_{2r}^{\infty} \int_{2r \leq |x_0-y| \leq t} |\Omega(x-y)| |b(y) - b_B| |f(y)| dy \frac{dt}{t^{n+1-\alpha}} \\ &\lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \int_{B(x_0,t)} |\Omega(x-y)| |b(y) - b_B| |f(y)| dy \frac{dt}{t^{n+1-\alpha}} \text{ holds.} \end{aligned}$$

Applying the Hölder's inequality and by (3.8), (4.1), (4.2), we get

$$\begin{aligned}
J_1 &\lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \int_{B(x_0,t)} |\Omega(x-y)| |b(y) - b_{B(x_0,t)}| |f(y)| dy \frac{dt}{t^{n+1-\alpha}} \\
&\quad + r^{\frac{n}{q}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |\Omega(x-y)| |f(y)| dy \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim r^{\frac{n}{q}} \int_{2r}^{\infty} \|\Omega(\cdot - y)\|_{L_s(B(x_0,t))} \| (b(\cdot) - b_{B(x_0,t)}) \|_{L_\mu(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1-\alpha}} \\
&\quad + r^{\frac{n}{q}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|\Omega(\cdot - y)\|_{L_s(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{s}} \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim \|b\|_* r^{\frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}}.
\end{aligned}$$

In order to estimate  $J_2$  note that

$$J_2 = \| (b(\cdot) - b_{B(x_0,t)}) \|_{L_q(B(x_0,t))} \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |f(y)| dy.$$

By (4.1), we get

$$J_2 \lesssim \|b\|_* r^{\frac{n}{q}} \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^{n-\alpha}} |f(y)| dy.$$

Thus, by (3.4) and (3.5)

$$J_2 \lesssim \|b\|_* r^{\frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

Summing up  $J_1$  and  $J_2$ , for all  $p \in (1, \infty)$  we get

$$(4.4) \quad \|T_{\Omega,b,\alpha} f\|_{L_q(B)} \lesssim \|b\|_* r^{\frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}}.$$

Finally, we have the following

$$\|T_{\Omega,b,\alpha} f\|_{L_q(B)} \lesssim \|b\|_* \|f\|_{L_p(2B)} + \|b\|_* r^{\frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{q}+1}},$$

which completes the proof of first statement by (3.7).

On the other hand when  $q < s$ , by the Fubini's theorem and the Minkowski inequality, we get

$$\begin{aligned}
J_1 &\lesssim \left( \int_B \left| \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}} \right|^q dx \right)^{\frac{1}{q}} \\
&+ \left( \int_B \left| \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}} \right|^q dx \right)^{\frac{1}{q}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} dy \frac{dt}{t^{n+1-\alpha}} \\
&+ \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} dy \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim |B|^{\frac{1}{q} - \frac{1}{s}} \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| \|\Omega(\cdot - y)\|_{L_s(B(x_0,t))} dy \frac{dt}{t^{n+1-\alpha}} \\
&+ |B|^{\frac{1}{q} - \frac{1}{s}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_s(B(x_0,t))} dy \frac{dt}{t^{n+1-\alpha}}.
\end{aligned}$$

Applying the Hölder's inequality and by (3.8), (4.1), (4.2), we get

$$\begin{aligned}
J_1 &\lesssim r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \|(b(\cdot) - b_{B(x_0,t)}) f\|_{L_1(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{s}} \frac{dt}{t^{n+1-\alpha}} \\
&+ r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|f\|_{L_p(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{s}} \frac{dt}{t^{\frac{n}{q}+1}} \\
&\lesssim r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \|(b(\cdot) - b_{B(x_0,t)})\|_{L_{p'}(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} t^{\frac{n}{s}} \frac{dt}{t^{n+1}} \\
&+ r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|f\|_{L_p(B(x_0,t))} t^{\frac{n}{s}} \frac{dt}{t^{\frac{n}{q}+1}} \\
&\lesssim \|b\|_* r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{s} - \frac{n}{q} - 1} \|f\|_{L_p(B(x_0,t))} dt.
\end{aligned}$$



Let  $\frac{1}{p} = \frac{1}{\nu} + \frac{1}{s}$ , then for  $J_2$ , by the Fubini's theorem, the Minkowski inequality, the Hölder's inequality and from (3.8), we get

$$\begin{aligned}
J_2 &\lesssim \left( \int_B \left| \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| |b(x) - b_B| |\Omega(x-y)| dy \frac{dt}{t^{n+1-\alpha}} \right|^q dx \right)^{\frac{1}{q}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|(b(\cdot) - b_B) \Omega(\cdot - y)\|_{L_q(B)} dy \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|b(\cdot) - b_B\|_{L_\nu(B)} \|\Omega(\cdot - y)\|_{L_s(B)} dy \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim \|b\|_* |B|^{\frac{1}{q} - \frac{1}{s}} \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_s(B)} dy \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim \|b\|_* r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \|f\|_{L_1(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{s}} \frac{dt}{t^{n+1-\alpha}} \\
&\lesssim \|b\|_* r^{\frac{n}{q} - \frac{n}{s}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{s} - \frac{n}{q} - 1} \|f\|_{L_p(B(x_0,t))} dt.
\end{aligned}$$

By combining the above estimates, we complete the proof of Lemma 4.1.  $\square$

Now we can give the following theorem (our main result).

**Theorem 4.1.** (Our main result) Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero and  $T_{\Omega, b, \alpha}$  is a sublinear operator satisfying condition (1.2) and bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ . Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ .

Let also, for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(4.5) \quad \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q} + 1}} dt \leq C \varphi_2(x, r),$$

and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(4.6) \quad \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt \leq C \varphi_2(x, r) r^{\frac{n}{s}},$$

where  $C$  does not depend on  $x$  and  $r$ .

Then, the operator  $T_{\Omega, b, \alpha}$  is bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$ . Moreover

$$\|T_{\Omega, b, \alpha} f\|_{M_{q, \varphi_2}} \lesssim \|b\|_* \|f\|_{M_{p, \varphi_1}}.$$

*Proof.* The statement of Theorem 4.1 follows by Lemma 1.2 and Lemma 4.1 in the same manner as in the proof of Theorem 3.1.  $\square$

By Theorem 4.1, we get the following new result.

**Corollary 4.1.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ . If for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition (4.5) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition (4.6). Then, the operators  $M_{\Omega, b, \alpha}$  and  $[b, \overline{T}_{\Omega, \alpha}]$  are bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$ .*

For the sublinear commutator of the fractional maximal operator is defined as follows

$$M_{b, \alpha}(f)(x) = \sup_{t > 0} |B(x, t)|^{-1 + \frac{\alpha}{n}} \int_{B(x, t)} |b(x) - b(y)| |f(y)| dy$$

by Theorem 4.1 we get the following new result.

**Corollary 4.2.** *Let  $0 < \alpha < n$ ,  $1 < p < \frac{n}{\alpha}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ ,  $b \in BMO(\mathbb{R}^n)$  and the pair  $(\varphi_1, \varphi_2)$  satisfies the condition (4.5). Then, the operators  $M_{b, \alpha}$  and  $[b, \overline{T}_{\alpha}]$  are bounded from  $M_{p, \varphi_1}$  to  $M_{q, \varphi_2}$ .*

Now using above results, we also obtain the boundedness of the operator  $T_{\Omega, b, \alpha}$  on the generalized vanishing Morrey spaces  $VM_{p, \varphi}$ .

**Theorem 4.2.** *(Our main result) Let  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , be homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ ,  $b \in BMO(\mathbb{R}^n)$ , and  $T_{\Omega, b, \alpha}$  is a sublinear operator satisfying condition (1.2) and bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$ . Let for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and*

$$(4.7) \quad \int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q}+1}} dt \leq C_0 \varphi_2(x, r),$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$ ,

$$(4.8) \quad \lim_{r \rightarrow 0} \frac{\ln \frac{1}{r}}{\inf_{x \in \mathbb{R}^n} \varphi_2(x, r)} = 0$$

and

$$(4.9) \quad c_\delta := \int_\delta^\infty (1 + \ln |t|) \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q}+1}} dt < \infty$$

for every  $\delta > 0$ , and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and also

$$(4.10) \quad \int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt \leq C_0 \varphi_2(x, r) r^{\frac{n}{s}},$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$ ,

$$\lim_{r \rightarrow 0} \frac{\ln \frac{1}{r}}{\inf_{x \in \mathbb{R}^n} \varphi_2(x, r)} = 0$$

and

$$(4.11) \quad c_{\delta'} := \int_{\delta'}^\infty (1 + \ln |t|) \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{q} - \frac{n}{s} + 1}} dt < \infty$$

for every  $\delta' > 0$ .

Then the operator  $T_{\Omega,b,\alpha}$  is bounded from  $VM_{p,\varphi_1}$  to  $VM_{q,\varphi_2}$ . Moreover,

$$(4.12) \quad \|T_{\Omega,b,\alpha}f\|_{VM_{q,\varphi_2}} \lesssim \|b\|_* \|f\|_{VM_{p,\varphi_1}}.$$

*Proof.* The norm inequality having already been provided by Theorem 4.1, we only have to prove the implication

$$(4.13) \quad \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|f\|_{L_p(B(x,r))}}{\varphi_1(x,r)} = 0 \text{ implies } \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{q}} \|T_{\Omega,b,\alpha}f\|_{L_q(B(x,r))}}{\varphi_2(x,r)} = 0.$$

To show that

$$\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{q}} \|T_{\Omega,b,\alpha}f\|_{L_q(B(x,r))}}{\varphi_2(x,r)} < \epsilon \text{ for small } r,$$

we use the estimate (4.3):

$$\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{q}} \|T_{\Omega,b,\alpha}f\|_{L_q(B(x,r))}}{\varphi_2(x,r)} \lesssim \frac{\|b\|_*}{\varphi_2(x,r)} \int_r^\infty \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x,t))} dt.$$

We take  $r < \delta_0$ , where  $\delta_0$  will be chosen small enough and split the integration:

$$(4.14) \quad \frac{r^{-\frac{n}{q}} \|T_{\Omega,b,\alpha}f\|_{L_q(B(x,r))}}{\varphi_2(x,r)} \leq C [I_{\delta_0}(x,r) + J_{\delta_0}(x,r)],$$

where  $\delta_0 > 0$  (we may take  $\delta_0 < 1$ ), and

$$I_{\delta_0}(x,r) := \frac{1}{\varphi_2(x,r)} \int_r^{\delta_0} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x,t))} dt,$$

and

$$J_{\delta_0}(x,r) := \frac{1}{\varphi_2(x,r)} \int_{\delta_0}^\infty \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{q}-1} \|f\|_{L_p(B(x,t))} dt$$

Now we choose any fixed  $\delta_0 > 0$  such that

$$\sup_{x \in \mathbb{R}^n} \frac{t^{-\frac{n}{p}} \|f\|_{L_p(B(x,t))}}{\varphi_1(x,t)} < \frac{\epsilon}{2CC_0}, \quad t \leq \delta_0,$$

where  $C$  and  $C_0$  are constants from (4.7) and (4.14). This allows to estimate the first term uniformly in  $r \in (0, \delta_0)$ :

$$\sup_{x \in \mathbb{R}^n} CI_{\delta_0}(x,r) < \frac{\epsilon}{2}, \quad 0 < r < \delta_0.$$

For the second term, writing  $1 + \ln \frac{t}{r} \leq 1 + |\ln t| + \ln \frac{1}{r}$ , we obtain

$$J_{\delta_0}(x,r) \leq \frac{c_{\delta_0} + \widetilde{c}_{\delta_0} \ln \frac{1}{r}}{\varphi_2(x,r)} \|f\|_{M_{p,\varphi}},$$

where  $c_{\delta_0}$  is the constant from (4.9) with  $\delta = \delta_0$  and  $\widetilde{c}_{\delta_0}$  is a similar constant with omitted logarithmic factor in the integrand. Then, by (4.8) we can choose small enough  $r$  such that

$$\sup_{x \in \mathbb{R}^n} J_{\delta_0}(x,r) < \frac{\epsilon}{2},$$

which completes the proof of (4.13).

For the case of  $q < s$ , we can also use the same method, so we omit the details.  $\square$

*Remark 4.2.* Conditions (4.9) and (4.11) are not needed in the case when  $\varphi(x, r)$  does not depend on  $x$ , since (4.9) follows from (4.7) and similarly, (4.11) follows from (4.10) in this case.

**Corollary 4.3.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ . If for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.9)-(4.7) and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.11)-(4.10). Then, the operators  $M_{\Omega, b, \alpha}$  and  $[b, \overline{T}_{\Omega, \alpha}]$  are bounded from  $VM_{p, \varphi_1}(\mathbb{R}^n)$  to  $VM_{q, \varphi_2}(\mathbb{R}^n)$ .*

In the case of  $q = \infty$  by Theorem 4.2, we get

**Corollary 4.4.** *Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$  and the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.9)-(4.7). Then the operators  $M_{b, \alpha}$  and  $[b, \overline{T}_{\alpha}]$  are bounded from  $VM_{p, \varphi_1}(\mathbb{R}^n)$  to  $VM_{q, \varphi_2}(\mathbb{R}^n)$ .*

## 5. SOME APPLICATIONS

In this section, we give the applications of Theorem 3.1, Theorem 3.2, Theorem 4.1, Theorem 4.2 for the Marcinkiewicz operator.

**5.1. Marcinkiewicz Operator.** Let  $S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$  be the unit sphere in  $\mathbb{R}^n$  equipped with the Lebesgue measure  $d\sigma$ . Suppose that  $\Omega$  satisfies the following conditions.

(a)  $\Omega$  is the homogeneous function of degree zero on  $\mathbb{R}^n \setminus \{0\}$ , that is,

$$\Omega(\mu x) = \Omega(x), \text{ for any } \mu > 0, x \in \mathbb{R}^n \setminus \{0\}.$$

(b)  $\Omega$  has mean zero on  $S^{n-1}$ , that is,

$$\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0,$$

where  $x' = \frac{x}{|x|}$  for any  $x \neq 0$ .

(c)  $\Omega \in Lip_{\gamma}(S^{n-1})$ ,  $0 < \gamma \leq 1$ , that is there exists a constant  $M > 0$  such that,

$$|\Omega(x') - \Omega(y')| \leq M|x' - y'|^{\gamma} \text{ for any } x', y' \in S^{n-1}.$$

In 1958, Stein [45] defined the Marcinkiewicz integral of higher dimension  $\mu_{\Omega}$  as

$$\mu_{\Omega}(f)(x) = \left( \int_0^{\infty} |F_{\Omega, t}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega, t}(f)(x) = \int_{|x-y| \leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy.$$

Since Stein's work in 1958, the continuity of Marcinkiewicz integral has been extensively studied as a research topic and also provides useful tools in harmonic analysis [29, 46, 47, 48].

The Marcinkiewicz operator is defined by (see [49])

$$\mu_{\Omega,\alpha}(f)(x) = \left( \int_0^\infty |F_{\Omega,\alpha,t}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,\alpha,t}(f)(x) = \int_{|x-y|\leq t} \frac{\Omega(x-y)}{|x-y|^{n-1-\alpha}} f(y) dy.$$

Note that  $\mu_{\Omega}f = \mu_{\Omega,0}f$ .

The sublinear commutator of the operator  $\mu_{\Omega,\alpha}$  is defined by

$$[b, \mu_{\Omega,\alpha}](f)(x) = \left( \int_0^\infty |F_{\Omega,\alpha,t,b}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,\alpha,t,b}(f)(x) = \int_{|x-y|\leq t} \frac{\Omega(x-y)}{|x-y|^{n-1-\alpha}} [b(x) - b(y)] f(y) dy.$$

We consider the space  $H = \{h : \|h\| = (\int_0^\infty |h(t)|^2 \frac{dt}{t^3})^{1/2} < \infty\}$ . Then, it is clear that  $\mu_{\Omega,\alpha}(f)(x) = \|F_{\Omega,\alpha,t}(x)\|$ .

By the Minkowski inequality, we get

$$\mu_{\Omega,\alpha}(f)(x) \leq \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-1-\alpha}} |f(y)| \left( \int_{|x-y|}^\infty \frac{dt}{t^3} \right)^{1/2} dy \leq C \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-\alpha}} |f(y)| dy.$$

Thus,  $\mu_{\Omega,\alpha}$  satisfies the condition (1.1). It is known that for  $b \in BMO(\mathbb{R}^n)$  the operators  $\mu_{\Omega,\alpha}$  and  $[b, \mu_{\Omega,\alpha}]$  are bounded from  $L_p(\mathbb{R}^n)$  to  $L_q(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_q(\mathbb{R}^n)$  for  $p = 1$  (see [49]), then by Theorems 3.1, 3.2, 4.1 and 4.2 we get

**Corollary 5.1.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let also, for  $s' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.11) and  $\Omega$  satisfies conditions (a)–(c). Then the operator  $\mu_{\Omega,\alpha}$  is bounded from  $M_{p,\varphi_1}$  to  $M_{q,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{q,\varphi_2}$  for  $p = 1$ .*

**Corollary 5.2.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $0 < \alpha < n$ ,  $1 \leq p < \frac{n}{\alpha}$  and  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$ . Let also, for  $s' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4) and (3.12)–(3.13) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4) and (3.14)–(3.15) and  $\Omega$  satisfies conditions (a)–(c). Then the operator  $\mu_{\Omega,\alpha}$  is bounded from  $VM_{p,\varphi_1}$  to  $VM_{q,\varphi_2}$  for  $p > 1$  and from  $VM_{1,\varphi_1}$  to  $WVM_{q,\varphi_2}$  for  $p = 1$ .*

**Corollary 5.3.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ . Let also, for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.5) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.6) and  $\Omega$  satisfies conditions (a)–(c). Then, the operator  $[b, \mu_{\Omega,\alpha}]$  is bounded from  $M_{p,\varphi_1}$  to  $M_{q,\varphi_2}$ .*

**Corollary 5.4.** *Suppose that  $\Omega \in L_s(S^{n-1})$ ,  $1 < s \leq \infty$ , is homogeneous of degree zero. Let  $1 < p < \infty$ ,  $0 < \alpha < \frac{n}{p}$ ,  $\frac{1}{q} = \frac{1}{p} - \frac{\alpha}{n}$  and  $b \in BMO(\mathbb{R}^n)$ . Let also, for  $s' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.9)-(4.7) and for  $q < s$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.11)-(4.10) and  $\Omega$  satisfies conditions (a)-(c). Then, the operator  $[b, \mu_{\Omega, \alpha}]$  is bounded from  $VM_{p, \varphi_1}$  to  $VM_{q, \varphi_2}$ .*

**Acknowledgement:** This study has been given as the plenary talk by the author at the “3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference (IFSCOM2016), August 29-September 1, 2016, Mersin, Turkey”.

#### REFERENCES

- [1] D.R. Adams, A note on Riesz potentials, *Duke Math. J.*, 42 (1975), 765-778.
- [2] A.S. Balakishiyev, V.S. Guliyev, F. Gurbuz and A. Serbetci, Sublinear operators with rough kernel generated by Calderon-Zygmund operators and their commutators on generalized local Morrey spaces, *Journal of Inequalities and Applications* 2015, 2015:61. doi:10.1186/s13660-015-0582-y.
- [3] X.N. Cao, D.X. Chen, The boundedness of Toeplitz-type operators on vanishing Morrey spaces. *Anal. Theory Appl.* 27 (2011), 309-319.
- [4] S. Chanillo, A note on commutators, *Indiana Univ. Math. J.*, 31 (1), (1982), 7-16.
- [5] S. Chanillo, D.K. Watson, R.L. Wheeden, Some integral and maximal operators related to starlike sets. *Studia Math.*, 107 (1993), 223-255.
- [6] F. Chiarenza, M. Frasca, Morrey spaces and Hardy-Littlewood maximal function, *Rend. Mat.*, 7 (1987), 273-279.
- [7] F. Chiarenza, M. Frasca, P. Longo, Interior  $W^{2,p}$ -estimates for nondivergence elliptic equations with discontinuous coefficients, *Ricerche Mat.*, 40 (1991), 149-168.
- [8] F. Chiarenza, M. Frasca, P. Longo,  $W^{2,p}$ -solvability of Dirichlet problem for nondivergence elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.*, 336 (1993), 841-853.
- [9] R.R. Coifman, R. Rochberg, G. Weiss, Factorization theorems for Hardy spaces in several variables, *Ann. of Math.*, 103 (3) (1976), 611-635.
- [10] R.R. Coifman, P. Lions, Y. Meyer, S. Semmes, Compensated compactness and Hardy spaces. *J. Math. Pures Appl.* 72 (1993), 247-286.
- [11] R.R. Coifman, Y. Meyer, Au delà des Opérateurs Pseudo-Différentiels, *Astérisque* 57. Société Mathématique de France, Paris, 1978, 185 pp.
- [12] Y. Ding and S.Z. Lu, Higher order commutators for a class of rough operators, *Ark. Mat.*, 37 (1999), 33-44.
- [13] Y. Ding and S.Z. Lu, Homogeneous fractional integrals on Hardy spaces, *Tohoku Math. J.*, 52 (2000), 153-162.
- [14] G. Di Fazio, D.K. Palagachev and M.A. Ragusa, Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients, *J. Funct. Anal.*, 166 (1999), 179-196.
- [15] J. Garcia-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Math. 16, Amsterdam, 1985.
- [16] L. Grafakos, *Classical and modern Fourier analysis*. Pearson Education. Inc. Upper Saddle River, New Jersey, 2004.
- [17] V.S. Guliyev, Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces, *J. Inequal. Appl.* 2009, Art. ID 503948, 20 pp.
- [18] F. Gurbuz, Boundedness of some potential type sublinear operators and their commutators with rough kernels on generalized local Morrey spaces [*Ph.D. thesis*], Ankara University, Ankara, Turkey, 2015.
- [19] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic Calderón-Zygmund operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, *Open Math.*, 14 (2016), 300-323.
- [20] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic fractional integral operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, *Advan. Math. (China)*, in press.

- [21] S. Janson, Mean oscillation and commutators of singular integral operators, *Ark. Mat.*, 16 (1978), 263-270.
- [22] F. John and L. Nirenberg, On functions of bounded mean oscillation. *Comm. Pure Appl. Math.*, 14 (1961), 415-426.
- [23] T. Karaman, Boundedness of some classes of sublinear operators on generalized weighted Morrey spaces and some applications [*Ph.D. thesis*], Ankara University, Ankara, Turkey, 2012 (in Turkish).
- [24] T. Kato, Strong  $L^p$  solutions of the Navier-Stokes equations in  $\mathbb{R}^m$  with applications to weak solutions, *Math. Z.*, 187 (1984), 471-480.
- [25] X. Li and D.C. Yang, Boundedness of some sublinear operators on Herz spaces, *Illinois J. of Math.*, 40 (1996), 484-501.
- [26] Y. Liang and W. Su, The relationship between fractal dimensions of a type of fractal functions and the order of their fractional calculus, *Chaos, Solitons and Fractals*, 34 (2007), 682-692.
- [27] G. Lu, S.Z. Lu, D.C. Yang, Singular integrals and commutators on homogeneous groups, *Anal. Math.*, 28 (2002), 103-134.
- [28] S.Z. Lu, D. Yang, Z. Zhou, Sublinear operators with rough kernel on generalized Morrey spaces. *Hokkaido Math. J.* 27 (1998), 219-232.
- [29] S.Z. Lu, Y. Ding, D.Y. Yan, *Singular integrals and related topics*, World Scientific Publishing, Singapore, 2006.
- [30] C. Miranda, Sulle equazioni ellittiche del secondo ordine di tipo non variazionale, a coefficienti discontinui. *Ann. Math. Pura E Appl.* 63 (4) (1963), 353-386.
- [31] T. Mizuhara, Boundedness of some classical operators on generalized Morrey spaces, *Harmonic Analysis* (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer - Verlag, Tokyo (1991), 183-189.
- [32] C.B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.*, 43 (1938), 126-166.
- [33] B. Muckenhoupt, On certain singular integrals. *Pacif. J. Math.*, 10 (1960), 239-261.
- [34] B. Muckenhoupt, R.L. Wheeden, Weighted norm inequalities for singular and fractional integrals. *Trans. Amer. Math. Soc.*, 161 (1971), 249-258.
- [35] K. Oldham and J. Spanier, *The fractional calculus*, New York, Academic press, 1974.
- [36] D.K. Palagachev, L.G. Softova, Singular integral operators, Morrey spaces and fine regularity of solutions to PDE's, *Potential Anal.*, 20 (2004), 237-263.
- [37] M. Paluszynski, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss, *Indiana Univ. Math. J.*, 44 (1995), 1-17.
- [38] J. Peetre, On the theory of  $M_{p,\lambda}$ , *J. Funct. Anal.*, 4 (1969), 71-87.
- [39] L.E. Persson, M.A. Ragusa, N. Samko, P. Wall, Commutators of Hardy operators in vanishing Morrey spaces. *AIP Conf. Proc.* 1493, 859 (2012); <http://dx.doi.org/10.1063/1.4765588>.
- [40] M.A., Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces. *J. Global Optim.* 368 (40) (2008), 1-3.
- [41] S.G. Shi, Z.W. Fu, F.Y. Zhao, Estimates for operators on weighted Morrey spaces and their applications to nondivergence elliptic equations, *J. Inequal. Appl.* 2013, 2013: 390.
- [42] S.G. Shi, S.Z. Lu, A characterization of Campanato space via commutator of fractional integral, *J. Math. Anal. Appl.*, 419 (2014), 123-137.
- [43] L.G. Softova, Singular integrals and commutators in generalized Morrey spaces, *Acta Math. Sin. (Engl. Ser.)*, 22 (2006), 757-766.
- [44] F. Soria, G. Weiss, A remark on singular integrals and power weights, *Indiana Univ. Math. J.*, 43 (1994) 187-204.
- [45] E.M. Stein, On the functions of Littlewood-Paley, Lusin and Marcinkiewicz, *Trans. Amer. Math. Soc.* 88 (1958) 430-466.
- [46] E.M. Stein, *Singular integrals and differentiability of functions*, Princeton University Press, Princeton, NJ, 1970.
- [47] E.M. Stein, *Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton NJ, 1993.
- [48] A. Torchinsky, *Real Variable Methods in Harmonic Analysis*, Pure and Applied Math. 123, Academic Press, New York, 1986.
- [49] A. Torchinsky and S. Wang, A note on the Marcinkiewicz integral, *Colloq. Math.* 60/61 (1990), 235-243.

- [50] C. Vitanza, Functions with vanishing Morrey norm and elliptic partial differential equations. In: Proceedings of Methods of Real Analysis and Partial Differential Equations, Capri, pp. 147-150. Springer (1990).
- [51] C. Vitanza, Regularity results for a class of elliptic equations with coefficients in Morrey spaces. *Ricerche Mat.* 42 (2) (1993), 265-281.
- [52] R.L. Wheeden and A. Zygmund, *Measure and Integral: An Introduction to Real Analysis*, vol. 43 of Pure and Applied Mathematics, Marcel Dekker, New York, NY, USA, 1977.

ANKARA UNIVERSITY, FACULTY OF SCIENCE, DEPARTMENT OF MATHEMATICS, TANDOĞAN 06100,  
ANKARA, TURKEY

*E-mail address:* feritgurbuz84@hotmail.com



IFSCOM2016 <sup>1</sup> PROCEEDING BOOK

No. 1 PP. 37-41 (2016)

ISBN: 978-975-6900-54-3

## TEACHERS' OPINIONS ABOUT MATHEMATIC PROGRAM REVISED WITH 4+4+4 EDUCATION SYSTEM

ADEM BAŞKAYA AND ORKUN ÇOŞKUNTUNCEL

**ABSTRACT.** The purpose of this research is taking the teachers' opinions about the re-prepared secondary school (5, 6, 7 and 8 grades) mathematics program with gradually changing education system in 2013-2014 academic year. The universe of research is secondary school mathematics teachers working in Mersin province, and the sample of research contains 149 teachers working in city center, districts and villages and all agree to participate on a voluntary basis to the research.

The scale of the research containing 5 factors (General features, activity, and applications of mathematics course, new education system, and textbook) with 28 questions has been developed by the researchers and the Cronbach Alfa rate of the scale is 0,886. For the analysis of the data, descriptive statistics, t-test and one-way ANOVA have been used.

As a result of the research, while the subdimension of achievement and content has come out at middle level, and also the applications of mathematics course has come out at medium level, the teaching period has been at good level but course books and education system has been at low level. It has been indicated that course books are not enough and the opinions of teachers hasn't been considered in research's renewing stage. Some has mentioned that the subjects has become simple regarding the omitted ones from curriculum while some mentioned that the curriculum has been completely cleared. Besides, it has also been mentioned that the curriculum has been able to be taught at fundamental level.

Received: 23–August–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

In our world maintaining to change continuously from past to present countries have tried to keep up with this change. They are still placing great emphasize on education in order to succeed it. Because reaching the way of the target indicated by Mustafa Kemal Atatürk as “Reach and pass the level of contemporary civilization” is provided with the education. The education programs and curriculums prepared for our today's conditions need to be improved most effectively and the programs related to all of the courses need to be structured regarding this purpose so that this education system can be performed and the individuals can succeed to get necessary knowledge and skills (Karagöz,2010; Olkun and Toluk, 2007). Our world has been in a complicated and quantitative status with the last technologic improvements; therefore, mathematical thinking has been more important and the need of teaching

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference

*Key words and phrases.* Mathematics, Secondary School Mathematics Program, 4+4+4 Education System, New Education System.

mathematics has increased (Willoughby, 1990). In this period, mathematics as a course has a great importance and our world's need of people understanding and interpreting the course, likely arises. As for considering either TEOG exam applied in our country or TIMMS and PISA exams applied in international area, we see which level we are at in terms of mathematics. So, it has been a must to change the education system and mathematics programs regarding these results. Because the education programs should follow the improvements all the time depending on the ones in science and technology. Besides, the teaching ones should also change according to time and conditions (Kemertaş, 1999). With these changes, redefining and reviewing of mathematics and it's education in accordance with the identified needs need to be performed (MEB, 2005). The reason of the aforesaid is that mathematics has always been the supporter of the forward but it hasn't been able to exceed the traditional status without it (Savaş and others, 2006). Considering these reasons, mathematics programs regarding 5.,6.,7.,8. grades of secondary school was renewed gradually in 2013-2014 school year in accordance with the decision dated 01.02.2013 prepared by the Ministry of National Education, the Board of Education and Discipline. This research has also been for analyzing teacher's opinions concerning the renewed mathematics program.

Consequently, this research has tried to seek answers for the question of "What are the opinions of secondary school mathematics teacher about the reprepared mathematics program with 4+4+4 education system?" and the following subproblems.

## 2. SUBPROBLEMS

1. How do the Secondary school mathematics teachers think about 4+4+4 education system, it's achievement and teaching period degree, the course of mathematical applications and the course books?

2. Is there a significant difference between the opinions of secondary school mathematics teachers about sex, seniority, place of duty, whether they take seminar or not, and whether the school buildings are the same or not when considered the program's general and subdimensions (System, achievement, teaching period, mathematical applications, and course books) ?

3. How do the mathematics teachers think about the omission of some subjects from the curriculum of secondary school mathematics program reprepared with 4+4+4 education system?

## 3. METHOD

In this study, the survey research design has been used. Survey models are the approaches aiming to describe the situations as either it is in the past or it is still continued (Karasar, 1995). Also, the study deals with that the different groups are compared in terms of some variables thereby performing a relational research. The data collection tool is a kind of likert scale developed by a researcher and the rate of Cronbach alpha reliability coefficient has obtained as 0,886. The scale has 5 factors and contains 28 articles.

## 4. FINDINGS AND RESULTS

According to findings obtained from the first subproblem of the research regarding the question "How do the teachers think about 4+4+4 education system,

achievement dimension, teaching methods and techniques' dimension, the course of mathematical applications and course books?", teachers have positively reacted 4+4+4 education system's being compulsory and gradual with the arithmetic mean 2,62 in middle level. According to teachers' opinions related with the research of Aybek ve Aslan (2015), teachers have positively reacted the 12 years and discontinuous education regarding 4+4+4 discontinuous compulsory education. In 4 interviews of 6, they have positively reacted the 12 years compulsory education in the study of Doğan, Demir ve Pınar (2014). But teachers have negatively reacted to the article "5 grades have been accepted as secondary school" with arithmetic mean 3,32 in middle level in terms of mathematics teaching. Teachers have expressed opinions to the article "The opinions of teachers working in the field have been received while passing to system" with the arithmetic mean 1,85 in weak level. Considering the general of the articles related with the system, they have expressed opinion with the arithmetic mean in weak level.

According to opinions obtained from the research regarding the achievements of the program, suitability has been identified in middle level that the achievement to the mental development is has arithmetic mean 3,24; the achievement to multi-directional thinking has arithmetic mean 2,98; the achievement to their readiness level has arithmetic mean 3,16. The study of Mercan (2011) done for the article about readiness in the past years has the quality to support. Also, the achievement associated with daily life has arithmetic mean 3,20; it's suitability to Turkish National Education and the general features of mathematics education has arithmetic mean 3,32 in middle level. The ordering of subject from concrete to abstract has arithmetic mean 3,42 in good level. The program organized from simple to complicated and it's having cyclical structure has arithmetic mean 3,53 in good level. Teachers have expressed opinions that the article "the content of program is consistent with the general features of mathematics education" has arithmetic mean 3,38; the article "the subject in the program's content is appropriate for the cognitive development of students" has arithmetic mean 3,26; the article "the achievement of the program are clearly understandable and applicable" has arithmetic mean 3,40; the article "the program have had the students like the mathematics course" has arithmetic mean 2,71 in middle level. The results obtained also from the same studies have quality to support the study (Mercan, 2011; Karagöz 2010; İyiol 2011).

In this research, the findings regarding the teaching period have been obtained are the followings: Teachers have expressed opinions in good level to the articles "Students are participating actively in course with the help of my applied activities" with arithmetic mean 3,54; "I am applying learning activities for increasing problems solving skills of students" with arithmetic mean 3,76; "I am benefiting from materials effectively while using learning methods" with arithmetic mean 3,50; "Learning and teaching activities I am using have quality to like mathematics" with arithmetic mean 3,69; "Teaching methods and techniques I am using addresses varied intelligence areas" with arithmetic mean 3,49; "Activities I am getting to be performed are at level which students can apply" with arithmetic mean 3,93; "I am trying to increase the interaction between students in learning and teaching period" with arithmetic mean 3,93.

When looked at the opinions of teachers about the mathematical applications course, the rate of arithmetic mean has come out at 3,17 with middle level regarding the article "Mathematics course has been more funny with the mathematical

applications”. Another article “I am using the mathematical applications most appropriately” has come out at arithmetic mean 3.20 and the other article “The expectations of both mathematical applications and parents are coincided” has come out at arithmetic mean 2,65 in middle level. The information missing of parents about elected courses and the sufficient information can't be transferred are the problems come across by the teachers that have been identified also in the study of Aslan ve Aybek (2015).

When looked at the opinions of the research regarding the course books, our teachers has expressed opinions to the article “no assistant source are needed as course books are enough for learning-teaching period” in weak level with the arithmetic mean 1,60. They have expressed opinions with arithmetic mean 1,97 to the article “The teaching style of course in course books is sufficiently clear”, 1,90 to the article “The exercises developing operation capabilities are sufficiently included in books” in weak level. Teachers has also mentioned in the research of Mutu (2008) done about the same subject in the past years that the content of 6. and 7. grade books is totally weak and the subject ordering is inconsistent; so they need assistant books because of this, and the sample questions are missing.

When looked at the answers to open ended questions we have asked regarding the omissions of some subjects, some teacher look positively at program's becoming simple but others has a number of concerns about that. They have considered that the subjects fully omitted will cause problems in high school and next education life, and they have defended that these subjects should be mentioned even a little. Furthermore, the positive contribution of the subjects for the students to understand and focus on other subjects is their another opinion and they have also considered that it has a positive impact in terms of time because of the mathematics course's heavy subjects.

#### REFERENCES

- [1] Aslan, S. ve Aybek, B. (2015). “Ortaokul Öğretmenlerinin 4+4+4 Kesintili Zorunlu Eğitim Sistemine Yönelik Yaşadıkları Sorunlar (Elazığ İli Örneği)”. *ElementaryEducation Online*, 14(2), 770-786, 2015. İlköğretim Online, 14(2), 770-786, 2015. [Online]: <http://ilkogretim-online.org.tr> <http://dx.doi.org/10.17051/io.2015.634> 54
- [2] Budak, M. ve Okur, M. (2012). *International Journal of New Trends in Arts, Sports & Science Education* - 2012, volume 1, issue 4
- [3] Cresswell, J.W. and Clark, V.L. (2015). *Karma Yöntem Araştırmaları (Çeviri Editörleri: Y. Dede, S.B. Demir)*. Ankara: Anı Yayıncılık (Eserin orijinali 2011'de yayımlandı).
- [4] Demir, S.B. ; Doğan, S. ve Pınar, M. A. (2013). “4+4+4 Eğitim Sisteminin Yansımaları: Beşinci Sınıflardaki Eğitim-Öğretim Sürecinin Branş Öğretmenlerinin Görüşleri Doğrultusunda Değerlendirilmesi”. *Turkish Studies - International Periodical For The Languages, Literature and History of Turkish or Turkic* Volume 8/9 Summer 2013, p. 1081-1098, ANKARA-TURKEY
- [5] Demir, S.B; Doğan, S. ve Pınar, M.A (2014). “4+4+4 Kesintili Zorunlu Eğitim Sisteminin Sınıf Öğretmenlerinin Görüşleri Doğrultusunda Değerlendirilmesi” *Elementary Education Online*, 13(2), 503-517, 2014. İlköğretim Online, 13(2), 503-517, 2014. [Online]: <http://ilkogretim-online.org.tr>
- [6] Greene, J.C., Caracelli, V.J., Graham, W.F. (1989). *Toward a conceptual framework for mixed-method evaluation designs*. *Educational Evaluation and Policy Analysis*, 11(3), 255-274.
- [7] İyiol, F.F (2011). İlköğretim 8.Sınıf Matematik Programının Öğretmenlerin Görüşlerine Göre Değerlendirilmesi. Selçuk Üniversitesi Eğitim Bilimleri Enstitüsü ilköğretim Anabilim Dalı Yüksek Lisans Tezi.

- [8] Karagöz, E. (2010). İlköğretim II. Kademe Matematik Dersi Öğretim Programının Öğretmen Görüşleri Doğrultusunda Değerlendirilmesi. Muğla Üniversitesi Sosyal Bilimler Enstitüsü Eğitim Programları ve Öğretimi Bilim Dalı Yüksek Lisans Tezi.
- [9] Karasar, N. (1995). Araştırmalarda Rapor Hazırlama 8. Basım Alkim yayınları Ankara.
- [10] Kemertaş, İ. (1999). Uygulamalı genel öğretim yöntemleri: "Öğretimde planlama ve değerlendirme İstanbul: Birsen Yayınevi
- [11] Kurtulmuş, Y. (2010). İlköğretim 8.Sınıf Matematik Ders Kitapları İle İlgili Öğretmen Görüşleri. Mustafa Kemal Üniversitesi Sosyal Bilimler Enstitüsü İlköğretim AnaBilim Dalı Yüksek Lisans Tezi.
- [12] MEB. (2005). İlköğretim Hayat Bilgisi, Matematik, Sosyal Bilgiler, Türkçe, Fen ve Teknoloji Dersi Öğretim Programlarında Değişiklik Yapılması. Milli Eğitim Bakanlığı Tebliğler Dergisi. 2575:
- [13] Mercan, Z. (2011). İlköğretim Matematik Dersi Öğretim Programının Eğitim Durumu Boyutunun Öğretmen ve Öğrenci Görüşleri Açısından Değerlendirilmesi. Muğla Üniversitesi Eğitim Bilimleri Enstitüsü Eğitim Programları ve Öğretimi Yüksek Lisans Tezi
- [14] Mutu, B.B (2008). 6. ve 7. Sınıf Matematik Ders Kitapları Hakkında Öğretmen Görüşleri. Osmangazi Üniversitesi Fen Bilimleri Enstitüsü İlköğretim Ana bilimdalı Yüksek Lisans Tezi
- [15] Olkun S.,Toluk, Z. (2007). İlköğretimde Etkinlik Temelli Matematik Öğretimi. (3.basım) Maya Akademi.
- [16] Savaş, E.;Obay, M. ve Duru A. (2006), Öğrenme Etkinliklerinin Öğrencilerin Matematik Başarıları Üzerindeki Etkisi, Journal Of OafqazUniversty, Yıl 2006, sayı:17
- [17] Tashakkari, A. ve Creswell, d.w. (2007). The eros of mixedmethods [Editorial]. Journal of mixedmethodsresearch, 1(1), 3-7.
- [18] Willoughby, S. (1990), MathematicsEducationFor A Changing World ASCD Virginia. www.matder.org.tr- www.ttkb.org.com.tr

**THE DIRECT AND INVERSE SPECTRAL PROBLEM FOR  
STURM-LIOUVILLE OPERATOR WITH DISCONTINUOUS  
COEFFICIENT**

KHANLAR R. MAMEDOV AND DÖNE KARAHAN

ABSTRACT. In this study, for Sturm-Liouville operator with discontinuous coefficient encountered in the non-homogeneous materials, direct and inverse problems are investigated. The spectral properties of the Sturm-Liouville problem with discontinuous coefficient such as the orthogonality of its eigenfunctions and simplicity of its eigenvalues are examined. Asymptotic formula is found for eigenvalues, and resolvent operator is constructed. The expansion formula with respect to eigenfunctions is obtained. It is shown that its eigenvalues are in the form of a complete system. Also, the Weyl solution and Weyl function are defined. Uniqueness theorems for the solution of the inverse problem according to spectral data are proved.

Received: 17–August–2016

Accepted: 29–August–2016

1. INTRODUCTION

In this study, the heat problem of a rod that consists of two parts with fixed cross section is examined. The side surfaces of the rod have been isolated and have different physical features [1]-[4]. When initial temperature is given arbitrary and the temperature at the ending points is not equal to zero, the heat problem of the rod takes the following form:

$$\begin{aligned}\rho(x)U_t &= U_{xx} + q(x)U, & 0 \leq x \leq \pi, \\ U(x, 0) &= \phi(x), \quad U_t(x, 0) = \psi(x), & 0 \leq x \leq \pi \\ U_x(0, t) &= 0, \quad U_x(\pi, t) = 0, & t > 0\end{aligned}$$

where the function  $U(x, t)$  is the temperature in the bar at the time  $t$ ,  $\rho(x)$  is a piecewise constant function and refers to the density of the rod and  $\phi(x)$ ,  $\psi(x)$  are enough smooth functions. By the method of separation of variables, the preceding equation is reduced to a boundary value problem for Sturm-Liouville equation:

$$(1.1) \quad -y'' + q(x)y = \lambda^2 \rho(x)y, \quad 0 \leq x \leq \pi,$$

$$(1.2) \quad y'(0) = y'(\pi) = 0,$$

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 34A55, 34B24, 47E05.

*Key words and phrases.* Sturm-Liouville operator, expansion formula, inverse problem, Weyl function.

where in particular,  $\rho(x)$  is chosen as

$$(1.3) \quad \rho(x) = \begin{cases} 1 & 0 \leq x < a \\ \alpha^2 & a \leq x \leq \pi. \end{cases}$$

$q(x) \in L_2(0, \pi)$  is a real valued function and  $\lambda$  is a complex parameter. Then, in finding the solution of the above diffusion problem, spectral problem (1.1), (1.2) must be examined [1]-[5]. The spectral problems with discontinuous coefficient on the bounded interval are investigated in [6]-[15]. The similar problems on the half line by different authors have been studied (see [16]-[18]). Let  $\varphi(x)$  and  $\psi(x)$  be solutions of (1.1), (1.2) boundary value problem satisfying the initial conditions

$$(1.4) \quad \varphi(0, \lambda) = 1, \varphi'(0, \lambda) = 0$$

and

$$(1.5) \quad \psi(\pi, \lambda) = 1, \psi'(\pi, \lambda) = 0.$$

Denote

$$(1.6) \quad \Delta(\lambda) = W[\varphi(x, \lambda), \psi(x, \lambda)] = \varphi'(x, \lambda)\psi(x, \lambda) - \varphi(x, \lambda)\psi'(x, \lambda).$$

The function  $\Delta(\lambda)$  is called the characteristic function of the problem (1.1), (1.2), and substituting  $x = 0$  and  $x = \pi$  into (1.6), we get

$$(1.7) \quad \Delta(\lambda) = \varphi'(\pi, \lambda) = -\psi'(0, \lambda).$$

**Lemma 1.1.** *The eigenfunctions  $y_1(x, \lambda_1)$  and  $y_2(x, \lambda_2)$  corresponding to different eigenvalues  $\lambda_1 \neq \lambda_2$  are orthogonal.*

*Proof.* Since  $y_1(x, \lambda_1)$  and  $y_2(x, \lambda_2)$  are eigenfunctions of problem (1.1), (1.2), we get

$$\begin{aligned} -y_1''(x, \lambda_1) + q(x)y_1(x, \lambda_1) &= \lambda_1^2 \rho(x)y_1(x, \lambda_1), \\ -y_2''(x, \lambda_2) + q(x)y_2(x, \lambda_2) &= \lambda_2^2 \rho(x)y_2(x, \lambda_2). \end{aligned}$$

Multiplying these equalities by  $y_1(x, \lambda_1)$  and  $-y_2(x, \lambda_2)$ , respectively, and adding together,

$$\frac{d}{dx} \{ \langle y_2(x, \lambda_2), y_1(x, \lambda_1) \rangle \} = (\lambda_1^2 - \lambda_2^2) \rho(x)y_1(x, \lambda_1)y_2(x, \lambda_2)$$

is found. Integrating from 0 to  $\pi$  and using the condition (1.2), we have

$$(\lambda_1^2 - \lambda_2^2) \int_0^\pi \rho(x)y_1(x, \lambda_1)y_2(x, \lambda_2)dx = 0.$$

Since  $\lambda_1 \neq \lambda_2$ ,

$$\int_0^\pi \rho(x)y_1(x, \lambda_1)y_2(x, \lambda_2)dx = 0.$$

□

**Corollary 1.1.** *The eigenvalues of the boundary value problem (1.1), (1.2) are real.*

**Lemma 1.2.** *The zeros  $\lambda_n$  of characteristic function  $\Delta(\lambda)$  coincide with the eigenvalues of the boundary value problem (1.1), (1.2). The functions  $\varphi(x, \lambda_n)$  and  $\psi(x, \lambda_n)$  are eigenfunctions and there exists a sequence  $\beta_n$  such that*

$$(1.8) \quad \psi(x, \lambda_n) = \beta_n \varphi(x, \lambda_n), \quad \beta_n \neq 0.$$

*Proof.* 1) Let  $\lambda_0$  be zero of  $\Delta(\lambda)$ . Then, because of (1.6),  $\psi(x, \lambda_0) = \beta_0 \varphi(x, \lambda_0)$  and the function  $\psi(x, \lambda_0)$  and  $\varphi(x, \lambda_0)$  satisfy the boundary condition (1.2). Thus,  $\lambda_0$  is an eigenvalue and  $\psi(x, \lambda_0)$ ,  $\varphi(x, \lambda_0)$  are corresponding eigenfunctions.

2) Let  $\lambda_0$  be an eigenvalue of the problem (1.1), (1.2) and let  $y_0(x)$  be a corresponding eigenfunction. Then,  $y_0(x)$  satisfies the boundary condition (1.2). Clearly,  $y_0(x) \neq 0$ . Without loss of generality, we put  $y_0(0) = 1$ . Then  $y_0'(0) = 0$ , and consequently,  $y_0(x) \equiv \varphi(x, \lambda)$ . Hence, from (1.7),  $\Delta_0(\lambda) = 0$ . We have proved that for each eigenvalue there exists only one eigenfunction.  $\square$

**Lemma 1.3.** *The eigenvalues of the boundary value problem (1.1), (1.2) are simple and*

$$(1.9) \quad \dot{\Delta}(\lambda_n) = 2\lambda_n \alpha_n \beta_n,$$

where

$$\alpha_n := \int_0^\pi \rho(x) \varphi^2(x, \lambda_n) dx.$$

is the normalizing number of (1.1), (1.2).

*Proof.* Since  $\varphi(x, \lambda_n)$  and  $\psi(x, \lambda)$  are the solutions of this problem,

$$\begin{aligned} -\varphi''(x, \lambda_n) + q(x)\varphi(x, \lambda_n) &= \lambda_n^2 \rho(x)\varphi(x, \lambda_n), \\ -\psi''(x, \lambda) + q(x)\psi(x, \lambda) &= \lambda^2 \rho(x)\psi(x, \lambda), \end{aligned}$$

are valid. Multiplying these equations by  $\psi(x, \lambda)$  and  $-\varphi(x, \lambda_n)$ , respectively, and adding them together, we get

$$\frac{d}{dx} \{ \langle \psi(x, \lambda), \varphi(x, \lambda_n) \rangle \} = (\lambda_n^2 - \lambda^2) \rho(x) \varphi(x, \lambda_n) \psi(x, \lambda).$$

Integrating from 0 to  $\pi$  and using the condition (1.2),

$$\int_0^\pi \rho(x) \varphi(x, \lambda_n) \psi(x, \lambda) = \frac{\Delta(\lambda_n) - \Delta(\lambda)}{\lambda_n^2 - \lambda^2}$$

is found. From Lemma 2, since  $\psi(x, \lambda_n) = \beta_n \varphi(x, \lambda_n)$ , as  $\lambda \rightarrow \lambda_n$ , we obtain

$$\dot{\Delta}(\lambda_n) = 2\lambda_n \alpha_n \beta_n$$

where  $\beta_n = \psi(0, \lambda_n)$ . Thus, it follows that  $\dot{\Delta}(\lambda_n) \neq 0$ .  $\square$

## 2. ON THE EIGENVALUES OF PROBLEM (1.1), (1.2) AT $q(x) \equiv 0$

Denote by  $\varphi_0(x, \lambda)$  the solution equation  $-y'' = \lambda^2 \rho(x)y$ , satisfying the condition (1.4). It has the following form:

$$(2.1) \quad \varphi_0(x, \lambda) = \frac{1}{2} \left( 1 + \frac{1}{\sqrt{\rho(x)}} \right) \cos \lambda \mu^+(x) + \frac{1}{2} \left( 1 - \frac{1}{\sqrt{\rho(x)}} \right) \cos \lambda \mu^-(x),$$

where  $\mu^\pm(x) = \pm x \sqrt{\rho(x)} + a(1 \mp \sqrt{\rho(x)})$ .

It is easy show that if  $(\lambda_n^0)^2$  are eigenvalues of problem (1.1), (1.2) at  $q(x) \equiv 0$ , then  $\lambda_n^0$  can be found from the equation  $\varphi_0'(\pi, \lambda) = 0$ , that is, from the equation

$$(2.2) \quad \begin{aligned} \Delta_0(\lambda) &= -\frac{1}{2} \lambda (\alpha + 1) \sin \lambda \mu^+(\pi) + \frac{1}{2} (\alpha - 1) \lambda \sin \lambda \mu^-(\pi) = 0 \\ \sin \lambda \mu^+(\pi) - \frac{\alpha - 1}{\alpha + 1} \sin \lambda \mu^-(\pi) &= 0. \end{aligned}$$



At last, from (2.2), it follows that

$$(2.3) \quad \lambda_n^0 = \frac{1}{\mu^+(\pi)} (n\pi + \epsilon_n)$$

where

$$\epsilon_n = (-1)^n \frac{\alpha - 1}{\alpha + 1} \sin \left( \frac{\mu^+(\pi)}{\mu^-(\pi)} n\pi \right) + O \left( \frac{1}{n} \right).$$

**Lemma 2.1.** *Roots of the function  $\Delta_0(\lambda)$  are isolated, that is,*

$$\inf_{n \neq k} |\lambda_n^0 - \lambda_k^0| = \beta > 0.$$

### 3. ASYMPTOTIC FORMULAS OF EIGENVALUES

Using representation for solution  $e(x, \lambda)$  of equation (1.1) satisfying the initial conditions  $e(0, \lambda) = 1$ ,  $e'(0, \lambda) = i\lambda$  (see[8]), it is easy to obtain the following integral representation for the solution  $\varphi(x, \lambda)$ :

$$\varphi(x, \lambda) = \varphi_0(x, \lambda) + \int_0^{\mu^+(x)} A(x, t) \cos \lambda t dt$$

where  $K(x, \cdot) \in L_1(-\mu^+(x), \mu^+(x))$  and  $A(x, t) = K(x, t) - K(x, -t)$ . The kernel  $A(x, t)$  processes the following properties:

- i)  $A(\pi, \mu^+(\pi)) = \frac{1}{4} \int_0^\pi \frac{1}{\sqrt{\rho(t)}} \left( 1 + \frac{1}{\sqrt{\rho(t)}} \right) q(t) dt$ ,
- ii)  $A(\pi, \mu^-(\pi) + 0) - A(\pi, \mu^-(\pi) - 0) = \frac{1}{4} \int_0^\pi \frac{1}{\sqrt{\rho(t)}} \left( 1 - \frac{1}{\sqrt{\rho(t)}} \right) q(t) dt$ .

**Theorem 3.1.** *Boundary value problem (1.1), (1.2) has a countable set of simple eigenvalues  $\{\lambda_n^2\}_{n \geq 1}$ , where*

$$(3.1) \quad \lambda_n = \lambda_n^0 + \frac{d_n}{\lambda_n^0} + \frac{k_n}{n}, \quad (\lambda_n > 0), \quad k_n \in l_2$$

where  $\lambda_n^0$  are zeros of the function

$$(3.2) \quad \Delta_0(\lambda) = -\frac{1}{2} \lambda (\alpha + 1) \sin \lambda \mu^+(\pi) + \frac{1}{2} (\alpha - 1) \lambda \sin \lambda \mu^-(\pi).$$

$\{\lambda_n^0\}^2$  are the eigenvalues of problem (1.1), (1.2), when  $q(x) \equiv 0$ ,  $d_n$  is a bounded sequence

$$(3.3) \quad d_n = \frac{h^+ \cos \lambda_n^0 \mu^+(\pi) + h^- \cos \lambda_n^0 \mu^-(\pi)}{\frac{1}{2} (\alpha + 1) \mu^+(\pi) \cos \lambda_n^0 \mu^+(\pi) - \frac{1}{2} (\alpha - 1) \mu^-(\pi) \cos \lambda_n^0 \mu^-(\pi)}.$$

*Proof.* Let  $\varphi(x, \lambda)$  be the solution of equation (1.1) at initial conditions  $\varphi(0, \lambda) = 1$ ,  $\varphi'(0, \lambda) = 0$ . Then the characteristic function  $\Delta(\lambda) = \varphi'(\pi, \lambda)$  is entire with respect to  $\lambda$  and it has the most countable set of zeros  $\lambda_n$  and numbers  $\lambda_n^2$  are eigenvalues of boundary value problem (1.1), (1.2). The standard method of variations of an arbitrary constants leads to the following integral equation for the solution  $\varphi(x, \lambda)$

$$(3.4) \quad \varphi(x, \lambda) = \varphi_0(x, \lambda) + \int_0^x g(x, t; \lambda) q(t) \varphi(t, \lambda) dt$$

where

$$(3.5) \quad g(x, t; \lambda) = \frac{1}{2} \left( \frac{1}{\sqrt{\rho(x)}} + \frac{1}{\sqrt{\rho(t)}} \right) \frac{\sin \lambda (\mu^+(x) - \mu^+(t))}{\lambda} + \frac{1}{2} \left( \frac{1}{\sqrt{\rho(x)}} - \frac{1}{\sqrt{\rho(t)}} \right) \frac{\sin \lambda (\mu^-(x) - \mu^+(t))}{\lambda}$$

and  $\varphi_0(x, \lambda)$  is the solution of equation (1.1) at  $q(x) = 0$ , satisfying the conditions  $\varphi(0, \lambda) = 1$ ,  $\varphi'(0, \lambda) = 0$ . From (3.4), after differentiating, we find

$$(3.6) \quad \varphi'(x, \lambda) = \varphi'_0(x, \lambda) + \int_0^x g_x(x, t; \lambda) q(t) \varphi(t, \lambda) dt$$

where

$$(3.7) \quad g_x(x, t; \lambda) = \sqrt{\rho(x)} \frac{1}{2} \left( \frac{1}{\sqrt{\rho(x)}} + \frac{1}{\sqrt{\rho(t)}} \right) \frac{\cos \lambda (\mu^+(x) - \mu^+(t))}{\lambda} + \sqrt{\rho(x)} \frac{1}{2} \left( \frac{1}{\sqrt{\rho(x)}} - \frac{1}{\sqrt{\rho(t)}} \right) \frac{\cos \lambda (\mu^-(x) - \mu^+(t))}{\lambda}$$

Consequently, if we put here  $x = \pi$ , we have

$$(3.8) \quad \Delta(\lambda) = \Delta_0(\lambda) + \int_0^\pi g'_\pi(x, t; \lambda) q(t) \varphi(t, \lambda) dt.$$

Now, from

$$(3.9) \quad \varphi(x, \lambda) = \varphi_0(x, \lambda) + O\left(\frac{e^{|Im \lambda| \mu^+(x)}}{|\lambda|}\right), \quad |\lambda| \rightarrow +\infty$$

we obtain

$$(3.10) \quad \Delta(\lambda) = \Delta_0(\lambda) + h^+ \cos \lambda_n^0 \mu^+(\pi) + h^- \cos \lambda_n^0 \mu^-(\pi) + K_0(\lambda),$$

where

$$(3.11) \quad h^\pm = \frac{1}{4} (1 \pm \alpha) \int_0^a q(t) dt + \frac{1}{4} \left(1 \pm \frac{1}{\alpha}\right) \int_a^\pi q(t) dt,$$

and

$$(3.12) \quad K_0(\lambda) = \frac{1}{4} \int_0^a [(1 + \alpha) \cos \lambda (2\mu^+(t) - \mu^+(\pi)) + (1 - \alpha) \cos \lambda (2\mu^+(t) - \mu^-(\pi))] q(t) dt + \frac{1}{4} \int_a^\pi \left[ \left(1 + \frac{1}{\alpha}\right) \cos \lambda (2\mu^+(t) - \mu^+(\pi)) \right] q(t) dt + \frac{1}{4} \int_a^\pi \left[ \left(1 - \frac{1}{\alpha}\right) \cos \lambda (\mu^+(\pi) + \mu^-(t) - \mu^+(t)) \right] q(t) dt + 0 \left( \frac{e^{|Im \lambda| \mu^+(\pi)}}{|\lambda|} \right).$$

Let us denote  $G_\delta = \{\lambda : |\lambda - \lambda_n^0| \geq \delta\}$ , where  $\delta$  is a sufficiently small positive number  $\delta < \frac{\beta}{2}$  (see lemma 4). It is easy to show that (see [3])

$$(3.13) \quad |\Delta_0(\lambda)| \geq |\lambda| C_\delta e^{|Im \lambda| \mu^+(\pi)}, \quad \lambda \in G_\delta, \quad C_\delta > 0.$$

On the other hand, we obtain

$$(3.14) \quad \Delta(\lambda) - \Delta_0(\lambda) \leq O\left(e^{|Im\lambda|\mu^+(\pi)}\right), \quad |\lambda| \rightarrow \infty.$$

Consider the contour  $\Gamma_n = \{\lambda : |\lambda| = |\lambda_n^0| + \frac{\beta}{2}\}$ , ( $n = 1, 2, \dots$ ). We have from (3.10)

$$(3.15) \quad |\Delta(\lambda) - \Delta_0(\lambda)| \leq \tilde{C}e^{|Im\lambda|\mu^+(\pi)}, \quad \lambda \in \Gamma_n,$$

for sufficiently large  $n$ , where  $\tilde{C} > 0$ . Applying now Rouché's theorem, we have that the number of zeros of  $\Delta_0(\lambda)$  inside  $\Gamma_n$  coincides with the number of zeros of  $\Delta(\lambda) = \{\Delta(\lambda) - \Delta_0(\lambda)\} + \Delta_0(\lambda)$ . Further applying the Rouché's theorem to the circle  $\gamma_n(\delta) = \{\lambda : |\lambda - \lambda_n^0| \leq \delta\}$ , we conclude that for sufficiently large  $n$ , there exist only one zero  $\lambda_n$  of the function  $\Delta(\lambda)$  in  $\gamma_n(\delta)$ . By virtue of the arbitrariness of  $\delta > 0$  we have

$$(3.16) \quad \lambda_n = \lambda_n^0 + \varepsilon_n, \quad \varepsilon_n = o(1), \quad n \rightarrow \infty.$$

Substituting (3.16) into (3.10), we obtain and taking into our account the relations

$$\Delta_0(\lambda_n^0) = -\frac{1}{2}\lambda_n^0(\alpha + 1)\sin\lambda_n^0\mu^+(\pi) + \frac{1}{2}\lambda_n^0(\alpha - 1)\sin\lambda_n^0\mu^-(\pi) = 0,$$

$$\sin\varepsilon_n\mu^+(\pi) \sim \varepsilon_n\mu^+(\pi), \quad \cos\varepsilon_n\mu^+(\pi) \sim 1, \quad n \rightarrow \infty$$

we get

$$(3.17) \quad \varepsilon_n = \frac{d_n}{\lambda_n^0 + \varepsilon_n} + \frac{\varepsilon_n}{\lambda_n^0 + \varepsilon_n}\tilde{d}_n + \frac{\tilde{K}_n}{\lambda_n^0 + \varepsilon_n}$$

where

$$d_n = \frac{h^+ \cos\lambda_n^0\mu^+(\pi) + h^- \cos\lambda_n^0\mu^-(\pi)}{\frac{1}{2}(\alpha + 1)\mu^+(\pi)\cos\lambda_n^0\mu^+(\pi) - \frac{1}{2}(\alpha - 1)\mu^-(\pi)\cos\lambda_n^0\mu^-(\pi)},$$

$\tilde{K}_n = K_0(\lambda_n^0 + \varepsilon_n)$  and

$$\tilde{d}_n = \frac{h^+\mu^+(\pi)\sin\lambda_n^0\mu^+(\pi) + h^-\mu^-(\pi)\sin\lambda_n^0\mu^-(\pi)}{\frac{1}{2}(\alpha + 1)\mu^+(\pi)\cos\lambda_n^0\mu^+(\pi) - \frac{1}{2}(\alpha - 1)\mu^-(\pi)\cos\lambda_n^0\mu^-(\pi)}.$$

Since  $\frac{1}{\lambda_n^0 + \varepsilon_n} = O(\frac{1}{n})$ ,  $\frac{\varepsilon_n}{\lambda_n^0 + \varepsilon_n} = o(\frac{1}{n})$ ,  $n \rightarrow \infty$  we have that  $d_n, \tilde{d}_n$  are bounded and (3.17) implies

$$\varepsilon_n = O\left(\frac{1}{n}\right), \quad n \rightarrow \infty.$$

Using (3.17) once more, we can obtain more precisely as  $n \rightarrow \infty$

$$(3.18) \quad \varepsilon_n = \frac{d_n}{\lambda_n^0} + \frac{k_n}{n}, \quad k_n \in l_2$$

where  $k_n = \frac{\mu^+(\pi)}{\pi}\tilde{K}_n + O(\frac{1}{n})$ ,  $n \rightarrow \infty$ . The theorem is proved.  $\square$

## 4. SPECTRAL EXPANSION FORMULA

**Theorem 4.1.** 1) The system of eigenfunctions  $\{\varphi(x, \lambda_n)\}_{n \geq 1}$  of boundary value problem (1.1), (1.2) is complete in  $L_2(0, \pi; \rho)$ ;

2) If  $f(x)$  is an absolutely continuous function on the segment  $[0, \pi]$ , and  $f'(0) = f'(\pi) = 0$ , then

$$(4.1) \quad f(x) = \sum_{n=1}^{\infty} a_n \varphi(x, \lambda_n),$$

where

$$(4.2) \quad a_n = \frac{1}{\alpha_n} \int_0^{\pi} f(t) \varphi(t, \lambda_n) \rho(t) dt,$$

and the series (4.1) converges uniformly on  $[0, \pi]$ ;

3) For  $f(x) \in L_2(0, \pi; \rho)$  the series (4.1) converges in  $L_2(0, \pi; \rho)$ , moreover the Parseval equality

$$(4.3) \quad \int_0^{\pi} |f(x)|^2 \rho(x) dx = \sum_{n=1}^{\infty} \alpha_n |a_n|^2$$

holds.

*Proof.* Let  $\psi(x, \lambda)$  be a solution of equation (1.1) under the initial conditions  $\psi(\pi, \lambda) = 1$ ,  $\psi'(\pi, \lambda) = 0$ . Denote

$$(4.4) \quad G(x, t; \lambda) = -\frac{1}{\Delta(\lambda)} \begin{cases} \psi(x, \lambda) \varphi(t, \lambda), & x \geq t \\ \varphi(x, \lambda) \psi(t, \lambda), & t \geq x \end{cases}$$

and let us consider the function

$$(4.5) \quad Y(x, \lambda) = \int_0^{\pi} \rho(t) f(t) G(x, t; \lambda) dt$$

which is a solution of the boundary value problem

$$(4.6) \quad -Y''(x, \lambda) + q(x)Y(x, \lambda) = \lambda^2 \rho(x)Y(x, \lambda) - f(x)\rho(x), \\ Y'(0, \lambda) = 0, Y'(\pi, \lambda) = 0.$$

Using (1.9), we obtain

$$(4.7) \quad \operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = \frac{1}{2\alpha_n \lambda_n} \varphi(x, \lambda_n) \int_0^{\pi} \rho(t) f(t) \varphi(t, \lambda_n) dt.$$

Let  $f(x) \in L_2(0, \pi; \rho)$  be such that

$$\int_0^{\pi} \rho(t) f(t) \varphi(t, \lambda_n) dt = 0 \quad n = 1, 2, 3, \dots$$

Then, from (4.7), we have  $\operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = 0$ ; consequently, for fixed  $x \in [0, \pi]$ , the function  $Y(x, \lambda)$  is entire with respect to  $\lambda$ . On the other hand, since

$$(4.8) \quad \Delta(\lambda) \geq |\lambda| \tilde{C}_\delta e^{|\operatorname{Im} \lambda| \mu^+(\pi)}, \quad \lambda \in G_\delta, \quad \tilde{C}_\delta > 0.$$

$$(4.9) \quad \varphi(x, \lambda) = O\left(e^{|\operatorname{Im} \lambda| \mu^+(x)}\right), \quad \psi(x, \lambda) = O\left(e^{|\operatorname{Im} \lambda| (\mu^+(\pi) - \mu^+(x))}\right), \quad |\lambda| \rightarrow \infty,$$

from (4.5), it follows that for fixed  $\delta > 0$  and sufficiently large  $\lambda^* > 0$ :

$$|Y(x, \lambda)| \leq \frac{C_\delta}{|\lambda|}, \quad \lambda \in G_\delta, \quad |\lambda| \geq \lambda^*.$$

Using the maximum principle for module of analytic functions and Liouville theorem, we conclude that  $Y(x, \lambda) \equiv 0$ . This fact and (4.6) imply that  $f(x) = 0$  a.e. on  $[0, \pi]$ . Thus, statement (1) of theorem is proved.

Let  $f(x) \in AC[0, \pi]$  be an arbitrary absolutely continuous function. Let us transform the function  $Y(x, \lambda)$  to the form

$$Y(x, \lambda) = -\frac{1}{\lambda^2 \Delta(\lambda)} \left\{ \psi(x, \lambda) \int_0^x (-\varphi''(t, \lambda) + q(t)\varphi(t, \lambda)) f(t) dt - \varphi(x, \lambda) \int_x^\pi (-\psi''(t, \lambda) + q(t)\psi(t, \lambda)) f(t) dt \right\}.$$

Integrating by parts the addends with the second-order derivatives and taking into account conditions  $f'(0) = 0$ ,  $f'(\pi) = 0$ , we have

$$(4.10) \quad Y(x, \lambda) = \frac{f(x)}{\lambda^2} - \frac{1}{\lambda^2} (Z_1(x, \lambda) + Z_2(x, \lambda)),$$

where

$$Z_1(x, \lambda) = \frac{1}{\Delta(\lambda)} \left[ \psi(x, \lambda) \int_0^x g(t) \varphi'(t, \lambda) dt + \varphi(x, \lambda) \int_x^\pi g(t) \psi'(t, \lambda) dt \right],$$

$$Z_2(x, \lambda) = \frac{1}{\Delta(\lambda)} \left[ \psi(x, \lambda) \int_0^x \varphi(t, \lambda) f(t) \rho(t) dt + \varphi(x, \lambda) \int_x^\pi \psi(t, \lambda) f(t) \rho(t) dt \right].$$

Here  $g(t) = f'(t)$ . Now consider the contour integral

$$(4.11) \quad I_N(x) = 2 \sum_{n=1}^N \operatorname{Res}_{\lambda=\lambda_n} Y(x, \lambda) = \sum_{n=1}^N a_n \varphi(x, \lambda_n)$$

where

$$a_n = \frac{1}{\alpha_n} \int_0^\pi \rho(t) f(t) \varphi(t, \lambda_n) dt.$$

On the other hand taking into account (4.10), we have

$$(4.12) \quad I_N(x) = f(x) - \frac{1}{2\pi i} \int_{\Gamma_N} \frac{1}{\lambda} (Z_1(x, \lambda) + Z_2(x, \lambda)) d\lambda.$$

Comparing (4.11) and (4.12), we obtain

$$f(x) = \sum_{n=1}^{\infty} a_n \varphi(x, \lambda_n) + \xi_N(x),$$

where

$$\xi_N(x) = \frac{1}{2\pi i} \int_{\Gamma_N} \frac{1}{\lambda} (Z_1(x, \lambda) + Z_2(x, \lambda)) d\lambda.$$

Therefore, in order to prove the item (2) of the theorem, it suffices to show that

$$(4.13) \quad \lim_{N \rightarrow \infty} \max_{0 \leq x \leq \pi} |\xi_N(x)| = 0.$$

From the estimates of solution  $\varphi(x, \lambda)$ ,  $\psi(x, \lambda)$  and the function  $\Delta(\lambda)$ , it follows that for fixed  $\delta > 0$  and sufficiently large  $\lambda^* > 0$ ,

$$(4.14) \quad \max_{0 \leq x \leq \pi} |Z_2(x, \lambda)| \leq \frac{C_2}{|\lambda|}, \quad \lambda \in G_\delta, \quad |\lambda| \geq \lambda^*, C_2 > 0.$$

Let us show that

$$(4.15) \quad \lim_{\substack{|\lambda| \rightarrow \infty \\ \lambda \in G_\delta}} \max_{0 \leq x \leq \pi} |Z_1(x, \lambda)| = 0.$$

At first, it was supposed that  $g(t)$  is absolutely continuous on  $[0, \pi]$ . In this case, integration by parts gives

$$Z_1(x, \lambda) = -\frac{1}{\Delta(\lambda)} \left\{ \psi(x, \lambda) \int_0^x \varphi(t, \lambda) g'(t) dt + \varphi(x, \lambda) \int_x^\pi \psi(t, \lambda) g'(t) dt \right\},$$

therefore, similarly to  $Z_2(x, \lambda)$ , we have

$$\max_{0 \leq x \leq \pi} |Z_1(x, \lambda)| \leq \frac{C_1}{|\lambda|}, \quad \lambda \in G_\delta, \quad |\lambda| \geq \lambda^*, C_1 > 0.$$

In the general case, we fix  $\varepsilon > 0$  and choose absolutely continuous function  $g_\varepsilon(t)$  such that

$$\int_0^\pi |g_\varepsilon(t) - g(t)| dt < \varepsilon.$$

Then, using the estimates  $\varphi(x, \lambda)$ ,  $\psi(x, \lambda)$ ,  $\Delta(\lambda)$ , one can find  $\lambda^{**} > 0$  such that when  $\lambda \in G_\delta$ ,  $|\lambda| \geq \lambda^{**}$ , from the relation

$$\begin{aligned} Z_1(x, \lambda) &= \frac{1}{\Delta(\lambda)} \left[ \psi(x, \lambda) \int_0^x \varphi'(t, \lambda) (g_\varepsilon(t) - g(t)) dt + \right. \\ &\quad \left. + \varphi(x, \lambda) \int_x^\pi \psi'(t, \lambda) (g_\varepsilon(t) - g(t)) dt \right] + \\ &+ \frac{1}{\Delta(\lambda)} \left[ \psi(x, \lambda) \int_0^x \varphi(t, \lambda) g'_\varepsilon(t) dt - \varphi(x, \lambda) \int_x^\pi \psi(t, \lambda) g'_\varepsilon(t) dt \right], \end{aligned}$$

we have

$$\max_{0 \leq x \leq \pi} |Z_1(x, \lambda)| \leq C \int_0^\pi |g_\varepsilon(t) - g(t)| dt + \frac{\tilde{C}(\varepsilon)}{|\lambda|} < C_\varepsilon + \frac{\tilde{C}(\varepsilon)}{|\lambda|}, \quad \lambda \in G_\delta, \quad |\lambda| \geq \lambda^{**}.$$

Consequently,

$$\overline{\lim}_{\substack{|\lambda| \rightarrow \infty \\ \lambda \in G_\delta}} \max_{0 \leq x \leq \pi} |Z_1(x, \lambda)| \leq C_\varepsilon.$$

Since  $\varepsilon$  is an arbitrary positive number, we obtain the validity of equality (4.15). Relations (4.14), (4.15) immediately imply (4.13), thus, statement (2) of theorem is proved.

System of eigenfunction  $\{\varphi(x, \lambda_n)\}_{n \geq 1}$  is complete and orthogonal in  $L_2(0, \pi; \rho)$ . Therefore, it forms the orthogonal basis in  $L_{2, \rho}(0, \pi)$  and Parseval equality from theorem is valid.  $\square$

## 5. WEYL SOLUTION, WEYL FUNCTION

Let  $\Phi(x, \lambda)$  be the solution of equation (1.1) that satisfied the conditions  $\Phi'(0, \lambda) = 1$ ,  $\Phi(\pi, \lambda) = 0$ . Denote by  $C(x, \lambda)$  the solution of equation (1.1), which satisfied the initial conditions  $C(0, \lambda) = 0$ ,  $C'(0, \lambda) = 1$ . Then, the solution  $\psi(x, \lambda)$  can be represented as follows

$$(5.1) \quad \psi(x, \lambda) = \psi(0, \lambda)\varphi(x, \lambda) - \Delta(\lambda)C(x, \lambda)$$

or

$$(5.2) \quad -\frac{\psi(x, \lambda)}{\Delta(\lambda)} = C(x, \lambda) - \frac{\psi(0, \lambda)}{\Delta(\lambda)}\varphi(x, \lambda).$$

Denote

$$(5.3) \quad M(\lambda) := -\frac{\psi(0, \lambda)}{\Delta(\lambda)}.$$

It is clear that

$$(5.4) \quad \Phi(x, \lambda) = C(x, \lambda) + M(\lambda)\varphi(x, \lambda).$$

The function  $\Phi(x, \lambda)$  and  $M(\lambda) = \Phi(0, \lambda)$  are respectively called the Weyl solution and the Weyl function of the boundary value problem (1.1), (1.2). The Weyl function is a meromorphic function having simple poles at points  $\lambda_n$  eigenvalues of boundary value problem (1.1), (1.2). Relations (5.2), (5.4) yield

$$(5.5) \quad \Phi(x, \lambda) = -\frac{\psi(x, \lambda)}{\Delta(\lambda)}.$$

It can be shown that

$$(5.6) \quad \langle \varphi(x, \lambda), \Phi(x, \lambda) \rangle = 1.$$

**Theorem 5.1.** *If  $M(\lambda) = \tilde{M}(\lambda)$ , then  $L = \tilde{L}$ ; that is, the boundary value problem (1.1), (1.2), is unique by the Weyl function.*

*Proof.* We describe the matrix  $P(x, \lambda) = [P_{ij}(x, \lambda)]_{i,j=1,2}$  with the formula

$$(5.7) \quad P(x, \lambda) \begin{pmatrix} \tilde{\varphi}(x, \lambda) & \tilde{\Phi}(x, \lambda) \\ \tilde{\varphi}'(x, \lambda) & \tilde{\Phi}'(x, \lambda) \end{pmatrix} = \begin{pmatrix} \varphi(x, \lambda) & \Phi(x, \lambda) \\ \varphi'(x, \lambda) & \Phi'(x, \lambda) \end{pmatrix}.$$

From (5.7), we have

$$(5.8) \quad \begin{aligned} \varphi(x, \lambda) &= P_{11}(x, \lambda)\tilde{\varphi}(x, \lambda) + P_{12}(x, \lambda)\tilde{\varphi}'(x, \lambda), \\ \Phi(x, \lambda) &= P_{11}(x, \lambda)\tilde{\Phi}(x, \lambda) + P_{12}(x, \lambda)\tilde{\Phi}'(x, \lambda), \end{aligned}$$

or

$$(5.9) \quad \begin{aligned} P_{11}(x, \lambda) &= \varphi(x, \lambda)\tilde{\Phi}'(x, \lambda) - \Phi(x, \lambda)\tilde{\varphi}'(x, \lambda), \\ P_{12}(x, \lambda) &= -\varphi(x, \lambda)\tilde{\Phi}(x, \lambda) + \Phi(x, \lambda)\tilde{\varphi}(x, \lambda). \end{aligned}$$

Taking equation (5.5) into consideration in (5.9), we get (5.4) into (5.9), then we get

$$(5.10) \quad \begin{aligned} P_{11}(x, \lambda) &= 1 + \frac{1}{\Delta(\lambda)} \left[ \varphi(x, \lambda)(\tilde{\psi}'(x, \lambda) - \psi'(x, \lambda)) - \psi(x, \lambda)(\tilde{\varphi}'(x, \lambda) - \varphi'(x, \lambda)) \right] \\ P_{12}(x, \lambda) &= \frac{1}{\Delta(\lambda)} \left[ \psi(x, \lambda)\tilde{\varphi}(x, \lambda) - \varphi(x, \lambda)\tilde{\psi}(x, \lambda) \right]. \end{aligned}$$

Now, from (4.8) and (4.9), we have from equation (5.10)

$$(5.11) \quad \lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} |P_{11}(x, \lambda) - 1| = \lim_{|\lambda| \rightarrow \infty} \max_{0 \leq x \leq \pi} |P_{12}(x, \lambda)| = 0.$$

Now, if we take into consideration equation (5.4) into (5.9), we get

$$\begin{aligned} P_{11}(x, \lambda) &= \varphi(x, \lambda) \tilde{C}'(x, \lambda) - C(x, \lambda) \tilde{\varphi}'(x, \lambda) + (\tilde{M}(\lambda) - M(\lambda)) \varphi(x, \lambda) \tilde{\varphi}'(x, \lambda) \\ P_{12}(x, \lambda) &= C(x, \lambda) \tilde{\varphi}(x, \lambda) - \tilde{C}(x, \lambda) \varphi(x, \lambda) - (\tilde{M}(\lambda) - M(\lambda)) \varphi(x, \lambda) \tilde{\varphi}(x, \lambda). \end{aligned}$$

Therefore if  $M(\lambda) = \tilde{M}(\lambda)$ , then  $P_{11}(x, \lambda)$  and  $P_{12}(x, \lambda)$  are entire functions for every fixed  $x$ . It can be easily seen from (5.11) that  $P_{11}(x, \lambda) = 1$  and  $P_{12}(x, \lambda) = 0$ . Substituting into (5.8), we get  $\varphi(x, \lambda) \equiv \tilde{\varphi}(x, \lambda)$  and  $\Phi(x, \lambda) \equiv \tilde{\Phi}(x, \lambda)$  for every  $x$  and  $\lambda$ . Hence, we arrive at  $q(x) \equiv \tilde{q}(x)$ .  $\square$

**Theorem 5.2.** *The expression*

$$(5.12) \quad M(\lambda) = - \sum_{n=0}^{\infty} \frac{1}{2\alpha_n \lambda_n (\lambda - \lambda_n)}$$

holds.

*Proof.* Using (5.3), we get for sufficiently large  $\lambda^* > 0$ ,

$$(5.13) \quad M(\lambda) \leq \frac{C_\delta}{|\lambda|}, \quad \lambda \in G_\delta, \quad |\lambda| > \lambda^*.$$

Further using (1.9) and (5.4) we calculate:

$$(5.14) \quad \operatorname{Res}_{\lambda=\lambda_n} M(\lambda) = - \frac{\psi(0, \lambda_n)}{\Delta(\lambda_n)} = - \frac{\beta_n}{\Delta(\lambda_n)} = - \frac{1}{2\lambda_n \alpha_n}.$$

Now, let's consider the contour integral

$$J_N(\lambda) = \frac{1}{2\pi i} \int_{\Gamma_N} \frac{M(\mu)}{\lambda - \mu} d\mu, \quad \lambda \in \operatorname{Int} \Gamma_N,$$

where  $\Gamma_N = \{\mu : |\mu| = |\lambda_N^0| + \frac{\gamma}{2}\}$  is a contour of counter-clockwise by pass.

By virtue of (5.13) we have  $\lim_{N \rightarrow \infty} J_N(\lambda) = 0$ . On the other hand, by residue theorem and (5.14) yield

$$J_N(\lambda) = -M(\lambda) - \sum_{n=-N}^N \frac{1}{2\lambda_n \alpha_n (\lambda - \lambda_n)}$$

and when  $N \rightarrow \infty$  we arrive at (5.12).  $\square$

**Theorem 5.3.** *If  $\lambda_n = \tilde{\lambda}_n, \alpha_n = \tilde{\alpha}_n$  for all  $n \in Z$  then  $L = \tilde{L}$ . That is, the problem (1.1), (1.2) is uniquely determined by spectral date.*

*Proof.* Since  $\lambda_n = \tilde{\lambda}_n, \alpha_n = \tilde{\alpha}_n$  for all  $n \in Z$  and considering the formula (5.12), we have  $M(\lambda) = \tilde{M}(\lambda)$ . Using Theorem 4,  $L = \tilde{L}$  is obtained.  $\square$



## REFERENCES

- [1] A. N. Tikhonov, A. A. Samarskii *Equation of Mathematical Physics* Dover Books on Physics and Chemistry. Dover. New York. 1990.
- [2] M. L. Rasulov *Methods of Contour Integration* Series in Applied Mathematics and Mechanics. v.3. Nort-Holland. Amsterdam. 1967.
- [3] G. Freiling, V. Yurko *Inverse Sturm- Liouville Problems and Their Applications* Nova Science Publishers. Inc. 2008.
- [4] A. Zettl *Sturm- Liouville Theory* Mathematical Surveys and Monographs. v.121. Am. Math. Soc. Providence. 2005.
- [5] A. M. Akhtyamov, A. V. Mouftakhov *Identification of boundary conditions using natural frequencies* Inverse Problems in Science and Engineering. 2004. 12(4). 393-408.
- [6] O. H. Hald *Discontinuous inverse eigenvalue problems* Comm. Pure Appl. Math. 1984. 37. 539-577.
- [7] R. Carlson *An inverse spectral problem for Sturm- Liouville operators with discontinuous coefficients* Pro. Amer. Math. Soc. 1994. 120(2). 5-9.
- [8] E. N. Akhmedova *On representation of a solution of Sturm- Liouville equation with discontinuous coefficients* Proceedings of IMM of NAS of Azarbaijan. 2002. 16(24). 5-9.
- [9] N. Altinisik, M. Kadakal, O. Mukhtarov *Eigenvalues and eigenfunctios of discontinuos Sturm-Liouville problems with eigenparameter dependent boundary conditions* Acta Math. Hung. 2004. 102(1-2). 159-175.
- [10] Kh. R. Mamedov *On a basis problem for a second order differential equation with a discontinuous coefficient and a spectral parameter in the boundary conditions* Geometry, Integrability and Quantization. 2006. 7. 218-225.
- [11] A. R. Aliev *Solvability of a class of boundary value problems for second-order operator-differential equations with a discontinuous coefficient in a weighted space* Differential Equations. 2007. 43(10)1459-1463.
- [12] A. A. Sedipkov *The inverse spectral problem for the Sturm-Liouville operator with discontinuous potantial* J. Inverse III-Posed Probl. 2012. 20. 139-167.
- [13] C. F. Yang *Inverse nodal problems of discontinuous Sturm-Liouville operator* Journal of Differential Equations. 2013. 254(4). 1992-2014.
- [14] B. Aliev, Y. S. Yakubov *Solvability of boundary value problems for second-order elliptic differential-operator equations with a spectral parameter and with a discontinuous coefficient at the highest derivative* Differential Equations. 2014. 50(4). 464-475.
- [15] Kh. R. Mamedov, F. A. Cetinkaya *An uniqueness theorem for a Sturm-Liouville equation with spectral parameter in boundary conditions* Appl. Math. Inf. Sci. 2015. 9(2). 981-988.
- [16] M. G. Gasymov *The direct and inverse problem of spectral analysis for a class of equations with a discontinuous coefficient* (Ed. M. M. Lavrent'ev), Non-Classical Methods in Geophysics, Nauka (Novosibirsk, Russia). 1977. 37-44.
- [17] Kh. R. Mamedov *Uniqueness of solution of the inverse problem of scattering theory with a spectral parameter in the boundary condition* Math. Notes 2003. 74 (1), 136-140.
- [18] Kh. R. Mamedov *On an inverse scattering problem for a discontinuous Sturm-Liouville equation with a spectral parameter in boundary condition* Bound. Value Prob. 2010. Article ID 171967, 1-17.

SCIENCE AND LETTER FACULTY, MATHEMATICS DEPARTMENT, MERSIN UNIVERSITY, 333343, MERSIN, TURKEY

*E-mail address:* hanlar@mersin.edu.tr

SCIENCE AND LETTERS FACULTY, MATHEMATICS DEPARTMENT, HARRAN UNIVERSITY, SANLIURFA, TURKEY

*E-mail address:* dkarahan@harran.edu.tr

IFSCOM2016 <sup>1</sup> PROCEEDING BOOK

No. 1 pp. 54-57 (2016)

ISBN: 978-975-6900-54-3

## ABOUT CERTAIN HOMOLOGICAL PROPERTIES OF SYMMETRIC DERIVATIONS OF KÄHLER MODULES

HAMIYET MERKEPCI, NECATI OLGUN, AND ELA AYDIN

ABSTRACT. In this study, we express more informations concerning with homological properties of symmetric derivations of Kahler modules acquainted by H. Osborn in [1].

Received: 19–August–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

The definition of  $n$ -th order symmetric derivations of Kähler modules were given by H. Osborn at 1965 in [1]. J. Johnson made known the structures of differential module on certain modules of Kähler differentials in [3]. Then, advanced principal theories about the calculus of high order derivations and a few functorial features of high order differential modules were presented by Y. Nakai in [4]. Higher derivations and universal differential operators of Kähler modules were studied by R. Hart in [2]. Olgun defined generalized symmetric derivations on high order Kähler modules in [6]. Komatsu presented right differential operators on a noncommutative ring extension in [10]. The more informations about these subjects were found in [5,7,8,9,11]. The aim of this study is to investigate these homological structures and is to give more knowldege about them.

### 2. PRELIMINARY

Throughout this paper we assume  $R$  be a commutative algebra over an algebraically closed field  $k$  with characteristic zero. When  $R$  is a  $k$ -algebra,  $J_n(R)$  denotes the universal module of  $n$ -th order differentials of  $R$  over  $k$  and  $\Omega_n(R)$  be the module of  $n$ -th order Kähler differentials of  $R$  over  $k$  and  $d_n$  be the canonical  $n$ -th order  $k$ -derivation  $R \rightarrow \Omega_n(R)$  of  $R$ . The pair  $\{\Omega_n(R), d_n\}$  has the universal mapping property with regard to the  $n$ -th order  $k$ -derivations of  $R$ .  $\Omega_n(R)$  is generated by the set  $\{d_n(r) : r \in R\}$ .

**Definition 2.1.** Let  $R$  be a commutative algebra over a field  $k$  of characteristic zero,  $A$  be an  $R$ -module,  $A \otimes_R A$  be the tensor product of  $A$  with itself and let  $K$  be the submodule of  $A \otimes_R A$  generated by the elements of the form  $a \otimes b - b \otimes a$  where  $a, b \in A$ . Consider the factor module  $\vee^2 A = A \otimes A / K$ . The module  $\vee^2 A$  is said to be the second symmetric power of  $A$ . The canonic balanced map is defined such that

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
*Key words and phrases.* Kähler module, symmetric derivation, projective dimension, hypersurface.

$$\begin{aligned}\otimes : A \times A &\longrightarrow A \otimes A \\ \otimes(a, b) &= a \otimes b\end{aligned}$$

and a natural surjective map defined such that  $\gamma : A \otimes A \longrightarrow \vee^2 A$ . Then the composite map is bilinear and called  $\gamma \otimes = \vee$ .

**Lemma 2.1.** *Let  $A$  and  $B$  be  $R$ -modules and let  $\zeta : A \times A \longrightarrow B$  be a bilinear alternating map. Then there exists an  $R$ -module homomorphism  $f : \vee^2 A \longrightarrow B$  such that the diagram*

$$\begin{array}{ccc} A \times A & \xrightarrow{\zeta} & B \\ \vee \searrow & & \nearrow f \\ & \vee^2 A & \end{array}$$

commutes.

**Definition 2.2.** Let  $R$  be any  $k$ -algebra (commutative with unit),  $R \rightarrow \Omega_1(R)$  be first order Kähler derivation of  $R$  and let  $\vee(\Omega_1(R))$  be the symmetric algebra  $\bigoplus_{p \geq 0} \vee^p(\Omega_1(R))$  generated over  $R$  by  $\Omega_1(R)$ .

A symmetric derivation is any linear map  $D$  of  $\vee(\Omega_1(R))$  into itself such that

- i)  $D(\vee^p(\Omega_1(R))) \subset \vee^{p+1}(\Omega_1(R))$
- ii)  $D$  is a first order derivation over  $k$  and
- iii) the restriction of  $D$  to  $R$  ( $R \simeq \vee^0(\Omega_1(R))$ ) is the Kähler derivation

$d_1 : R \rightarrow \Omega_1(R)$ .

**Definition 2.3.** Let  $R$  be any  $k$ -algebra (commutative with unit),  $R \rightarrow \Omega_n(R)$  be  $n$ -th order Kähler derivation of  $R$  and let  $\vee(\Omega_n(R))$  be the symmetric algebra  $\bigoplus_{p \geq 0} \vee^p(\Omega_n(R))$  generated over  $R$  by  $\Omega_n(R)$ .

A generalized symmetric derivation is any  $k$ -linear map  $D$  of  $\vee(\Omega_n(R))$  into itself such that

- i)  $D(\vee^p(\Omega_n(R))) \subset \vee^{p+1}(\Omega_n(R))$
- ii)  $D$  is a  $n$ -th order derivation over  $k$  and
- iii) the restriction of  $D$  to  $R$  ( $R \simeq \vee^0(\Omega_n(R))$ ) is the Kähler derivation

$d_n : R \rightarrow \Omega_n(R)$ .

**Proposition 2.1.** *Let  $R = k[x_1, \dots, x_s]$  be a polynomial algebra of dimension  $s$ .*

*Then  $\Omega_n(R)$  is a free  $R$ -module of rank  $\binom{n+s}{s} - 1$  with basis*

$$\{ d_n(x_1^{i_1} \dots x_s^{i_s}) : i_1 + \dots + i_s \leq n \}$$

$$\vee^2(\Omega_n(R)) \text{ is a free } R\text{-module of rank } \binom{t+1}{t-1}$$

where  $t = \binom{n+s}{s} - 1$  with basis  $\{ d_n(x_1^{i_1} \dots x_s^{i_s}) \otimes d_n(x_1^{i_1} \dots x_s^{i_s}) : i_1 + \dots + i_s \leq n \}$

### 3. SYMMETRIC POWERS OF KÄHLER MODULES

In this section, we consider the tensor, exterior and symmetric algebras of Kähler modules and define the symmetric powers of a given module  $A$  over a  $k$ -algebra and a few elementary properties.

**Definition 3.1.** Let  $A$  be a  $R$ -module.

i) By  $\otimes^n A$  we shall denote the  $R$ -module with a universal  $R$ -bilinear map of  $A^n \rightarrow \otimes^n A$  written  $(x_1, \dots, x_n) \rightarrow x_1 \otimes \dots \otimes x_n$ . This module is called the  $n$ -fold tensor power of  $A$ .

ii) By  $\Lambda^n A$  we shall denote the  $R$ -module with a universal alternating  $R$ -bilinear map of  $A^n \rightarrow \Lambda^n A$  written  $(x_1, \dots, x_n) \rightarrow x_1 \wedge \dots \wedge x_n$ . This module is called the  $n$ -fold exterior power of  $A$ .

iii) By  $\vee^n A$  we shall denote the  $R$ -module with a universal symmetric  $R$ -bilinear map of  $A^n \rightarrow \vee^n A$  written  $(x_1, \dots, x_n) \rightarrow x_1 \dots x_n$ . This module is called the  $n$ -fold symmetric power of  $A$ .

Let us the conventin that  $\otimes^1 A, \Lambda^1 A$  and  $\vee^1 A$  are all identified with  $A$ , while  $\otimes^0 A, \Lambda^0 A$  and  $\vee^0 A$  are all identified with  $R$ .

**Theorem 3.1.** *Let  $A$  be a free  $R$ -module on a basis  $X = \{x_1, \dots, x_d\}$ . If  $A$  is generated by  $x_1, \dots, x_d$  then  $A^{\otimes k}$  is generated by  $x_{i_1} \otimes \dots \otimes x_{i_k}$  as an  $R$ -module. Where  $1 \leq i_1, \dots, i_k \leq d$  and  $\dim_R(\otimes^k A) = d^k$ .*

*Since  $\Lambda^k(A)$  is factor module of  $A^{\otimes k}$ , so  $\Lambda^k(A)$  is generated by  $x_{i_1} \wedge \dots \wedge x_{i_k}$  as an  $R$ -module where  $1 \leq i_1, \dots, i_k \leq d$ . For any  $x_1, x_2 \in X$ , it satisfied  $x_1 \wedge x_1 = 0$  and  $x_1 \wedge x_2 + x_2 \wedge x_1 = 0$ . If  $0 \neq A$  is affine free with  $x_1, \dots, x_d$  then  $x_{i_1} \wedge \dots \wedge x_{i_k}$  is basis for  $\Lambda^k A$  and  $\dim_R(\Lambda^k A) = \binom{d}{k}$ .*

$\vee A$  may be presented by the generating set  $X$ , and relation  $xy = yx$  ( $x, y \in X$ ) and is the (commutative) polynomial algebra  $R[X]$ . Then an  $R$ -module basis for  $\vee A$  is given by those products  $x_1 \dots x_n$  with  $x_1 \leq \dots \leq x_n \in X$ . If  $X$  is a finite set  $\{x_1, \dots, x_r\}$ , then the elements of this basis can be written  $x_1^{i_1} \dots x_r^{i_r}$  with  $i_1, \dots, i_r \geq 0$ , and for each  $n$ ,  $\dim(\vee^n A) = \binom{r+n-1}{r-1}$ .

#### 4. HOMOLOGICAL PROPERTIES OF SYMMETRIC DERIVATIONS

**Theorem 4.1.** *Let  $R$  be an affine  $k$ -algebra. Then we have a long exact sequence of  $R$ -modules*

$$0 \rightarrow \ker \eta \rightarrow \Omega_{(2n)}(R) \xrightarrow{\eta} J_n(\Omega_n(R)) \rightarrow \operatorname{coker} \eta \rightarrow 0.$$

for all  $n \geq 0$ .

**Example 4.1.**  $R = k[a, b]$  be a polynomial algebra of dimension 2. Then  $\Omega_1(R)$  is a free  $R$ -module of rank 2 with basis  $\{d_1(a), d_1(b)\}$  and  $\Omega_2(R)$  is a free  $R$ -module of rank 5 with basis  $\{d_2(a), d_2(b), d_2(a^2), d_2(ab), d_2(b^2)\}$ .

$J_1(\Omega_1(R))$  is a free  $R$ -module generated by  $\{\Delta_1(d_1(a)), \Delta_1(d_1(b)), \Delta_1(ad_1(a)), \Delta_1(ad_1(b)), \Delta_1(bd_1(a)), \Delta_1(bd_1(b))\}$

**Theorem 4.2.** *Let  $R$  be an affine  $k$ -algebra. Then we have a long exact sequence of  $R$ -modules*

$$0 \rightarrow \ker \gamma \rightarrow J_n(\Omega_n(R)) \xrightarrow{\gamma} \vee^2(\Omega_n(R)) \rightarrow \operatorname{coker} \gamma \rightarrow 0.$$

for all  $n \geq 0$ .

**Lemma 4.1.** *Let  $R$  be an affine domain. Then  $\Omega_n(R)$  is a free  $R$ -module if and only if  $\vee^2(\Omega_n(R))$  is a free  $R$ -module.*

**Theorem 4.3.** *Let  $R$  be an affine  $k$ -algebra and  $\vee(\Omega_1(R))$  has at least one symmetric derivations.  $\Omega_1(R)$  is a projective  $R$ -module if and only if  $\Omega_2(R)$  is a projective  $R$ -module.*

**Corollary 4.1.** *Let  $R$  be an affine local  $k$ -algebra and  $\vee(\Omega_1(R))$  has at least one symmetric derivation.  $\Omega_1(R)$  is a free  $R$ -module if and only if  $\Omega_2(R)$  is a free  $R$ -module.*

## REFERENCES

- [1] Osborn, H., Modules of Differentials II, Math. Ann.175, 146-158, (1968).
- [2] Hart, R., Higher Derivations and Universal Differentials, Journal of Alg. 184, 175-181, 1996.
- [3] Johnson J., Kähler Differentials and differential algebra, The Annals of Mathematics, Second series, Vol. 89, No 1, 92-98, (1969).
- [4] Nakai, Y., High Order Derivations, Osaka J. Math. 7, 1-27, (1970).
- [5] Erdogan, A., Differential Operators and Their Universal Modules, Phd. Thesis, Universty of Leeds, (1993).
- [6] Olgun N., Symmetric Derivations On *Kähler* Modules, [arXiv:1602.01343](https://arxiv.org/abs/1602.01343) [math.AC]
- [7] Bourbaki, N., Algebra, Hermann, Paris, 1974.
- [8] Sweedler, M.E. and Heyneman, R.G., Affine Hopf Algebras, J.Algebra 13, 192-241, (1969).
- [9] Vasconcelos, W.V., A note On Normality And The Module of Differentials, Math. Z., 105, 291-293, 1968.
- [10] Komatsu, H., High order Kähler Modules of Noncommutative ring extensions, Comm.Algebra 29, 5499-5524, (2001)
- [11] Bergman, G., Tensor algebras, exterior algebras and symmetric algebras, <http://math.berkeley.edu/gbergman>, (1997).

DEPARTMENT OF MATHEMATICS, CUKUROVA UNIVERSITY, ADANA  
*E-mail address:* [hamiyetmerkepçi@hotmail.com](mailto:hamiyetmerkepçi@hotmail.com)

DEPARTMENT OF MATHEMATICS, GAZIANTEP UNIVERSITY, GAZIANTEP  
*E-mail address:* [olgun@gantep.edu.tr](mailto:olgun@gantep.edu.tr)

DEPARTMENT OF MATHEMATICS, CUKUROVA UNIVERSITY, ADANA  
*E-mail address:* [eyadin@cu.edu.tr](mailto:eyadin@cu.edu.tr)

## ON A NONLOCAL BOUNDARY VALUE PROBLEM

OLGUN CABRI AND KHANLAR R. MAMEDOV

ABSTRACT. In this study, parabolic partial differential equation with two integral boundary conditions are considered for distribution of family savings for a family set. By separation of variables method, eigenvalues and eigenfunctions of the problem are obtained and solution is written. Moreover, Method of lines method and Crank Nicolson method are applied to the problem and errors of numerical methods are presented.

Received: 08–August–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Integral boundary conditions for parabolic equations are well known problem in applications (see, for example, Cannon[1], Ionkin[7], Kamynin[8], Day[3], Erofeeko and Kozlovski[6]). Such a boundary condition are called nonlocal boundary condition or nonclassical boundary condition. Similar problems are also used for hyperbolic equations.

In this study, we deal with a family saving model which can be represented by Kolmogorov equation with two integral boundary conditions.

Suppose that  $x(t)$  denotes the saving of a family at time  $t$  and satisfy the differential equation

$$(1.1) \quad dx = F(x, t) dt + G(x, t) dX, \quad G \geq 0$$

where  $X$  is the Markov process,  $F(x, t)$  is the rate of the change for the family saving and  $G(x, t) dX$  is the random change of the family income.

For a family set let us assume that equation (1.1) describes the saving of all families by ignoring the dynamic of individual family saving. The density distribution of the saving of families  $u(x, t)$  satisfies

$$(1.2) \quad \frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} ((c(x, t) + F(x, t)) u) + \frac{1}{2} \frac{\partial^2}{\partial x^2} (b(x, t)u) + f(x, t)$$

with initial condition

$$(1.3) \quad u(x, 0) = \varphi(x), \quad 0 \leq x \leq l$$

and boundary conditions

$$(1.4) \quad \int_0^l u(x, t) dx = N(t), \quad t \geq 0$$

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
*Key words and phrases.* Nonlocal Boundary Condition, Family Saving Model, Method of Lines Method, Crank Nicolson Method.

$$(1.5) \quad \int_0^l x u(x, t) dx = K(t), t \geq 0$$

where  $c(x, t)$ ,  $b(x, t)$ ,  $K(t)$ ,  $N(t)$ ,  $\varphi(x)$  and  $f(x, t)$  are continuously differentiable functions.  $N(t)$ ,  $K(t)$  denote total number of families and total amount of family saving in  $[0, l]$  respectively [6].

## 2. SPECIAL CASE OF THE MODEL

We will consider special case of problem (1.2)-(1.5) on region  $D = (0 < t < \infty) \times (0 < x < l)$

$$(2.1) \quad \frac{\partial u}{\partial t} = a^2 \frac{\partial^2 u}{\partial x^2} + f(x, t),$$

$$(2.2) \quad u(x, 0) = \varphi(x), 0 \leq t \leq T,$$

$$(2.3) \quad \int_0^1 u(x, t) dx = N(t), t \geq 0,$$

$$(2.4) \quad \int_0^1 x u(x, t) dx = K(t), t \geq 0,$$

where  $f(x, t)$ ,  $K(t)$ ,  $N(t)$ ,  $\varphi(x)$  are continuously differentiable functions on region  $D$ . Compatibility conditions of this problem are

$$\int_0^1 x \varphi(x) dx = N(0) \text{ and } \int_0^1 \varphi(x) dx = K(0).$$

Using the transform

$$u(x, t) = v(x, t) + (12K(t) - 6N(t))x + 4N(t) - 6K(t)$$

boundary conditions of equation (2.1)-(2.4) become homogenous:

$$(2.5) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2} + F(x, t),$$

$$(2.6) \quad v(x, 0) = \psi(x),$$

$$(2.7) \quad \int_0^1 v(x, t) dx = 0,$$

$$(2.8) \quad \int_0^1 x v(x, t) dx = 0,$$

where

$$F(x, t) = f(x, t) - (12K'(t) - 6N'(t))x + 4N'(t) - 6K'(t)$$

and

$$\psi(x) = \varphi(x) - (12K(0) - 6N(0))x + 4N(0) - 6K(0).$$

Equations (2.5)-(2.8) are linear with respect to  $v(x,t)$ , then this problem can split into two auxiliary problems:

i)

$$(2.9) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2},$$

$$(2.10) \quad v(x, 0) = \psi(x),$$

$$(2.11) \quad \int_0^1 v(x, t) dx = 0$$

$$(2.12) \quad \int_0^1 xv(x, t) dx = 0$$

ii)

$$(2.13) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2} + F(x, t),$$

$$(2.14) \quad v(x, 0) = 0,$$

$$(2.15) \quad \int_0^1 v(x, t) dx = 0$$

$$(2.16) \quad \int_0^1 xv(x, t) dx = 0$$

If solution of the problem (i) is  $v_1(x, t)$  and solution of the problem (ii) is  $v_2(x, t)$  then solution of the problem (2.5)-(2.8) is  $v(x, t) = v_1(x, t) + v_2(x, t)$ .

Integrating both sides of (2.9) with respect to  $x$  from 0 to 1 and using integration by parts, integral boundary conditions in (2.11) and (2.12) become, respectively,

$$v_x(1, t) - v_x(0, t) = 0,$$

$$v_x(1, t) - v(1, t) + v(0, t) = 0.$$

Substituting these equations in (2.9)-(2.12), we have

$$(2.17) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2},$$

$$(2.18) \quad v(x, 0) = \psi(x),$$

$$(2.19) \quad v_x(1, t) - v_x(0, t) = 0,$$

$$(2.20) \quad v_x(1, t) - v(1, t) + v(0, t) = 0.$$



By the separation of variables, a Sturm-Liouville problem and an ODE are, respectively, obtained as

$$(2.21) \quad X''(x) + \lambda X(x) = 0,$$

$$(2.22) \quad X'(1) - X'(0) = 0,$$

$$(2.23) \quad X'(1) - X(1) + X(0) = 0,$$

and

$$(2.24) \quad T'(t) + \lambda a^2 T(t) = 0.$$

Sturm-Liouville problem (2.21)-(2.23) is self adjoint and boundary conditions are regular, and also strongly regular. Therefore, the eigenfunctions of the Sturm-Liouville problem are the Riesz basis on  $L^2[0, 1]$  (Naimark[11], Kesselman[9], Mikhailov [10]).

Characteristic equation of the Sturm-Liouville problem is

$$(2.25) \quad 2 - 2 \cos k - k \sin k = 0,$$

where  $\sqrt{\lambda} = k$ .

It is easily seen that  $k_0 = 0$  and  $k_{2n} = 2n\pi$ , ( $n = 1, 2, \dots$ ) are roots of the equation (2.25). There is also another root of equation (2.25) in  $\left[n\pi, \frac{(2n+1)}{2}\pi\right]$ . By using Langrange-Burmam formula root is calculated asymptotically as

$$\begin{aligned} k_{2n+1} &= (2n+1)\pi - 4((2n+1)\pi)^{-1} - \frac{32}{3}((2n+1)\pi)^{-3} - \frac{832}{15}((2n+1)\pi)^{-5} \\ &+ O\left(\frac{1}{n^7}\right). \end{aligned}$$

Corresponding eigenfunctions are obtained by

$$X_0(x) = 1,$$

$$X_{2n} = \cos(2\pi n)x, \quad n = 1, 2, \dots$$

$$X_{2n+1} = \frac{-k_n}{2} \cos(k_n x) + \sin(k_n x), \quad n = 1, 2, \dots$$

Therefore, solution of the problem (2.17)-(2.20) is

$$v_1(x, t) = \sum_{n=0}^{\infty} A_{2n} \cos(2\pi n x) e^{-a^2 4\pi^2 n^2 t} + \sum_{n=1}^{\infty} B_n \left( \frac{-k_n}{2} \cos(k_n x) + \sin(k_n x) \right) e^{-a^2 k_n^2 t},$$

where

$$A_0 = \int_0^1 \psi(x) dx,$$

$$A_n = 2 \int_0^1 \psi(x) \cos(2\pi n x) dx, \quad n = 1, 2, \dots$$

$$B_n = \frac{1}{\|X_{2n+1}(x)\|^2} \int_0^1 \psi(x) \left( \frac{-k_n}{2} \cos(k_n x) + \sin(k_n x) \right) dx, \quad n = 1, 2, \dots$$

Solution of the problem (2.13)-(2.16) can be easily obtained by

$$v_2(x, t) = \sum_{n=0}^{\infty} \left[ \int_0^t F_{2n}(\tau) e^{-k_n^2(t-\tau)} d\tau \right] X_{2n}(x) \\ + \left[ \int_0^t F_{2n+1}(\tau) e^{-k_n^2(t-\tau)} d\tau \right] X_{2n+1}(x),$$

where

$$F_0(\tau) = \int_0^1 F(x, \tau) x dx, \\ F_{2n}(\tau) = \int_0^1 F(x, \tau) X_{2n}(x) dx, \quad n = 1, 2, \dots \\ F_{2n+1}(\tau) = \int_0^1 F(x, \tau) X_{2n+1}(x) dx. \quad n = 1, 2, \dots$$

### 3. NUMERICAL SOLUTION

Method of Lines [12] and the Crank-Nicolson method [13] are used for numerical solution of problem (2.1)-(2.4). In both methods, the Simpson's rule is used to approximate the integral in (2.3) and (2.4) numerically. We display here a few of numerical results.

#### Example 3.1.

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + (x^2 - 2)e^t,$$

$$u(x, 0) = x^2,$$

$$\int_0^1 u(x, t) dx = (1/6) - 2t,$$

$$\int_0^1 xu(x, t) dx = (1/12) - t.$$

Exact solution of example 1 is  $u(x, t) = x - x^2 - 2t$ . The absolute relative errors at various spatial lengths for  $u(0.5, 0.5)$  are shown in Table 1.

Relative Error at  $u(0.5, 0.5)$  in Example 1

Spatial Length	MOL Method	Crank-Nicolson Method
h=0.1	1.3471E-14	2.9606E-16
h=0.05	8.4510E-13	2.9606E-16
h=0.025	1.7494E-12	6.8094E-15
h=0.0125	4.6876E-12	1.9244E-15

**Example 3.2.**

$$\begin{aligned}\frac{\partial u}{\partial t} &= \frac{\partial^2 u}{\partial x^2}, \\ u(x, 0) &= \sin(\pi x), \\ \int_0^1 u(x, t) dx &= \frac{2}{\pi} \exp(-\pi^2 t), \\ \int_0^1 x u(x, t) dx &= (1/12) - t.\end{aligned}$$

Exact solution of example 2 is  $u(x, t) = \sin(\pi x) \exp(-\pi^2 t)$ . The absolute relative errors at various spatial lengths for  $u(0.5, 0.5)$  are shown in Table 2.

Relative Error at  $u(0.5, 0.5)$  in Example 2

Spatial Length	MOL Method	Crank-Nicolson Method
h=0.1	0.0029	0.0075
h=0.05	4.5074E-4	0.0023
h=0.025	6.4370E-5	5.6888E-4
h=0.0125	4.4595E-6	9.1160E-5

## 4. CONCLUSION

Diffusion equation with two integral boundary conditions is studied. Integral boundary conditions are transformed to local one and by separation of variables, analytic solution of this problem is found. In addition, by applying the Method of Lines [12] and Crank Nicolson method [13], numerical solution of the problem is found.

## REFERENCES

- [1] Cannon, J.R., *The solution of the heat equation subject to specification of energy*, Quarterly of Applied Mathematics, Vol. **21**, 155-160, 1963.
- [2] Cannon, J. R. ve Van Der Hoek, J., *An implicit finite difference scheme for the diffusion of mass in a portion of the domain*, Numerical Solutions of Partial Differential Equations (J. Noye, Ed), 527-539. 1982
- [3] Day, W.A., *Extension of a property of the equation to linear thermoelasticity and other theories*, Quarterly of Applied Mathematics, 1982, 40.3: 319-330.
- [4] Dehghan, M. , *On the numerical solution of the diffusion equation with a nonlocal boundary condition*, Mathematical Problems in Engineering, 2003(2), 81-92, 2003
- [5] Ekolin, G., *Finite difference methods for a nonlocal boundary value problem for the heat equation*, BIT Numerical Mathematics, 1991, 31.2: 245-261.
- [6] Erofeenko, V.T. ve Kozlovski, I.Y., *Equation Partial Differential Equation and Mathematical Models in Economic*, URRS, 2011
- [7] Ionkin N. I, *Solution of a boundary-value problem in heat conduction with a nonclassical boundary condition*, Diff. Eqs., Vol. 13, No. 2,1977, pp. 294-304.
- [8] Kamynin, L. I., *A boundary value problem in the theory of heat conduction with a nonclassical boundary condition*, USSR Computational Mathematics and Mathematical Physics, 1964, 4.6: 33-59.
- [9] Kesselman, G. M., *On the unconditional convergence of eigenfunction expansions of certain differential operators*, Izv. Vyssh. Uchebn. Zaved. Mat. 39(2),(1964), 8293, (Russian).
- [10] Mikhailov, V. P., *On Riesz bases in  $L^2(0, 1)$* , Dokl. Akad. Nauk SSSR, 1962, 144, 981-984, (Russian)

- [11] Naimark, M.A., *Linear Differential Operator*, 1968
- [12] Rehmana, M. A. ve Taj, M. S., *A Numerical technique for heat equation subject to integral specification*, Sci.nt(Lahore),24(1),21-26,2011
- [13] Shingmin W. ve Lin, Y. *A numerical method for the diffusion equation with nonlocal boundary specifications*, International Journal of Engineering Science, 1990, 28.6: 543-546.

ARTVIN CORUH UNIVERSITY  
*E-mail address:* [olguncabri@artvin.edu.tr](mailto:olguncabri@artvin.edu.tr)

MERSIN UNIVERSITY  
*E-mail address:* [hanlarm@yahoo.com](mailto:hanlarm@yahoo.com)

## FUZZY RIGHT FRACTIONAL OSTROWSKI INEQUALITIES

MEHMET CITIL AND MUHAMMED GEZERCAN

ABSTRACT. In this paper, firstly fuzzy basic concept is studied. We investigated other Ostrowski type inequalities in literature. We obtained the very general fuzzy fractional Ostrowski type inequality with right fractional Caputo derivative using the Hölder inequality in this type.

Received: 29–June–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Mathematical inequalities take an important place among mathematical concepts. These enable us to find the values of these quantities approximately. Mathematical inequalities have also important applications in functional analysis. For example when building norms on some linear spaces.

The following result is known in the literature as an Ostrowski's inequality. In 1938, the classical integral inequality was proved by A.M. Ostrowski [9].

The inequality of Ostrowski gives us an estimation for the deviation of the values of a smooth function from its mean value. More precisely, if  $f : [a, b] \rightarrow \mathbb{R}$  is a differentiable function with bounded derivative, then

$$\left| f(x) - \frac{1}{b-a} \int_a^b f(t) dt \right| \leq \left[ \frac{1}{4} + \frac{(x - \frac{a+b}{2})^2}{(b-a)^2} \right] (b-a) \|f'\|_\infty$$

for every  $x \in [a, b]$ . Moreover the constant  $1/4$  in the right side of the inequality is the best possible value for the better result.

The theory of fractional calculus has known an intensive development over the last few decades. It is shown that derivatives and integrals of fractional type provide an adequate mathematical modelling of real objects and processes see [7] – [8].

We notice that the first generalization of Ostrowski's inequality was given by Milanovic and Pecaric in [2].

In [10] Pachpatte has proved the Ostrowski inequality in three independent variables. In the past few years, many authors have obtained various generalizations of this type of inequality and many researchers worked on a fractional form of it as well as on time scale calculus [11].

Univariate right fractional Ostrowski inequalities has been shown by Anastassiou [12].

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference

*Key words and phrases.* Fuzzy fractional Ostrowski inequality, fuzzy right Caputo fractional derivative, Hukuhara difference.

Fuzzy sets were defined in [1] A standard fuzzy set in  $X$  is characterized by a membership function  $\mu : X \rightarrow [0, 1]$  A standard fuzzy set is called normalized if  $\sup_{x \in X} \mu(x) = 1$

Fuzzy fractional calculus and the Ostrowski inequalities have been studied by Anatassiou [5].

The main purpose of this manuscript is to establish Ostrowski-type inequality involving right Caputo differentiability. First of all; we give basic information about the fuzzy set .Then ,we introduce the very general univariate fuzzy fractional Ostrowski type inequality.We show this inequality in fuzzy space.

## 2. BACKGROUND

We need the following basic concepts

**Definition 2.1.** [5] Let  $\mu : \mathbb{R} \rightarrow [0, 1]$  with the following properties

- i)  $\mu$  is normal ,i.e., $\exists x_0 \in \mathbb{R}; \mu(x_0) = 1$
- ii)  $\mu(\lambda x + (1 - \lambda)y) \geq \min\{\mu(x), \mu(y)\}, \forall x, y \in \mathbb{R}, \forall \lambda \in [0, 1]$  ( $\mu$  is called a convex fuzzy subset).
- iii)  $\mu$  is upper semicontinuous on  $\mathbb{R}$ , i.e.  $\forall x_0 \in \mathbb{R}$  and  $\forall \epsilon > 0, \exists$  neighborhood  $V(x_0) : \mu(x) \leq \mu(x_0) + \epsilon, \forall x \in V(x_0)$
- iv) The set  $\text{supp}(\mu)$  is compact in  $\mathbb{R}$ . where  $(\text{supp}(\mu) = \{x \in \mathbb{R} : \mu(x) > 0\})$

We call  $\mu$  a fuzzy real number.Denote the set of all  $\mu$  with  $\mathbb{R}_F$ .E.g.,  $\chi_{\{x_0\}} \in \mathbb{R}_F$ , for any  $x_0 \in \mathbb{R}$ , where  $\chi_{\{x_0\}}$  is the characteristic function at  $x_0$ .

For  $0 < r \leq 1$  and  $\mu \in \mathbb{R}_F$  define

$$[\mu]^r = \{x \in \mathbb{R} : \mu(x) \geq r\}$$

and

$$[\mu]^0 = \overline{\{x \in \mathbb{R} : \mu(x) \geq r\}}$$

Then it is well known that for each  $r \in [0, 1]$ ,  $[\mu]^r$  is a closed and bounded interval of  $\mathbb{R}$ .For  $u, v \in \mathbb{R}_F$  and  $\lambda \in \mathbb{R}$ ,we define uniquely the sum  $u \oplus v$  and the product  $\lambda \odot u$  by

$$[u \oplus v]^r = [u]^r + [v]^r, [\lambda \odot u]^r = \lambda [u]^r, \forall r \in [0, 1]$$

Notice  $1 \odot u = u$  and its holds

$$u \oplus v = v \oplus u, \lambda \odot u = u \odot \lambda$$

If  $0 \leq r_1 \leq r_2 \leq 1$  then  $[\mu]^{r_2} \subseteq [\mu]^{r_1}$ . Actually  $[u]^r = [u_-^r, u_+^r]$ ,  $u_-^r \leq u_+^r$ ,  $u_-^r, u_+^r \in \mathbb{R}, \forall r \in [0, 1]$

For  $\lambda > 0$  one has  $\lambda \pm = (\lambda \odot u)_{\pm}^r$ , respectively.

**Definition 2.2.** [5]  $D : \mathbb{R}_F \times \mathbb{R}_F \rightarrow \mathbb{R}_+ \cup \{0\}$

$$D(u, v) = \sup_{r \in [0, 1]} \max\{|u_-^r - v_-^r|, |u_+^r - v_+^r|\}$$

where  $[u]^r = [u_-^r, u_+^r]$ ;  $u, v \in \mathbb{R}_F$ . We have that  $D$  is a metric on  $\mathbb{R}_F$ . Then  $(\mathbb{R}_F, D)$  is a complete metric space with the following properties:

- i)  $D(u \oplus w, v \oplus w) = D(u, v) \quad \forall u, v, w \in \mathbb{R}_F$
- ii)  $D(\lambda \odot u, \lambda \odot v) = |\lambda| D(u, v) \quad \forall \lambda \in \mathbb{R}, \forall u, v \in \mathbb{R}_F$
- iii)  $D(u \oplus v, w \oplus e) \leq D(u \oplus w) + D(v \oplus e), \forall u, v, w, e \in \mathbb{R}_F$

Here  $\sum^*$  is stands for fuzzy summation and  $\tilde{0} : \chi_{\{0\}} \in \mathbb{R}_F$  is the neutral element with respect to  $\oplus$ , i.e.,

$$u \oplus \tilde{0} = \tilde{0} \oplus u = u, \forall u \in \mathbb{R}_F$$

Denote

$$D^*(f, g) = \sup_{x \in [a, b]} D(f, g)_x$$

Where  $f, g : [a, b] \rightarrow \mathbb{R}_F$ .

We define  $C_F^U([a, b])$  the space of uniformly continuous functions from  $[a, b] \rightarrow \mathbb{R}_F$ , also  $C_F([a, b])$  the space of fuzzy continuous functions on  $[a, b]$ . It is clear that

$$C_F^U([a, b]) = C_F([a, b])$$

and  $L_F([a, b])$  is the space of Lebesgue integrable functions.

**Definition 2.3.** [13] Let  $u, v \in \mathbb{R}_F$ . If there exists  $w \in \mathbb{R}_F$  such that  $u = v + w$ , the  $w$  is called the Hukuhara difference of  $u$  and  $v$ , and it is denoted by  $u \ominus v$ .

**Definition 2.4.** [13] Let  $u, v \in \mathbb{R}_F$ . If there exists  $w \in \mathbb{R}_F$  such that

$$u \ominus_{gH} v = w \iff \begin{cases} (i) & u = v + w \\ \text{or} & (ii) & u = v + (-1)w \end{cases}$$

Then  $w$  is called the generalized Hukuhara difference of  $u$  and  $v$ .

Please note that a function  $f : [a, b] \rightarrow \mathbb{R}_F$  so called fuzzy-valued function. The  $r$ -level representation of fuzzy-valued function  $f$  is expressed by

$$f_r(t) = [f_r^-(t), f_r^+(t)], t \in [a, b], r \in [0, 1]$$

Here,  $f^r(t) = f_r(t)$

**Definition 2.5.** [5] Let  $f : [a, b] \rightarrow \mathbb{R}_F$ . We say that  $f$  is Fuzzy-Riemann integrable to  $I \in \mathbb{R}_F$  if for any  $\epsilon > 0$ , there exists  $\delta > 0$  such that for any division  $P = \{[u, v]; \xi\}$

of  $[a, b]$  with the norms  $\Delta(P) < \delta$ , we have

$$D\left(\sum_P^* (v - u) \odot f(\xi), I\right) < \epsilon$$

We write

$$I = (FR) \int_a^b f(x) dx$$

**Theorem 2.1.** [9] *Let  $f : [a, b] \rightarrow \mathbb{R}_F$  be fuzzy continuous. Then  $(FR) \int_a^b f(x) dx$  exists and belongs to  $\mathbb{R}_F$ , furthermore it holds*

$$\left[ (FR) \int_a^b f(x) dx \right]^r = \left[ \int_a^b f_-^{(r)}(x) dx, \int_a^b f_+^{(r)}(x) dx \right], \quad r \in [0, 1]$$

**Theorem 2.2.** [5] *Let  $f \in C_F([a, b])$  and  $c \in [a, b]$ . Then*

$$(FR) \int_a^b f(x) dx = (FR) \int_a^c f(x) dx + (FR) \int_c^b f(x) dx$$

**Theorem 2.3.** [5] *Let  $f, g \in C_F([a, b])$  and  $c_1, c_2 \in \mathbb{R}$ . Then*

$$(FR) \int_a^b (c_1 f(x) + c_2 g(x)) dx = c_1 (FR) \int_a^b f(x) dx + c_2 (FR) \int_a^b g(x) dx$$

also we need

**Lemma 2.1.** [5] *If  $f, g : [a, b] \subseteq \mathbb{R} \rightarrow \mathbb{R}_F$  are fuzzy continuous functions, then the function  $F : [a, b] \rightarrow \mathbb{R}_+$  defined by  $F(x) = D(f(x), g(x))$  is continuous on  $[a, b]$ :*

$$D \left( (FR) \int_a^b f(x) dx, (FR) \int_a^b g(x) dx \right) \leq (FR) \int_a^b D(f(x), g(x)) dx$$

**Definition 2.6.** [4] *Let  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $0 < v \leq 1$ .*

The fuzzy Riemann-Liouville integral of fuzzy-valued function  $f$  is defined as following:

$$(I_{\alpha+}^v f)(x) = \frac{1}{\Gamma(v)} \odot \int_a^x (x-t)^{v-1} \odot f(t) dt, \quad x \in [a, b]$$

$$I_{\alpha+}^0 f(x) = f$$

Let us consider the  $r$ -level representation of fuzzy-valued function  $f$  as  $f_r(t) = [f_r^-(t), f_r^+(t)]$ ,  $t \in [a, b]$ ,  $r \in [0, 1]$

Also, we define the fuzzy fractional right Riemann-Liouville operator by

$$I_{b-}^v f(x) = \frac{1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot f(t) dt, \quad x \in [a, b]$$

$$I_{b-}^0 f(x) = f$$

Above,  $\Gamma$  denotes the gamma function:

$$\Gamma(v) = \int_0^{\infty} e^{-t} t^{v-1} dt$$



**Definition 2.7.** [4] Let  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $x_0$  in  $(a, b)$  and  $\Phi(x) = \frac{1}{\Gamma(1-v)} \int_a^x \frac{f(t)}{(x-t)^v} dt$ . We say that  $f$  is Riemann-Liouville H-differentiable about order  $0 < v < 1$  at  $x_0$ , if there exists an element  $({}^{RL}D_{\alpha+}^v)(x_0) \in \mathbb{R}_F$ , such that for  $h > 0$  sufficiently small

$$\begin{aligned} i) \quad & ({}^{RL}D_{\alpha+}^v)(x_0) = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0+h) \ominus \Phi(x_0)}{h} = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0) \ominus \Phi(x_0-h)}{h} \\ \text{or} \\ ii) \quad & ({}^{RL}D_{\alpha+}^v)(x_0) = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0) \ominus \Phi(x_0+h)}{-h} = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0-h) \ominus \Phi(x_0)}{-h} \\ \text{or} \\ iii) \quad & ({}^{RL}D_{\alpha+}^v)(x_0) = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0+h) \ominus \Phi(x_0)}{h} = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0-h) \ominus \Phi(x_0)}{-h} \\ \text{or} \\ iv) \quad & ({}^{RL}D_{\alpha+}^v)(x_0) = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0) \ominus \Phi(x_0+h)}{-h} = \lim_{h \rightarrow 0^+} \frac{\Phi(x_0) \ominus \Phi(x_0-h)}{h} \end{aligned}$$

### 3. MAIN RESULTS

**Definition 3.1.** [14] Let  $f \in C_F([a, b]) \cap L_F([a, b])$  be a fuzzy set-valued function. Then  $f$  is said to be Caputo's H-differentiable at  $x$  when

$$({}^{D_{\alpha+}^v} f)(x) = \frac{1}{\Gamma(1-v)} \int_a^x \frac{f'(t)}{(x-t)^v} dt$$

where  $0 < \alpha < 1$  and  $0 < v < 1$ .

Also, we adopt the same procedure to present Caputo's H-differentiability, we say  $f$  is

[(i) - v]-differentiable if Eq. (8) holds while  $f$  is (i)-differentiable, and  $f$  is [(ii) - v]-differentiable if Eq. (21) holds while  $f$  is (ii)-differentiable.

**Definition 3.2.** [15] Let  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $f^n$  is integrable. Then the right fuzzy Caputo derivate of  $f$  for  $n-1 < v < n$ , and  $x \in [a, b]$ ,  $D_{b-}^v f(x) \in \mathbb{R}_F$  and defined by

$$D_{b-}^v f(x) = \frac{(-1)^n}{\Gamma(n-v)} \odot \int_x^b (t-x)^{-v+n-1} \odot f^n(t) dt$$

and for  $n = 1$

$$D_{b-}^v f(x) = \frac{(-1)}{\Gamma(1-v)} \odot \int_x^b (t-x)^{-v} \odot f'(t) dt$$

**Theorem 3.1.** [14] Let  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $0 < v < 1$ ,  $0 \leq r \leq 1$ ,

i) Let  $f$  be (ii)-differentiable, then we have [(i) - v] differentiable right fuzzy Caputo derivative and

$$({}^{D_{b-}^v} f)(x, r) = [ ({}^{D_{b-}^v} f_-)(x, r), ({}^{D_{b-}^v} f_+)(x, r) ]$$

ii) Let  $f$  be  $(i)$ -differentiable, then we have  $[(i) - v]$  differentiable right fuzzy Caputo derivative and

$$(D_{b-}^v f)(x, r) = [ (D_{b-}^v f_+)(x, r), (D_{b-}^v f_-)(x, r) ]$$

**Theorem 3.2.** [14] Let  $0 < v < 1$ ,  $D_{b-}^v f(x) = g(x, f(x))$  with the fuzzy initial condition  $f_0 = f(b)$ , the fuzzy fractional differential equation is equivalent to one of the following integral equations:

i) if  $f$  is a  $[(i) - v]$  differentiable fuzzy-valued function, then

$$f(x) = f(b) \oplus \frac{1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt$$

ii) if  $f$  is a  $[(ii) - v]$  differentiable fuzzy-valued function, then

$$f(x) = f(b) \oplus \frac{-1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt$$

**Theorem 3.3.** Let  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $0 < v < 1$ ,  $p, q > 0$  such that  $\frac{1}{p} + \frac{1}{q} = 1$ , and  $(D_{b-}^v f)(x) \in \mathbb{R}_F$ ;  $(t \in [a, b])$

$$D \left( \frac{1}{b-a} \odot (FR) \int_a^b f(x) dx, f(b) \right) \leq \frac{\sup D((D_{b-}^v f)(t), \tilde{0})}{\Gamma(v)(p(v-1) + 1)^{\frac{1}{p}}(v + \frac{1}{p})} (b-a)^{v-1 + \frac{1}{p}}$$

*Proof.* We have

$$\begin{aligned} D \left( \frac{1}{b-a} \odot (FR) \int_a^b f(x) dx, f(b) \right) &= D \left( \frac{1}{b-a} \odot (FR) \int_a^b f(x) dx, \frac{f(b)}{b-a} \int_a^b dx \right) \\ &= D \left( \frac{1}{b-a} \odot (FR) \int_a^b f(x) dx, \frac{1}{b-a} \odot (FR) \int_a^b f(b) dx \right) \\ &= \frac{1}{b-a} D \left( (FR) \int_a^b f(x) dx, (FR) \int_a^b f(b) dx \right) \\ &\leq \frac{1}{b-a} \int_a^b D(f(x), f(b)) dx \quad (*) \end{aligned}$$

Here  $[(i) - v]$  differentiable .

We notice that  $f \in C_F([a, b]) \cap L_F([a, b])$ ,  $0 < v < 1$ ,

$$f(x) = f(b) \oplus \frac{1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt$$

For  $a \leq x \leq b$ , we have

$$\begin{aligned}
 D(f(x), f(b)) &= D\left(f(b) \oplus \frac{1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt, f(b)\right) \\
 &= D\left(\frac{1}{\Gamma(v)} \odot \int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt, \tilde{0}\right) \\
 &\leq \frac{1}{\Gamma(v)} D\left(\int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt, \tilde{0}\right) \\
 &\leq \frac{1}{\Gamma(v)} D\left(\int_x^b (t-x)^{v-1} \odot (D_{b-}^v f)(t) dt, \int_x^b \tilde{0} dt\right) \\
 &\leq \frac{1}{\Gamma(v)} \int_x^b (t-x)^{v-1} \left(D\left((D_{b-}^v f)(t), \tilde{0}\right)\right) dt \\
 &\leq \frac{1}{\Gamma(v)} \left(\int_x^b (t-x)^{p(v-1)}\right)^{\frac{1}{p}} \left(\int_x^b \left(D\left((D_{b-}^v f)(t), \tilde{0}\right)\right)^q dt\right)^{\frac{1}{q}} \\
 &\leq \frac{1}{\Gamma(v)} \frac{(b-x)^{v-1+\frac{1}{p}}}{(p(v-1)+1)^{\frac{1}{p}}} \left(\int_x^b \left(D\left((D_{b-}^v f)(t), \tilde{0}\right)\right)^q dt\right)^{\frac{1}{q}}
 \end{aligned}$$

Now,  $\forall x \in [a, b]$  and for (\*)

$$\begin{aligned}
 D\left(\frac{1}{b-a} \odot (FR) \int_a^b f(x) dx, f(b)\right) &\leq \frac{1}{b-a} \int_a^b D(f(x), f(b)) dx \\
 &\leq \frac{\sup\left(D\left((D_{b-}^v f)(t), \tilde{0}\right)\right)}{(b-a)\Gamma(v)(p(v-1)+1)^{\frac{1}{p}}} \left(\int_a^b (b-x)^{v-1+\frac{1}{p}} dx\right) \\
 &= \frac{\sup\left(D\left((D_{b-}^v f)(t), \tilde{0}\right)\right)}{\Gamma(v)(p(v-1)+1)^{\frac{1}{p}}} \frac{(b-a)^{v-1+\frac{1}{p}}}{v+\frac{1}{p}} \quad (t \in [a, b])
 \end{aligned}$$

□

## REFERENCES

- [1] L.A. Zadeh, Fuzzy Sets, Information and Control (1965) 8, 338-35.
- [2] Milovanović, G.V., Pecarić, J.E.: On generalization of the inequality of A. Ostrowski and some related applications. (1976) 44, 155-158,
- [3] Anastassiou, G.A., On right fractional calculus. Chaos Solitons Fractals (2009). 42, 365-376
- [4] Salahshour, S. Allahviranloo, T. Abbasbandy, Solving fuzzy fractional differential equations by fuzzy Laplace transforms, Commun Nonlinear sci numer simulat (2012)17-1372-1381
- [5] Anastassiou, G.A., Fuzzy Fractional Calculus and the Ostrowski Integral Inequality. Intelligent Mathematics: Computational Analysis (2011), 553-574. Springer, Berlin
- [6] Anastassiou, G.A., Ostrowski type inequalities. Proc. AMS (1995), 123, 3775-3791.
- [7] Kilbas A. A., Srivastava H. M., Trujillo J. J., Theory and Applications of Fractional Differential Equations, in: North-Holland Mathematics Studies (2006), vol. 204, Elsevier Science B.V., Amsterdam.
- [8] Samko S. G., Kilbas A. A., Marichev O. I., Fractional Integrals and Derivatives Theory and Application, Gordon and Breach Science, New York, (1993)
- [9] Ostrowski, A, Über die absolutabweichung einer differentiebaren funktion von ihrem intetegralmittelwert. Comment. Math. Helv.(1938). 10, 226-227.
- [10] Pachpatte, B.G., New Ostrowski and Gruss type inequalities. An. Ştiinţ. Univ. "Al. I. Cuza" Iaşi, Mat.. LI (2005),2, 377-386.
- [11] Bohner, M, Matthews, T: Ostrowski inequalities on time scales. J. Inequal. Pure Appl. Math (2008),9, 6-8

- [12] Anastassiou, G.A., Univariate right fractional Ostrowski inequalities. CUBO A. Mathematical Journal (2012), Vol , 14.1.1-7.
- [13] Armand A., Mohammadi S., Existence and uniqueness for fractional differential equations with uncertainty (2014) ID jums-00011, 9 Pages doi: 10.5899/2014/jums-00011
- [14] Salahshour, S. Allahviranloo, T. Abbasbandy, S. Baleanu, D. Existence and uniqueness results for fractional differential equations with uncertainty ( 2012), Adv. Diff. Equ. 112 ,
- [15] Allahviranloo, T. Abbasbandy, S. Salahshour, S. Shahryari, M.R. Baleanu, D. On Solutions of Linear Fractional Differential Equations with the Uncertainty, Hindawi Publishing Corporation abstract and Applied Analysis (2013), ID 178378, 13 pages

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, SÜTCÜ İMAM UNIVERSITY,  
KAHRAMANMARAŞ, TURKEY

*E-mail address:* `citil@ksu.edu.tr`

*E-mail address:* `m.gezercan@hotmail.com`

## NEW INTUITIONISTIC FUZZY LEVEL SETS

GÖKHAN ÇUVALCIOĞLU AND YELDA YORULMAZ

ABSTRACT. The concept of Intuitionistic Fuzzy Sheet  $t$ -Cut Set and Intuitionistic Fuzzy  $\alpha - t$  Block Cut Set are introduced. The differences between  $C_{\alpha,\beta}$  level set and new intuitionistic fuzzy sets is shown.

Received: 27-June-2016

Accepted: 29-August-2016

### 1. INTRODUCTION

The function  $\mu : X \rightarrow [0, 1]$  is called a fuzzy set over  $X(FS(X))[?]$ . For  $x \in X$ ,  $\mu(x)$  is the membership degree of  $x$  and the non-membership degree is  $1 - \mu(x)$ . Intuitionistic fuzzy sets have been introduced by Atanassov [2], as an extension of fuzzy sets. If  $X$  is a universal then a intuitionistic fuzzy set  $A$ , the membership and non-membership degree for each  $x \in X$  respectively,  $\mu_A(x)(\mu_A : X \rightarrow [0, 1])$  and  $\nu_A(x)(\nu_A : X \rightarrow [0, 1])$  such that  $0 \leq \mu_A(x) + \nu_A(x) \leq 1$ . The class of intuitionistic fuzzy sets on  $X$  is denoted by  $IFS(X)$ .

**Definition 1.1.** [2] An intuitionistic fuzzy set (shortly IFS) on a set  $X$  is an object of the form

$$A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$$

where  $\mu_A(x), (\mu_A : X \rightarrow [0, 1])$  is called the “degree of membership of  $x$  in  $A$ ”,  $\nu_A(x), (\nu_A : X \rightarrow [0, 1])$  is called the “degree of non-membership of  $x$  in  $A$ ”, and where  $\mu_A$  and  $\nu_A$  satisfy the following condition:

$$\mu_A(x) + \nu_A(x) \leq 1, \text{ for all } x \in X.$$

**Definition 1.2.** [1] An intuitionistic fuzzy set  $A$  is said to be contained in an intuitionistic fuzzy set  $B$  if and only if, for all  $x \in X : \mu_A(x) \leq \mu_B(x)$  and  $\nu_A(x) \geq \nu_B(x)$ . If fuzzy set  $B$  contains fuzzy set  $A$  then it is shown by  $A \sqsubseteq B$ .

It is clear that  $A = B$  if and only if  $A \sqsubseteq B$  and  $B \sqsubseteq A$ .

**Definition 1.3.** [2] Let  $A \in IFS(X)$  and let  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$  then the set

$$A^c = \{ \langle x, \nu_A(x), \mu_A(x) \rangle : x \in X \}$$

is called the complement of  $A$ .

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* Primary 05C38, 15A15; Secondary 05A15, 15A18.  
*Key words and phrases.* Intuitionistic Fuzzy Sets, Intuitionistic Fuzzy Sheet  $t$ -Cut Sets, Intuitionistic Fuzzy  $\alpha - t$  Block Cut Sets.




FIGURE 1

The intersection and the union of two IFSs  $A$  and  $B$  on  $X$  are defined by

$$A \sqcap B = \{ \langle x, \mu_A(x) \wedge \mu_B(x), \nu_A(x) \vee \nu_B(x) \rangle : x \in X \}$$

$$A \sqcup B = \{ \langle x, \mu_A(x) \vee \mu_B(x), \nu_A(x) \wedge \nu_B(x) \rangle : x \in X \}$$

Some special Intuitionistic Fuzzy Sets on  $X$  are defined as following;

$$O^* = \{ \langle x, 0, 1 \rangle : x \in X \}$$

$$U^* = \{ \langle x, 0, 0 \rangle : x \in X \}$$

**Definition 1.4.** [4] Let  $A \in IFS(X)$ . Then  $(\alpha, \beta)$ -cut of  $A$  is a crisp subset  $C_{\alpha, \beta}(A)$  of the IFS  $A$  is given by

$$C_{\alpha, \beta}(A) = \{ x : x \in X \text{ such that } \mu_A(x) \geq \alpha, \nu_A(x) \leq \beta \}$$

where  $\alpha, \beta \in [0, 1]$  with  $\alpha + \beta \leq 1$ .

## 2. SHEET AND BLOCK CUT INTUITIONISTIC FUZZY LEVEL SETS

**Definition 2.1.** Let  $X$  be a set and  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \} \in IFS(X)$ . If  $t \in [0, 1]$  then sheet  $t$ -cut of  $A$  defined as following

$$A(t) = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : \mu_A(x) + \nu_A(x) = t, x \in X \}$$

**Proposition 2.1.** Let  $X$  be a set and  $A, B \in IFS(X)$ . For every  $t \in [0, 1]$ ,

- (1)  $(A \sqcup B)(t) = A(t) \sqcup B(t)$
- (2)  $A(t) \sqcap B(t) = (A \sqcap B)(t)$
- (3)  $(A^c(t))^c = A(t)$

*Proof.* (1)

$$A(t) \sqcup B(t) = \left\{ \begin{array}{l} \langle x, \max(\mu_A(x), \mu_B(x)), \min(\nu_A(x), \nu_B(x)) \rangle : \\ \mu_A(x) + \nu_A(x) = t \wedge \mu_B(x) + \nu_B(x) = t, x \in X \end{array} \right\}$$

If  $\mu_A(x) \geq \mu_B(x)$  then from  $\mu_A(x) + \nu_A(x) = t = \mu_B(x) + \nu_B(x)$  we obtain  $\nu_A(x) \leq \nu_B(x)$ .

$$\max(\mu_A(x), \mu_B(x)) + \min(\nu_A(x), \nu_B(x)) = \mu_A(x) + \nu_A(x) = t$$

If  $\mu_A(x) \leq \mu_B(x)$  then from  $\mu_A(x) + \nu_A(x) = t = \mu_B(x) + \nu_B(x)$  we obtain  $\nu_A(x) \leq \nu_B(x)$ .

$$\max(\mu_A(x), \mu_B(x)) + \min(\nu_A(x), \nu_B(x)) = \mu_B(x) + \nu_B(x) = t$$

Thence,  $(A \sqcup B)(t) = A(t) \sqcup B(t)$ .

(2)

$$A(t) \sqcap B(t) = \left\{ \begin{array}{l} \langle x, \min(\mu_A(x), \mu_B(x)), \max(\nu_A(x), \nu_B(x)) \rangle : \\ \mu_A(x) + \nu_A(x) = t \wedge \mu_B(x) + \nu_B(x) = t, x \in X \end{array} \right\}$$

If  $\mu_A(x) \geq \mu_B(x)$  then from  $\mu_A(x) + \nu_A(x) = t = \mu_B(x) + \nu_B(x)$  we obtain  $\nu_A(x) \leq \nu_B(x)$ .

$$\min(\mu_A(x), \mu_B(x)) + \max(\nu_A(x), \nu_B(x)) = \mu_B(x) + \nu_B(x) = t$$

If  $\mu_A(x) \leq \mu_B(x)$  then from  $\mu_A(x) + \nu_A(x) = t = \mu_B(x) + \nu_B(x)$  we obtain  $\nu_A(x) \leq \nu_B(x)$ .

$$\min(\mu_A(x), \mu_B(x)) + \max(\nu_A(x), \nu_B(x)) = \mu_A(x) + \nu_A(x) = t$$

Therefore, we obtain that  $A(t) \sqcap B(t) = (A \sqcap B)(t)$ .

(3) It is clear. □

*Remark 2.1.* Let  $X$  be a set and  $A \in IFS(X)$ .  $A(t)$  is a fuzzy set on  $[0, t]$ .

**Proposition 2.2.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $t, s \in [0, 1]$  then

$$\text{Either } A(t) \sqcap A(s) = O^* \text{ or } t = s$$

*Proof.* If  $A(t) \sqcap A(s) \neq O^*$  and  $t \neq s$  then there exists  $x \in X$ ,

$$\begin{aligned} \mu_A(x) + \nu_A(x) &= t \text{ and } \mu_A(x) + \nu_A(x) = s \\ &\Rightarrow t = s \end{aligned}$$

□

**Corollary 2.1.** There exist an equivalence relation on  $X$  such that the sheet  $t$ -cuts are equivalence class of that relation.

**Definition 2.2.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $t \in [0, 1]$  and  $\alpha \in [0, t]$  then

$$A(t)_\alpha = \{x : x \in X, A(t)(x) \geq (\alpha, t - \alpha)\}$$

is called  $\alpha - t$  block cut of  $A$ .

From definitions, it is easily seen that for every  $t \in [0, 1]$ ,  $A(t) \in FS(X)$ . Because  $A(t) : X \rightarrow [0, t]$  and  $[0, t] \sim [0, 1]$ . For short notation, if  $A(t) : X \rightarrow [0, t]$  then  $A(t)$  will be called  $t$ -fuzzy set on  $X$  ( $A(t) \in FS_t(X)$ ). It is clear that  $A(t)_\alpha$  is a crisp set.

**Proposition 2.3.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $t \in [0, 1]$  then

- (1)  $A(t)_t = \{x : x \in X, \mu_A(x) = t \wedge \nu_A(x) = 0\}$
- (2)  $A(t)_0 = \{x : x \in X, \mu_A(x) \geq 0 \wedge \nu_A(x) \leq t\}$




FIGURE 2

$$(3) A(0) = U^*$$

**Example 2.1.** Let  $X = \{a, b, c, d, e\}$  and  $A = \{(a, 0.5, 0.4), (b, 0.2, 0.3), (c, 0.5, 0.3), (d, 0.4, 0.4), (e, 0.4, 0.1)\}$ .

- (1)  $A(0.5)_{0.3} = \{e\}$  but  $C_{0.3,0.5}(A) = \{a, c, d, e\}$  and  $C_{0.5,0.3}(A) = \{c\}$ .
- (2)  $A(0.8)_{0.5} = \{c\}$  but  $0.8 + 0.5 > 1$  so, we can not obtain  $C_{0.5,0.8}(A)$  or  $C_{0.8,0.5}(A)$ .

**Example 2.2.** Let  $X = \{a, b, c, d, e\}$  and  $A = \{(a, 0.1, 0.2), (b, 0.4, 0.3), (c, 0.6, 0.2), (d, 0.7, 0.1), (e, 0.2, 0.5)\}$ .

$$A(0.3)_{0.2} = \emptyset \text{ but } C_{0.2,0.3}(A) = \{b, c, d\} \text{ and } C_{0.3,0.2}(A) = \{c, d\}.$$

That is seen from the examples,  $(\alpha, \beta)$ -cut of an intuitionistic fuzzy set  $A$  and  $\alpha - t$  block cut of  $A$  are different sets. For all  $t \in [0, 1]$  and  $\alpha \in [0, t]$ , we can determine  $\alpha - t$  block cut of  $A$ , if  $\alpha + t > 1$  then we can not determine  $(\alpha, \beta)$ -cut of  $A$ . Consequently,  $\alpha - t$  block cut of an intuitionistic fuzzy set allows a more extensive studying area.

**Proposition 2.4.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $t \in [0, 1]$  and  $\alpha, \beta \in [0, t]$  such that  $\alpha \leq \beta$  then  $A(t)_\beta \subseteq A(t)_\alpha$ .

*Proof.* Let  $\alpha \leq \beta$ . If  $x \in A(t)_\beta$  then

$$A(t)(x) \geq (\beta, t - \beta) \geq (\alpha, t - \alpha)$$

Therefore  $x \in A(t)_\alpha$ .  $\square$

**Proposition 2.5.** Let  $X$  be a set and  $A, B \in IFS(X)$ . If  $t \in [0, 1]$  and  $\alpha \in [0, t]$  then

- (1)  $A(t)_\alpha \cup B(t)_\alpha = (A(t) \sqcup B(t))_\alpha$
- (2)  $A(t)_\alpha \cap B(t)_\alpha = (A(t) \sqcap B(t))_\alpha$



- (3)  $(A(t)^c)_\alpha = t - A(t)_\alpha, (t(x) = t)$   
 (4)  $(A^c(t)^c)_\alpha = t - A(t)_\alpha = t - A^c(t)_\alpha$

*Proof.* (1)

$$\begin{aligned}
 x \in A(t)_\alpha \cup B(t)_\alpha &\Leftrightarrow A(t)(x) \geq (\alpha, t - \alpha) \vee B(t)(x) \geq (\alpha, t - \alpha) \\
 &\Leftrightarrow (\mu_{A(t)}(x) \geq \alpha \wedge \nu_{A(t)}(x) \leq t - \alpha) \vee (\mu_{B(t)}(x) \geq \alpha \wedge \nu_{B(t)}(x) \leq t - \alpha) \\
 &\Leftrightarrow (\mu_{A(t)}(x) \geq \alpha \vee \mu_{B(t)}(x) \geq \alpha) \wedge (\nu_{A(t)}(x) \leq t - \alpha \vee \nu_{B(t)}(x) \leq t - \alpha) \\
 &\Leftrightarrow (\mu_{A(t)}(x) \vee \mu_{B(t)}(x)) \geq \alpha \wedge (\nu_{A(t)}(x) \wedge \nu_{B(t)}(x)) \leq t - \alpha \\
 &\Leftrightarrow \mu_{A(t) \sqcup B(t)}(x) \geq \alpha \wedge \nu_{A(t) \sqcup B(t)}(x) \leq t - \alpha \\
 &\Leftrightarrow x \in (A(t) \sqcup B(t))_\alpha
 \end{aligned}$$

(2)

$$\begin{aligned}
 A(t)_\alpha \cap B(t)_\alpha &= \{x \in X : A(t)(x) \geq (\alpha, t - \alpha) \wedge B(t)(x) \geq (\alpha, t - \alpha)\} \\
 &= \{x \in X : (\mu_{A(t)}(x) \geq \alpha \wedge \nu_{A(t)}(x) \leq t - \alpha) \wedge (\mu_{B(t)}(x) \geq \alpha \wedge \nu_{B(t)}(x) \leq t - \alpha)\} \\
 &= \{x \in X : (\mu_{A(t)}(x) \geq \alpha \wedge \mu_{B(t)}(x) \geq \alpha) \wedge (\nu_{A(t)}(x) \leq t - \alpha \wedge \nu_{B(t)}(x) \leq t - \alpha)\} \\
 &= \{x \in X : (\mu_{A(t)}(x) \wedge \mu_{B(t)}(x)) \geq \alpha \wedge (\nu_{A(t)}(x) \vee \nu_{B(t)}(x)) \leq t - \alpha\} \\
 &= \{x \in X : \mu_{A(t) \sqcap B(t)}(x) \geq \alpha \wedge \nu_{A(t) \sqcap B(t)}(x) \leq t - \alpha\} \\
 &= (A(t) \sqcap B(t))_\alpha
 \end{aligned}$$

(3)

$$\begin{aligned}
 (A(t)^c)_\alpha &= \{x \in X : A(t)^c(x) \geq (\alpha, t - \alpha)\} \\
 &= \{x \in X : \nu_{A(t)}(x) \geq \alpha \wedge \mu_{A(t)}(x) \leq t - \alpha\} \\
 &= \{x \in X : t - \nu_{A(t)}(x) \leq t - \alpha \wedge t - \mu_{A(t)}(x) \geq \alpha\} \\
 &= \{x \in X : (t - A(t))(x) \geq (\alpha, t - \alpha)\} \\
 &= t - A(t)_\alpha
 \end{aligned}$$

□

## REFERENCES

- [1] Atanassov K.T., Intuitionistic Fuzzy Sets, VII ITKR'sSession, Sofia, June (1983).
- [2] Atanassov K.T., Intuitionistic Fuzzy Sets, Fuzzy Sets and Systems, (1986), 20, 87-96
- [3] Çuvalcioglu G., Yılmaz S., On Block and Sheet Level of Intuitionistic Fuzzy Sets, 2nd Int. IFS Conference, Mersin, October(2015).
- [4] Sharma P.K.,  $(\alpha, \beta)$ -Cut of Intuitionistic Fuzzy Groups, International Mathematical forum, (2011),53(6), 2605-2614.
- [5] Zadeh L.A., Fuzzy Sets, Information and Control, (1965), 8, 338-353.

UNIVERSITY OF MERSIN DEPARTMENT OF MATHEMATICS MERSIN, TURKEY.

*E-mail address:* gcuvalcioglu@mersin.edu.tr

*E-mail address:* yeldayorulmaz@gmail.com

## ENERGY SPECTRUM OF SPINLESS PARTICLES IN ELECTROMAGNETIC FIELDS

ÖZGÜR MIZRAK AND KENAN SOGUT

ABSTRACT. Dynamics of the non-relativistic and relativistic charged spinless particles subjected to space-dependent parallel and orthogonal electromagnetic fields is investigated by solving Schrödinger and Klein-Gordon equations. Exact solutions of the motion are used to obtain the quantized energy spectrum and momentum of the particles. Some numerical results for the first few quantum levels are determined with the help of MATHEMATICA software.

Received: 22–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Finding the exact solutions of the wave equations for the external fields is one of the old problems. Among these equations Schrödinger and Klein-Gordon are the most studied ones. Besides by the increase in the applications of the electric and magnetic fields in fundamental areas of technology, especially in electromechanics, health physics and so forth, a significant interest has been given to these solutions. Such efforts have been performed for different configurations of the external fields [1-3].

These studies provide remarkable information regarding the quantum mechanical systems. Some of these attempts are the interpretation of the physical processes. The most important ones are Compton scattering by a laser source, Brownian motion, coherent states, and energy levels of electrons.

There are very few studies in the literature on the solution of the wave equation of the spinless particles in the presence of both electric and magnetic fields. The aim of this study is to move this attempt one step further by obtaining the exact solutions of the spinless particles for two orientations of decaying electric and magnetic fields given by *Case(i)*  $A_0 = \frac{E_0}{z}$ ,  $A_1 = \frac{B_0}{y}$  and *Case(ii)*  $A_0 = \frac{E_0}{y}$ ,  $A_1 = \frac{B_0}{z}$ , where  $E_0$  and  $B_0$  are constants. The first and second cases belong to the parallel and orthogonal fields, respectively. We note that  $y$  and  $z$  variables are defined in the region  $(0, \infty)$  to keep the finite external fields. Such kind of varying electromagnetic field is encountered in semiconductor heterostructures.

In the following sections, the exact solutions for nonrelativistic and relativistic cases will be obtained, respectively. By comparing the solutions of the nonrelativistic and relativistic wave equations of the spinless particles, contributions coming

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
*Key words and phrases.* Relativistic particles, electromagnetic fields, energy spectrum.

from the relativistic effects will be considered and by using the mathematical properties of the wave functions we will obtain the energy spectrum and exact solutions for both cases.

## 2. SOLUTION OF THE SCHRÖDINGER EQUATION

Motion of the nonrelativistic spinless particles is described by the Schrödinger equation and in the existence of the external electromagnetic fields, it is given by (we take  $\hbar = 1$ )

$$(2.1) \quad \left[ \frac{(\vec{P} - e\vec{A})^2}{2m} \right] \Phi = (i\frac{\partial}{\partial t} - eA_0)\Phi$$

where  $e$  is charge,  $m$  is mass of the particle,  $\vec{A}$  is the vector electromagnetic potential. In the following steps we solve the Schrödinger equation for the cases where electric and magnetic fields are parallel and perpendicular to each other.

**2.1. Case (i) Parallel EM Fields.** For the choice of  $A_0 = \frac{E_0}{z}$ ,  $A_1 = \frac{B_0}{y}$ ,  $\vec{E} \parallel \vec{B}$ . We define the solution of (2.1) by

$$(2.2) \quad \Phi_{\parallel} = e^{i(P_x - \epsilon t)} H(y)K(z)$$

Plugging this solution into (2.1) we obtain,

$$\left[ \underbrace{\left( P_x - \frac{eB_0}{y} \right)^2}_{\hat{Q}(y)} + \underbrace{P_y^2 + P_z^2 - 2m \left( \epsilon - \frac{eE_0}{z} \right)}_{\hat{D}(z)} \right] H(y)K(z) = 0$$

In short we can write

$$\left[ \hat{Q}(y) + \hat{D}(z) \right] H(y)K(z) = 0$$

Separating this equation with respect to  $y$  and  $z$ , we obtain

$$(2.3) \quad \left[ \hat{Q}(y) + b \right] H(y) = 0$$

$$(2.4) \quad \left[ \hat{D}(z) - b \right] K(z) = 0$$

where  $b$  is the constant of separation.

Let  $\gamma^2 = (P_x^2 + b)$ , and making  $\rho = 2\gamma y$  change of variable (2.3) becomes Whittaker equation [4]

$$(2.5) \quad \left[ \frac{d^2}{d\rho^2} - \frac{e^2 B_0^2}{\rho^2} + \frac{2eB_0 P_x}{\gamma \rho} - \frac{1}{4} \right] H(\rho) = 0$$

So exact solution of (2.3) is

$$(2.6) \quad H(y) = W_{\lambda, \mu}(2\gamma y)$$

where  $\mu^2 = \frac{1}{4} + e^2 B_0^2$ , and  $\lambda = \frac{eB_0 P_x}{\gamma}$ .

In order Whittaker function to be bounded [4]

$$\mu - \lambda = -\left(n + \frac{1}{2}\right) = -N, \quad n = 0, 1, 2, \dots$$

So from this equality we find

$$b = P_x^2 \left( \frac{1}{1 + \frac{(1/4+N^2)}{e^2 B_0^2}} + 2NeB_0 \sqrt{1 + \frac{1}{4e^2 B_0^2}} - 1 \right)$$

Now for the solution of (2.4), this equation is written as

$$(2.7) \quad \begin{aligned} & \left[ P_z^2 - 2m \left( \epsilon - \frac{eE_0}{z} \right) - b \right] K(z) = 0 \\ & \left[ \frac{d^2}{dz^2} - \frac{2meE_0}{z} + (2m\epsilon + b) \right] K(z) = 0 \end{aligned}$$

(2.7) is similar to the below equation

**Definition 2.1.**

$$xy'' + (ax + b)y' + (cx + d)y = 0$$

For  $a^2 > 4c$  solution is given by [5]

$$y = x^{-\frac{b}{a}} e^{-\frac{cx}{a}} {}_1F_1 \left( \frac{2d - ab}{2\sqrt{a^2 - 4c}}, \frac{b - 1}{2}, x\sqrt{a^2 - 4c} \right)$$

Returning to the equation (2.7),

$$zK''(z) + [(2m\epsilon + b)z - 2meE_0] K(z) = 0$$

for  $0 > 2m\epsilon + b$

$$K(z) = {}_1F_1 \left( \frac{-meE_0}{\sqrt{-4(2m\epsilon + b)}}, -\frac{1}{2}, z\sqrt{-4(2m\epsilon + b)} \right)$$

From the requirement of Hypergeometric functions to be finite

$$\frac{-2meE_0}{\sqrt{-4(2m\epsilon + b)}} = -n$$

where  $n = 0, 1, 2, \dots$  we obtain the energy spectrum of Schrödinger equation for the parallel case as

$$\epsilon_{\parallel} = \frac{P_x^2}{2m} \left[ 1 - \frac{1}{1 + \frac{(1/4+N^2)}{e^2 B_0^2}} + 2NeB_0 \sqrt{1 + \frac{1}{4e^2 B_0^2}} \right] - \frac{me^2 E_0^2}{2n^2}$$

So the exact solution of (2.1) for parallel case is written as

$$\Phi_{\parallel} = e^{i(xP_x - \epsilon t)} W_{\lambda, \mu}(2\gamma y) {}_1F_1(z)$$

**2.2. Case (ii) Orthogonal EM Fields.** For the choice of  $A_0 = \frac{E_0}{y}$ ,  $A_1 = \frac{B_0}{y}$ ,  $\vec{E} \perp \vec{B}$ . In this case, we will look for the solution of (2.1) as

$$\Phi_{\perp} = e^{i(xP_x + zP_z - \epsilon t)} M(y)$$

Writing this in (2.1), we obtain

$$\left[ \frac{d^2}{dy^2} - \frac{e^2 B_0^2}{y^2} + \frac{2e(P_x B_0 - mE_0)}{y} + (2m\epsilon - P_x^2 - P_z^2) \right] M(y) = 0$$

Again solution of this equation is given by Whittaker function as

$$M(y) = W_{\kappa, \sigma}(2uy)$$

where  $\kappa = \frac{e(P_x - mE_0)}{iu}$ ,  $\sigma^2 = \frac{1}{4} - e^2 B_0^2$ ,  $u^2 = (2m\epsilon - P_x^2 - P_z^2)$ . For Whittaker functions,

$$\sigma - \kappa = -(n + 1/2)$$

should be satisfied. From this condition, we obtain the energy spectrum for the Schrödinger equation for the orthogonal case

$$\epsilon_{\perp} = \frac{1}{2m} \left[ P_x^2 + P_z^2 - \frac{e^2 P_x^2 B_0^2 + e^2 m^2 E_0^2 - 2e^2 m P_x B_0 E_0}{\frac{1}{4} + e^2 B_0^2 + N^2 + 2N\sqrt{\frac{1}{4} + e^2 B_0^2}} \right]$$

So the exact solution of (2.1) in orthogonal case is written as

$$\Phi_{\perp} = e^{i(xP_x + zP_z - \epsilon t)} W_{\kappa, \sigma}(2uy)$$

### 3. SOLUTION OF THE KLEIN-GORDON EQUATION

The Klein-Gordon equation for the relativistic spinless particles is given by (we take  $\hbar = 1$ )

$$(3.1) \quad \left[ (\vec{P} - e\vec{A})^2 + m^2 \right] \phi = (P_0 - eA_0)^2 \phi$$

**3.1. Case (i) Parallel EM Fields.** Again we will look for the solution as

$$\phi_{\parallel} = e^{i(xP_x - \epsilon t)} F(y)G(z)$$

writing this in (3.1) we obtain

$$\left[ \underbrace{-\frac{d^2}{dy^2} + (P_x - \frac{eB_0}{y})^2}_{\hat{Q}(y)} - \underbrace{\frac{d^2}{dz^2} - (\epsilon - \frac{eE_0}{z})^2}_{\hat{R}(z)} + m^2 \right] F(y)G(z) = 0$$

In short we can write

$$\left[ \hat{Q}(y) + \hat{R}(z) + m^2 \right] F(y)G(z) = 0$$

Separating this equation with respect to y and z, we obtain

$$(3.2) \quad \left[ \hat{Q}(y) + s \right] F(y) = 0$$

$$(3.3) \quad \left[ \hat{R}(z) + m^2 - s \right] G(z) = 0$$

where s is the separation constant.

Equation (3.2) is written as

$$(3.4) \quad \left[ -\frac{d^2}{dy^2} + (P_x - \frac{eB_0}{y})^2 + s \right] F(y) = 0$$

This equation is the same equation obtained in the Schrödinger case. So the solution is

$$F(y) = W_{\lambda, \mu}(2\gamma y)$$

where  $\mu = \pm \sqrt{\frac{1}{4} + e^2 B_0^2}$ ,  $\lambda = \frac{eB_0}{\sqrt{1 + \frac{s}{P_x^2}}}$  and  $\gamma = \sqrt{P_x^2 + s}$

As before

$$s = P_x^2 \left( \frac{1}{1 + \frac{(1/4+N^2)}{e^2 B_0^2} + 2NeB_0 \sqrt{1 + \frac{1}{4e^2 B_0^2}}} - 1 \right)$$

Equation (3.3) is written as

$$(3.5) \quad \left[ -\frac{d^2}{dz^2} - \left( \epsilon - \frac{eE_0}{z} \right)^2 + m^2 - s \right] G(z) = 0$$

Again solution of this equation is given by Whittaker functions

$$G(z) = W_{\tilde{\lambda}, \tilde{\mu}}(2\alpha z)$$

where  $\tilde{\lambda} = \frac{ieE_0\epsilon}{\alpha}$ ,  $\tilde{\mu} = \pm \sqrt{\frac{1}{4} - e^2 E_0^2}$ , and  $\alpha^2 = \epsilon^2 - m^2 + s$  and the energy spectrum for the parallel case is given by

$$\epsilon_{\parallel} = \pm \left[ \frac{(m^2 - s) \left( \frac{1}{4} - e^2 E_0^2 + N^2 + 2\tilde{N} \sqrt{\frac{1}{4} - e^2 E_0^2} \right)}{\frac{1}{4} + \tilde{N}^2 + 2\tilde{N} \sqrt{\frac{1}{4} - e^2 E_0^2}} \right]^{\frac{1}{2}}$$

and the exact solution of Klein-Gordon equation for the parallel case is given by

$$\phi_{\parallel} = e^{i(xP_x - et)} W_{\lambda, \mu}(2\gamma y) W_{\tilde{\lambda}, \tilde{\mu}}(2\alpha z)$$

**3.2. Case (ii) Orthogonal EM Fields.** For the choice of  $A_0 = \frac{E_0}{y}$ ,  $A_1 = \frac{B_0}{y}$ ,  $\vec{E} \perp \vec{B}$ .

Again we will look for the solution of (3.1) as

$$\phi_{\perp} = e^{i(xP_x + zP_z - et)} N(y)$$

Writing this in (3.1), we obtain

$$\left[ \frac{d^2}{dy^2} + \frac{e^2(E_0^2 - B_0^2)}{y^2} + \frac{2e(P_x B_0 - \epsilon E_0)}{y} + (\epsilon^2 - m^2 - P_x^2 - P_z^2) \right] N(y) = 0$$

Again solution of this equation is given by Whittaker function as

$$N(y) = W_{\tilde{\kappa}, \tilde{\sigma}}(2vy)$$

where  $\tilde{\kappa} = \frac{e(P_x - \epsilon E_0)}{iv}$ ,  $\tilde{\sigma}^2 = \frac{1}{4} - e^2(E_0^2 - B_0^2)$ ,  $v^2 = (\epsilon^2 - m^2 - P_x^2 - P_z^2)$ . For Whittaker functions,

$$\tilde{\sigma} - \tilde{\kappa} = -(\tilde{n} + 1/2)$$

should be satisfied. From this condition, we obtain the energy spectrum for the Klein-Gordon equation for the orthogonal case from below quadratic equation

$$\epsilon^2 \underbrace{(w^2 + e^2 E_0^2)}_a + \epsilon \underbrace{(-2e^2 P_x B_0 E_0)}_b + \underbrace{e^2 P_x^2 B_0^2 - w^2(m^2 + P_x^2 + P_z^2)}_c = 0$$

where  $w = \sqrt{\frac{1}{4} - e^2(E_0^2 - B_0^2)} + \tilde{N}$ .

$$\epsilon_{\perp} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Exact solution of the (3.1) for the orthogonal case is

$$\phi_{\perp} = e^{i(xP_x + zP_z - et)} W_{\tilde{\kappa}, \tilde{\sigma}}(2vy)$$




FIGURE 1. Parallel Case




FIGURE 2. Orthogonal Case

#### 4. CONCLUSION

We investigated the motion of the spin-0 particles in electromagnetic fields for parallel and orthogonal orientations. Analysis is performed for Schrödinger and Klein-Gordon cases and that present us the contribution of the relativistic effects. In the case of  $\vec{E} \parallel \vec{B}$ , the relativistic effects arise only for the motion in the  $z$ -direction. In that case the Whittaker functions that occurred in the relativistic solutions are replaced by the confluent hypergeometric function for nonrelativistic solutions. In case of the orthogonal fields  $\vec{E} \perp \vec{B}$ , exact solutions of the Schrödinger and Klein-Gordon equations are found in terms of the Whittaker functions.

#### REFERENCES

- [1] Lam, L., *Motion in electric and magnetic fields. I. Klein-Gordon particles*, Journal of Mathematical Physics, Vol. **12**, No. 2, 299-303, 1971.
- [2] Sogut K. and Havare A., *Spinless Particles in Exponentially Varying Electric and Magnetic Fields*, Advances in High Energy Physics, Volume **2014**, Article ID 493120, 2014.
- [3] Greiner W., *Relativistic Quantum Mechanics*, Springer, Berlin, Germany, 1997.
- [4] Abramowitz M. and Stegun I. A., *Handbook of Mathematical Functions*, Dover, New York, NY, USA, 1970.
- [5] Komke E., *Mathematik und Ihre Anwedungen in Physik und Technik*, Geest und Portig, Leipzig, Germany, 1967.

MERSIN UNIVERSITY, DEPARTMENT OF MATHEMATICS, MERSIN, 33343, TURKEY.  
*E-mail address:* [ozgurmizrak@gmail.com](mailto:ozgurmizrak@gmail.com)

MERSIN UNIVERSITY, DEPARTMENT OF PHYSICS, MERSIN, 33343, TURKEY.  
*E-mail address:* [kenansogut@gmail.com](mailto:kenansogut@gmail.com)

## SOME INTUITIONISTIC FUZZY MODAL OPERATORS OVER INTUITIONISTIC FUZZY IDEALS AND GROUPS

SINEM TARSUSLU(YILMAZ), GÖKHAN ÇUVALCIOĞLU, AND ARIF BAL

ABSTRACT. K.T. Atanassov generalized fuzzy sets in to Intuitionistic Fuzzy Sets in 1983[1]. Intuitionistic Fuzzy Modal Operator was firstly defined by same author and the other operators were defined by several researchers[2, 3, 4]. Intuitionistic fuzzy algebraic structures and their properties were studied in[5, 6, 7].

In this paper, we studied some intuitionistic fuzzy operators on intuitionistic fuzzy ideals and groups.

Received: 14–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

The original concept of fuzzy sets in Zadeh [9] was introduced as an extension of crisp sets by enlarging the truth value set to the real unit interval  $[0, 1]$ . In fuzzy set theory, if the membership degree of an element  $x$  is  $\mu(x)$  then the nonmembership degree is  $1 - \mu(x)$  and thus it is fixed. Intuitionistic fuzzy sets have been introduced by Atanassov in 1983 [1] and form an extension of fuzzy sets by enlarging the truth value set to the lattice  $[0, 1] \times [0, 1]$ .

**Definition 1.1.** [1] An intuitionistic fuzzy set (shortly IFS) on a set  $X$  is an object of the form

$$A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$$

where  $\mu_A(x), (\mu_A : X \rightarrow [0, 1])$  is called the “degree of membership of  $x$  in  $A$ ”,  $\nu_A(x), (\nu_A : X \rightarrow [0, 1])$  is called the “degree of non- membership of  $x$  in  $A$ ”, and where  $\mu_A$  and  $\nu_A$  satisfy the following condition:

$$\mu_A(x) + \nu_A(x) \leq 1, \text{ for all } x \in X.$$

The hesitation degree of  $x$  is defined by  $\pi_A(x) = 1 - \mu_A(x) - \nu_A(x)$

**Definition 1.2.** [1] An IFS  $A$  is said to be contained in an IFS  $B$  (notation  $A \sqsubseteq B$ ) if and only if, for all  $x \in X : \mu_A(x) \leq \mu_B(x)$  and  $\nu_A(x) \geq \nu_B(x)$ .

It is clear that  $A = B$  if and only if  $A \sqsubseteq B$  and  $B \sqsubseteq A$ .

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 03E72,47S40.

*Key words and phrases.* Intuitionistic Fuzzy Modal Operator, Intuitionistic Fuzzy Algebraic Structures.



**Definition 1.3.** [1] Let  $A \in IFS$  and let  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$  then the above set is called the complement of  $A$

$$A^c = \{ \langle x, \nu_A(x), \mu_A(x) \rangle : x \in X \}$$

**Definition 1.4.** [2] Let  $X$  be a set and  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \} \in IFS(X)$ .

$$(1) \square A = \{ \langle x, \mu_A(x), 1 - \mu_A(x) \rangle : x \in X \}$$

$$(2) \diamond A = \{ \langle x, 1 - \nu_A(x), \nu_A(x) \rangle : x \in X \}$$

**Definition 1.5.** [3] Let  $X$  be a set and  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \} \in IFS(X)$ , for  $\alpha, \beta \in I$

$$(1) \boxplus(A) = \left\{ \left\langle x, \frac{\mu_A(x)}{2}, \frac{\nu_A(x)+1}{2} \right\rangle : x \in X \right\}$$

$$(2) \boxtimes(A) = \left\{ \left\langle x, \frac{\mu_A(x)+1}{2}, \frac{\nu_A(x)}{2} \right\rangle : x \in X \right\}$$

$$(3) \boxplus_\alpha(A) = \{ \langle x, \alpha\mu_A(x), \alpha\nu_A(x) + 1 - \alpha \rangle : x \in X \}$$

$$(4) \boxtimes_\alpha(A) = \{ \langle x, \alpha\mu_A(x) + 1 - \alpha, \alpha\nu_A(x) \rangle : x \in X \}$$

$$(5) \text{ for } \max\{\alpha, \beta\} + \gamma \in I, \boxplus_{\alpha, \beta, \gamma}(A) = \{ \langle x, \alpha\mu_A(x), \beta\nu_A(x) + \gamma \rangle : x \in X \}$$

$$(6) \text{ for } \max\{\alpha, \beta\} + \gamma \in I, \boxtimes_{\alpha, \beta, \gamma}(A) = \{ \langle x, \alpha\mu_A(x) + \gamma, \beta\nu_A(x) \rangle : x \in X \}$$

**Definition 1.6.** [8] Let  $X$  be a set and  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \} \in IFS(X)$ ,  $\alpha, \beta, \alpha + \beta \in I$

$$(1) \boxplus_{\alpha, \beta}(A) = \{ \langle x, \alpha\mu_A(x), \alpha\nu_A(x) + \beta \rangle : x \in X \}$$

$$(2) \boxtimes_{\alpha, \beta}(A) = \{ \langle x, \alpha\mu_A(x) + \beta, \alpha\nu_A(x) \rangle : x \in X \}$$

The operators  $\boxplus_{\alpha, \beta, \gamma}, \boxtimes_{\alpha, \beta, \gamma}$  are an extensions of  $\boxplus_{\alpha, \beta}, \boxtimes_{\alpha, \beta}$  (resp.).

In 2007, the author [4] defined a new operator and studied some of its properties. This operator is named  $E_{\alpha, \beta}$  and defined as follows:

**Definition 1.7.** [4] Let  $X$  be a set and  $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \} \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$ . We define the following operator:

$$E_{\alpha, \beta}(A) = \{ \langle x, \beta(\alpha\mu_A(x) + 1 - \alpha), \alpha(\beta\nu_A(x) + 1 - \beta) \rangle : x \in X \}$$

If we choose  $\alpha = 1$  and write  $\alpha$  instead of  $\beta$  we get the operator  $\boxplus$ . Similarly, if  $\beta = 1$  is chosen and written instead of  $\beta$ , we get the operator  $\boxtimes_\alpha$ .

In 2007, Atanassov introduced the operator  $\square_{\alpha, \beta, \gamma, \delta}$  which is a natural extension of all these operators in [3].

**Definition 1.8.** [3] Let  $X$  be a set,  $A \in IFS(X)$ ,  $\alpha, \beta, \gamma, \delta \in [0, 1]$  such that

$$\max(\alpha, \beta) + \gamma + \delta \leq 1$$

then the operator  $\square_{\alpha, \beta, \gamma, \delta}$  defined by

$$\square_{\alpha, \beta, \gamma, \delta}(A) = \{ \langle x, \alpha\mu_A(x) + \gamma, \beta\nu_A(x) + \delta \rangle : x \in X \}$$

In 2010, the author [4] defined a new operator which is a generalization of  $E_{\alpha, \beta}$ .

**Definition 1.9.** [4] Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \omega \in [0, 1]$ . We define the following operator:

$$Z_{\alpha, \beta}^{\omega}(A) = \{ \langle x, \beta(\alpha\mu_A(x) + \omega - \omega.\alpha), \alpha(\beta\nu_A(x) + \omega - \omega.\beta) \rangle : x \in X \}$$

We have defined a new OTMO on IFS, that is generalization of the some OTMOs.  $Z_{\alpha, \beta}^{\omega, \theta}$  defined as follows:

**Definition 1.10.** [4] Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \omega, \theta \in [0, 1]$ . We define the following operator:

$$Z_{\alpha, \beta}^{\omega, \theta}(A) = \{ \langle x, \beta(\alpha\mu_A(x) + \omega - \omega.\alpha), \alpha(\beta\nu_A(x) + \theta - \theta.\beta) \rangle : x \in X \}$$

The operator  $Z_{\alpha, \beta}^{\omega, \theta}$  is a generalization of  $Z_{\alpha, \beta}^{\omega}$ , and also,  $E_{\alpha, \beta}, \boxplus_{\alpha, \beta}, \boxtimes_{\alpha, \beta}$ .

**Definition 1.11.** [5] Let  $G$  be a groupoid,  $A \in IFS(G)$ . If for all  $x, y \in G$ ,

$$A(xy) \geq \min(A(x), A(y))$$

then  $A$  called an intuitionistic fuzzy subgroupoid over  $G$ .

**Definition 1.12.** [6] Let  $G$  be a grupoid,  $A \in IFS(G)$ . If for all  $x, y \in G$ ,

$$A(xy) \geq \max(A(x), A(y))$$

then  $A$  called an intuitionistic fuzzy ideal over  $G$ , shortly  $IFI(G)$ .

**Definition 1.13.** [6] Let  $G$  be a grup and  $A \in IFS(G)$  a grupoid. If for all  $x \in G$ ,

$$A(x^{-1}) \geq A(x)$$

then  $A$  called an intuitionistic fuzzy subgroup over  $G$ , shortly  $IFG(G)$ .

## 2. MAIN RESULTS

**Theorem 2.1.** Let  $G$  be a groupoid and  $A \in IFS(G)$ .

- (1) If  $A \in IFI(G)$  then  $\square A \in IFI(G)$
- (2) If  $A \in IFI(G)$  then  $\diamond A \in IFI(G)$

*Proof.* (1) For  $x, y \in G$ ,

$$\mu_{\square A}(xy) = \mu_A(xy) \geq \mu_A(x) \vee \mu_A(y)$$

and

$$\begin{aligned} \nu_{\square A}(xy) &= 1 - \mu_A(xy) \leq (1 - \mu_A(x)) \wedge (1 - \mu_A(y)) \\ &= \nu_{\square A}(x) \wedge \nu_{\square A}(y) \end{aligned}$$

So,

$$\square A(xy) \geq \square A(x) \vee \square A(y)$$

□

**Theorem 2.2.** Let  $G$  be a groupoid and  $A \in IFS(G)$ .

- (1) If  $A \in IFI(G)$  then  $\boxplus(A) \in IFI(G)$
- (2) If  $A \in IFI(G)$  then  $\boxtimes(A) \in IFI(G)$

*Proof.* (1) For  $x, y \in G$ ,

$$\begin{aligned}\mu_{\boxplus(A)}(xy) &= \frac{\mu_A(xy)}{2} \geq \frac{\mu_A(x)}{2} \vee \frac{\mu_A(y)}{2} \\ &= \mu_{\boxplus(A)}(x) \vee \mu_{\boxplus(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{\boxplus(A)}(xy) &= \frac{\nu_A(xy) + 1}{2} \leq \frac{\nu_A(x) + 1}{2} \wedge \frac{\nu_A(y) + 1}{2} \\ &= \nu_{\boxplus(A)}(x) \wedge \nu_{\boxplus(A)}(y)\end{aligned}$$

So,

$$\boxplus(A)(xy) \geq \boxplus(A)(x) \vee \boxplus(A)(y)$$

□

**Theorem 2.3.** *Let  $G$  be a groupoid and  $A \in IFS(G)$ .*

- (1) If  $A \in IFI(G)$  then  $\boxplus_\alpha(A) \in IFI(G)$
- (2) If  $A \in IFI(G)$  then  $\boxtimes_\alpha(A) \in IFI(G)$

*Proof.* (1) For  $x, y \in G$ ,

$$\begin{aligned}\mu_{\boxtimes_\alpha(A)}(xy) &= \alpha\mu_A(xy) + 1 - \alpha \geq (\alpha\mu_A(x) + 1 - \alpha) \vee (\alpha\mu_A(y) + 1 - \alpha) \\ &= \mu_{\boxtimes_\alpha(A)}(x) \vee \mu_{\boxtimes_\alpha(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{\boxtimes_\alpha(A)}(xy) &= \alpha\nu_A(xy) \leq (\alpha\nu_A(x)) \wedge (\alpha\nu_A(y)) \\ &= \nu_{\boxtimes_\alpha(A)}(x) \vee \nu_{\boxtimes_\alpha(A)}(y)\end{aligned}$$

So,

$$\boxtimes_\alpha(A)(xy) \geq \boxtimes_\alpha(A)(x) \vee \boxtimes_\alpha(A)(y)$$

□

**Theorem 2.4.** *Let  $G$  be a groupoid and  $A \in IFS(G)$ .*

- (1) If  $A \in IFI(G)$  then  $\boxplus_{\alpha,\beta}(A) \in IFI(G)$
- (2) If  $A \in IFI(G)$  then  $\boxtimes_{\alpha,\beta}(A) \in IFSI(G)$
- (3) If  $A \in IFI(G)$  then  $\boxplus_{\alpha,\beta,\gamma}(A) \in IFI(G)$
- (4) If  $A \in IFI(G)$  then  $\boxtimes_{\alpha,\beta,\gamma}(A) \in IFI(G)$

*Proof.* For  $x, y \in G$ ,

$$\begin{aligned}\mu_{\boxplus_{\alpha,\beta,\gamma}(A)}(xy) &= \alpha\mu_A(xy) \geq \alpha\mu_A(x) \vee \alpha\mu_A(y) \\ &= \mu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x) \vee \mu_{\boxplus_{\alpha,\beta,\gamma}(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{\boxplus_{\alpha,\beta,\gamma}(A)}(xy) &= \beta\nu_A(xy) + \gamma \leq (\beta\nu_A(x) + \gamma) \wedge (\beta\nu_A(y) + \gamma) \\ &= \nu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x) \wedge \nu_{\boxplus_{\alpha,\beta,\gamma}(A)}(y)\end{aligned}$$

So,

$$\boxplus_{\alpha,\beta,\gamma}(A)(xy) \geq \boxplus_{\alpha,\beta,\gamma}(A)(x) \vee \boxplus_{\alpha,\beta,\gamma}(A)(y)$$

The other properties can proof with same way.

□

**Theorem 2.5.** *Let  $G$  be a groupoid and  $A \in IFS(G)$  an ideal then  $E_{\alpha,\beta}(A) \in IFS(G)$  is an ideal.*

*Proof.* For  $x, y \in G$ ,

$$\begin{aligned}\mu_{E_{\alpha,\beta}(A)}(xy) &= \beta(\alpha\mu_A(xy) + 1 - \alpha) \geq \beta(\alpha\mu_A(x) + 1 - \alpha) \vee \beta(\alpha\mu_A(y) + 1 - \alpha) \\ &= \mu_{E_{\alpha,\beta}(A)}(x) \vee \mu_{E_{\alpha,\beta}(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{E_{\alpha,\beta}(A)}(xy) &= \alpha(\beta\nu_A(xy) + 1 - \beta) \leq \alpha(\beta\nu_A(x) + 1 - \beta) \wedge \alpha(\beta\nu_A(y) + 1 - \beta) \\ &= \nu_{E_{\alpha,\beta}(A)}(x) \wedge \nu_{E_{\alpha,\beta}(A)}(y)\end{aligned}$$

So,

$$E_{\alpha,\beta}(A)(xy) \geq E_{\alpha,\beta}(A)(x) \vee E_{\alpha,\beta}(A)(y)$$

□

**Theorem 2.6.** *Let  $G$  be a groupoid and  $A \in IFS(G)$  an ideal then  $\square_{\alpha,\beta,\gamma,\delta}(A) \in IFS(G)$  is an ideal.*

*Proof.* For  $x, y \in G$ ,

$$\begin{aligned}\mu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(xy) &= \alpha\mu_A(xy) + \gamma \geq (\alpha\mu_A(x) + \gamma) \vee (\alpha\mu_A(y) + \gamma) \\ &= \mu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x) \vee \mu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(xy) &= \beta\nu_A(xy) + \delta \leq (\beta\nu_A(x) + \delta) \wedge (\beta\nu_A(y) + \delta) \\ &= \nu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x) \wedge \nu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(y)\end{aligned}$$

$$\text{So, } \square_{\alpha,\beta,\gamma,\delta}(A)(xy) \geq \square_{\alpha,\beta,\gamma,\delta}(A)(x) \vee \square_{\alpha,\beta,\gamma,\delta}(A)(y)$$

□

**Theorem 2.7.** *Let  $G$  be a groupoid and  $A \in IFS(G)$  an ideal then  $Z_{\alpha,\beta}^{\omega,\theta}(A) \in IFS(G)$  is an ideal.*

*Proof.* For  $x, y \in G$ ,

$$\begin{aligned}\mu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(xy) &= \beta(\alpha\mu_A(xy) + \omega - \omega.\alpha) \geq \beta(\alpha\mu_A(x) + \omega - \omega.\alpha) \vee \beta(\alpha\mu_A(y) + \omega - \omega.\alpha) \\ &= \mu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(x) \vee \mu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(y)\end{aligned}$$

and

$$\begin{aligned}\nu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(xy) &= \alpha(\beta\nu_A(xy) + \theta - \theta.\beta) \leq \alpha(\beta\nu_A(x) + \theta - \theta.\beta) \wedge \alpha(\beta\nu_A(y) + \theta - \theta.\beta) \\ &= \nu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(x) \wedge \nu_{Z_{\alpha,\beta}^{\omega,\theta}(A)}(y)\end{aligned}$$

$$\text{Therefore, we obtain } Z_{\alpha,\beta}^{\omega,\theta}(A)(xy) \geq Z_{\alpha,\beta}^{\omega,\theta}(A)(x) \vee Z_{\alpha,\beta}^{\omega,\theta}(A)(y).$$

□

**Theorem 2.8.** *Let  $G$  be a group and  $A \in IFS(G)$ .*

- (1) If  $A \in IFG(G)$  then  $\square A \in IFG(G)$ .
- (2) If  $A \in IFG(G)$  then  $\diamond A \in IFG(G)$ .

*Proof.* It is clear that, if  $A \in IFG(G)$  then it means  $A \in IFI(G)$  and for all  $x \in G$ ,  $A(x^{-1}) \geq A(x)$ .

So, it will be enough to prove the correctness of the second condition.

(2) For  $x \in G$

$$\mu_{\diamond A}(x^{-1}) = 1 - \nu_A(x^{-1}) \geq 1 - \nu_A(x) = \mu_{\diamond A}(x)$$

and

$$\nu_{\diamond A}(x^{-1}) = \nu_A(x^{-1}) \leq \nu_A(x) = \nu_{\diamond A}(x)$$

The other property can be proved same way.  $\square$

**Theorem 2.9.** *Let  $G$  be a group and  $A \in IFS(G)$ .*

- (1) If  $A \in IFG(G)$  then  $\boxplus(A) \in IFG(G)$
- (2) If  $A \in IFG(G)$  then  $\boxtimes(A) \in IFG(G)$

*Proof.* (2) For  $x, y \in G$ , If  $A \in IFG(G)$  then  $\boxtimes(A) \in IFI(G)$ . So,  $\boxtimes(A)(xy) \geq \boxtimes(A)(x) \wedge \boxtimes(A)(y)$ .

Now,

$$\mu_{\boxtimes(A)}(x^{-1}) = \frac{\mu_A(x^{-1}) + 1}{2} \geq \frac{\mu_A(x) + 1}{2} = \mu_{\boxtimes(A)}(x)$$

and

$$\nu_{\boxtimes(A)}(x^{-1}) = \frac{\nu_A(x^{-1})}{2} \leq \frac{\nu_A(x)}{2} = \nu_{\boxtimes(A)}(x)$$

Therefore,

$$\boxtimes(A)(x^{-1}) \geq \boxtimes(A)(x)$$

$\square$

**Theorem 2.10.** *Let  $G$  be a group and  $A \in IFS(G)$ .*

- (1) If  $A \in IFG(G)$  then  $\boxplus_\alpha(A) \in IFG(G)$
- (2) If  $A \in IFG(G)$  then  $\boxtimes_\alpha(A) \in IFG(G)$

*Proof.* (1) For  $x, y \in G$ , it is clear that  $\boxplus_\alpha(A)(xy) \geq \boxplus_\alpha(A)(x) \wedge \boxplus_\alpha(A)(y)$ . On the other hand,

$$\mu_{\boxplus_\alpha(A)}(x^{-1}) = \alpha\mu_A(x^{-1}) \geq \alpha\mu_A(x) = \mu_{\boxplus_\alpha(A)}(x)$$

and

$$\nu_{\boxplus_\alpha(A)}(x^{-1}) = \alpha\nu_A(x^{-1}) + 1 - \alpha \leq \alpha\nu_A(x) + 1 - \alpha = \nu_{\boxplus_\alpha(A)}(x)$$

So,

$$\boxplus_\alpha(A)(x^{-1}) \geq \boxplus_\alpha(A)(x)$$

$\square$

**Theorem 2.11.** *Let  $G$  be a group and  $A \in IFS(G)$ .*

- (1) If  $A \in IFG(G)$  then  $\boxplus_{\alpha,\beta}(A) \in IFG(G)$
- (2) If  $A \in IFG(G)$  then  $\boxtimes_{\alpha,\beta}(A) \in IFG(G)$
- (3) If  $A \in IFG(G)$  then  $\boxplus_{\alpha,\beta,\gamma}(A) \in IFG(G)$
- (4) If  $A \in IFG(G)$  then  $\boxtimes_{\alpha,\beta,\gamma}(A) \in IFG(G)$

*Proof.* For  $x, y \in G$ ,

$$\mu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x^{-1}) = \alpha\mu_A(x^{-1}) + \gamma \geq \alpha\mu_A(x) + \gamma = \mu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x)$$

and

$$\nu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x^{-1}) = \beta\nu_A(x^{-1}) \leq \beta\nu_A(x) = \nu_{\boxplus_{\alpha,\beta,\gamma}(A)}(x)$$

So,

$$\boxplus_{\alpha,\beta,\gamma}(A)(x^{-1}) \geq \boxplus_{\alpha,\beta,\gamma}(A)(x)$$

The other properties can proof with same way.  $\square$

**Theorem 2.12.** *Let  $G$  be a group and  $A \in IFS(G)$ . If  $A$  is an intuitionistic fuzzy subgroup on  $G$  then  $E_{\alpha,\beta}(A) \in IFG(G)$ .*

*Proof.* It is clear that for  $x, y \in G$ ,  $E_{\alpha,\beta}(A)(xy) \geq E_{\alpha,\beta}(A)(x) \wedge E_{\alpha,\beta}(A)(y)$ .

$$\begin{aligned}\mu_{E_{\alpha,\beta}(A)}(x^{-1}) &= \beta(\alpha\mu_A(x^{-1}) + 1 - \alpha) \geq \beta(\alpha\mu_A(x) + 1 - \alpha) \\ &= \mu_{E_{\alpha,\beta}(A)}(x)\end{aligned}$$

and

$$\begin{aligned}\nu_{E_{\alpha,\beta}(A)}(x^{-1}) &= \alpha(\beta\nu_A(x^{-1}) + 1 - \beta) \leq \alpha(\beta\nu_A(x) + 1 - \beta) \\ &= \nu_{E_{\alpha,\beta}(A)}(x)\end{aligned}$$

So,  $E_{\alpha,\beta}(A) \in IFG(G)$ .  $\square$

**Theorem 2.13.** *Let  $G$  be a group and  $A \in IFS(G)$  an intuitionistic fuzzy group then  $\square_{\alpha,\beta,\gamma,\delta}(A) \in IFS(G)$  is an intuitionistic fuzzy subgroup.*

*Proof.* For  $x \in G$ ,

$$\begin{aligned}\mu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x^{-1}) &= \alpha\mu_A(x^{-1}) + \gamma \geq \alpha\mu_A(x) + \gamma \\ &= \mu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x)\end{aligned}$$

and

$$\begin{aligned}\nu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x^{-1}) &= \beta\nu_A(x^{-1}) + \delta \leq \beta\nu_A(x) + \delta \\ &= \nu_{\square_{\alpha,\beta,\gamma,\delta}(A)}(x)\end{aligned}$$

Therefore  $\square_{\alpha,\beta,\gamma,\delta}(A) \in IFG(G)$ .  $\square$

**Theorem 2.14.** *Let  $G$  be a group and  $A \in IFS(G)$ . If  $A$  is an intuitionistic fuzzy subgroup on  $G$  then  $Z_{\alpha,\beta}^{\omega,\theta}(A) \in IFG(G)$ .*

*Proof.* It can shown easily.  $\square$

## REFERENCES

- [1] K.T. Atanassov, Intuitionistic Fuzzy Sets , VII ITKR.s Session, Sofia, June 1983.
- [2] K. T. Atanassov, Intuitionistic Fuzzy Sets, ,Spinger, Heidelberg, 1999.
- [3] K.T. Atanassov, Studies in Fuzziness and Soft Computing-On Intuitionistic Fuzzy Sets Theory, ISBN 978-3-642-29126-5, Springer Heidelberg, New York, 2012.
- [4] G. Çuvalcıođlu , New Tools Based on Intuitionistic Fuzzy Sets and Generalized Nets, ISBN 978-3-319-26301-4, Springer International Publishing Switzerland, (2016) 55-71.
- [5] K.Hur ,Y. Jang and H. W. Kang, Intuitionistic fuzzy Subgroupoid , Int. Jour. of Fuzzy Logic and Int. Sys., 3(1) (2003) 72-77.
- [6] R. Biswas , Intuitionistic fuzzy subgroups, Mathematical Forum, Vol. X,(1989) 37-46.
- [7] S. Yilmaz, "Intuitionistic fuzzy Modal operatörlerin, Intuitionistic fuzzy Yapılarka İlişkileri", Mersin Üniversitesi Fen Bilimleri Enstitüsü, Yüksek lisans Tezi, 2012, 75 s.
- [8] Dencheva K., Extension of intuitionistic fuzzy modal operators  $\boxplus$  and  $\boxtimes$ ,Proc.of the Second Int. IEEE Symp. Intelligent systems, Varna, June 22-24, (2004), Vol. 3, 21-22.
- [9] L.A. Zadeh, Fuzzy Sets, Information and Control, 8(1965) 338-353.

MERSIN UNIVERSITY FACULTY OF ARTS AND SCIENCES DEPARTMENT OF MATHEMATICS

*E-mail address:* [sinemyilmaz@gmail.com](mailto:sinemyilmaz@gmail.com)

*E-mail address:* [gcuvalcioglu@gmail.com](mailto:gcuvalcioglu@gmail.com)

*E-mail address:* [arif.bal.math@gmail.com](mailto:arif.bal.math@gmail.com)

## STABILITY ANALYSIS ON EFFECT OF SYSTEM RESTORE ON EPIDEMIC MODEL FOR COMPUTER VIRUSES

MEHMET EMRE ERDOGAN AND KEMAL USLU

ABSTRACT. More than 317 million new pieces of malware computer viruses or other malicious software were created last year. That means nearly one million new threats were released each day. Every year computer viruses cost homes and businesses billions of dollars in lost time and equipment. Computer viruses are continually evolving and their structures increasingly becoming more complex and transmission capabilities are becoming more powerful. So, we consider a SEIS model to demonstrate the system restore has a more effective role than antivirus softwares on virus defense. Also we have investigated the global behavior of the endemic equilibrium and we have supported our results with numerical simulation.

Received: 28–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Epidemiology is an area of medicine concerned with the identification of factors and conditions associated with the spread of an infectious process in a community. Because of a virus programs behavior is similar to the infectious process, this areas detects and strategies that may be useful for us [1]. Biological viruses enter their host through an opening after passively being breathed in, swallowed or via direct contact. Virtual viruses also enter their host passively when you insert an infected disk or open an infected e-mail attachment. Similarly to a biological virus which has to have the correct host and tissue specificity to gain a foothold a horse virus would not make a human being sick a computer virus has to be compatible with the system to gain a foothold. The damage these viruses do is also similar. Biological viruses replicate at the cost of the host damage can include pain, suffering and even death. Computer viruses slow down the computer files can become inaccessible and even lost, and sometimes the complete hard disk gets damaged [2]. Community, population, carrier, portal of entry, vector, symptom, modes of transmission, extra-host survival, immunity, susceptibility, sub-clinical, indicator, effective transfer rate, quarantine, isolation, infection, medium and culture are all terms from epidemiology that are useful in understanding and fighting computer viruses [1].

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 93A30; 97Mxx.

*Key words and phrases.* Computer Virus; Epidemiology; Mathematical Model; Global Stability; Endemic Equilibrium.

A computer virus is a program that can infect other programs by modifying them to include a possibly evolved version of itself. With this infection property, a virus can spread to the transitive closure of information flow, corrupting the integrity of information as it spreads. Given the widespread use of sharing in current computer systems, the threat of a virus causing widespread integrity corruption is significant [3]. Viruses, worms and trojans are all part of a class of software called malware. Malware or malicious code (malcode) is short for malicious software. It is code or software that is specifically designed to damage, disrupt, steal, or in general inflict some other bad or illegitimate action on data, hosts, or networks. A computer virus is a type of malware that propagates by inserting a copy of itself into and becoming part of another program. It spreads from one computer to another, leaving infections as it travels. Viruses can range in severity from causing mildly annoying effects to damaging data or software and causing denial-of-service conditions. Almost all viruses are attached to an executable file, which means the virus may exist on a system but will not be active or able to spread until a user runs or opens the malicious host file or program. When the host code is executed, the viral code is executed as well. Normally, the host program keeps functioning after it is infected by the virus. However, some viruses overwrite other programs with copies of themselves, which destroys the host program altogether. Viruses spread when the software or document they are attached to is transferred from one computer to another using the network, a disk, file sharing, or infected e-mail attachments.

The proper assessment of computer viruses in the management of information security and integrity depends on estimates of the risk and impact of computer virus incidents and an analysis of how they are influenced by various factors in the computing environment. Mathematical or computer simulation models of the transmission and control of computer viruses can be useful in synthesizing available information and providing a theoretical basis for control strategies [4]. Predicting virus outbreaks is extremely difficult due to human nature of the attacks but more importantly, detecting outbreaks early with a low probability of false alarms seems quite difficult [5]. By developing models it is possible to characterize essential properties of the attacks [6]. Consequently, anti-virus software has been developed to take precautions. In order to understand the effectiveness of the antivirus technologies, numbers of mathematical models were suggested to investigate the epidemic behaviors of computer virus. Due to analogical similarity between the computer viruses and infectious diseases biological counterparts, several propagation models of computer viruses have been proposed, and the obtained results indicate that the long-term behavior of computer virus can be predicted. One of the early triumphs of mathematical epidemiology was the formulation of a simple model that predicted behaviour very similar to the behaviour observed in countless epidemics [7]. The Kermack McKendrick model is a compartmental model based on relatively simple assumptions on the rates of flow between different classes of members of the population [8]. Thenceforward many computer viruses modelling studies have been made. These are; SI models [9-11], SIS models [12-14], SIR models [15-20], SIRS models [21-26], SAIC models [27], SEIR models [28], SEIQR-SEIQRS models [29-32], SLBS models [33-36], and some other models [37-41], have been proposed that every compartment which are Susceptible computers, Infected computers including the latent and breaking-out computers based on some



models, Recovered computers including the quarantine computers based on some models are considered to have the same connecting and disconnecting constants, on some models for the latent and breaking-out computers have the same infecting, corrupting and recovered constants. When considering today's conditions, it is clear that this situation is how inadequate. That's what we have done in this study, eliminating this deficiency we tried to develop an advanced epidemic model by thinking total of 14 separate constants for connecting, disconnecting, infecting, recovering and corrupting to the each compartment, instead of previous models have been constructed by 4-5 parametres. Furthermore, we optimized our model for the present day including the effect of system restore which is not considered so far.

So, we consider a SEIS model (see also Fig.1), where the compartments are:  $S(t)$  susceptible,  $E(t)$  exposed,  $I(t)$  infective at time  $t$ , respectively. The parameters of the model are defined as:

- D1.:** Every computer connects to the Internet with constant rate  $\alpha > 0$ .  
Respectively positive constant rates for each compartments are:  $\alpha_1$  for Susceptible,  $\alpha_2$  for Exposed,  $\alpha_3$  for Infective. Let  $\alpha = \alpha_1 + \alpha_2 + \alpha_3$ .
- D2.:** Every computer disconnects from the Internet with constant rate  $\delta > 0$ . Respectively positive constant rates for each compartments are:  $\delta_1$  for Susceptible,  $\delta_2$  for Exposed and  $\delta_3$  for Infective. Let  $\delta = \delta_1 + \delta_2 + \delta_3$ .
- D3.:** Every susceptible computer is infected with constant rate  $\beta > 0$  by infected removable storage media.
- D4.:** Every susceptible computer is infected by exposed computers with constant rate  $\gamma_1 > 0$  and infective computers with constant rate  $\gamma_2 > 0$ , where  $\gamma = \gamma_1 + \gamma_2$ .
- D5.:** Every exposed computer is transformed into infective computer with constant rate  $\eta > 0$ , and recovered with constant rate  $\theta_1 > 0$ .
- D6.:** Every infective computer is recovered with constant rate  $\theta_2 > 0$ .
- D7.:** Every infective computer is returned to susceptible computer with constant rate  $\mu_1 > 0$ , and returned to exposed computer with constant rate  $\mu_2 > 0$  by using system restore. Let  $\mu = \mu_1 + \mu_2$ .




FIGURE 1. The SEIS Model

From the definitions, model can be shown by:

$$\begin{aligned}
 (1.1) \quad & S'(t) = \alpha_1 + \theta_1 E(t) + (\theta_2 + \mu_1) I(t) - (\beta + \gamma_1 E(t) + \gamma_2 I(t) + \delta_1) S(t) \\
 & E'(t) = \alpha_2 + (\beta + \gamma_1 E(t) + \gamma_2 I(t)) S(t) - (\theta_1 + \eta + \delta_2) E(t) + \mu_2 I(t) \\
 & I'(t) = \alpha_3 + \eta E(t) - (\theta_2 + \mu_1 + \mu_2 + \delta_3) I(t)
 \end{aligned}$$

with the initial conditions  $(S(0), E(0), I(0)) \in R_+^3$ . Let  $N(t) = S(t) + E(t) + I(t)$ . Summing the system and simplifying, we get  $\frac{dN(t)}{dt} = \alpha - \delta N(t)$ , it is easy to get  $\lim_{t \rightarrow \infty} N(t) = \frac{\alpha}{\delta}$ . Hence, the following system would be obtained by system (1.1):

$$\begin{aligned}
 (1.2) \quad & E' = \alpha_2 + (\beta + \gamma_1 E + \gamma_2 I)(N - E - I) - (\theta_1 + \eta + \delta_2) E + \mu_2 I \\
 & I' = \alpha_3 + \eta E - (\theta_2 + \mu_1 + \mu_2 + \delta_3) I
 \end{aligned}$$

with initial conditions  $(E(0), I(0)) \in R_+^2$ . It is easy to verify that  $\Delta = \{(E, I) | E, I \geq 0, N \geq E + I\}$  is positively invariant for the system (1.2). As thus, we will consider the global stability of (1.2) on the set  $\Delta$ . Let us shortly overview the theory of asymptotically autonomous systems. An ordinary differential equation in  $R^n$ ,

$$(1.3) \quad x = f(t, x)$$

is called asymptotically autonomous with limit equation

$$(1.4) \quad y = g(y)$$

if  $f(t, x) \rightarrow g(x)$ ,  $t \rightarrow \infty$ , locally uniformly in  $x \in R^n$ , i.e. for  $x$  in any compact subset of  $R^n$ . For simplicity we assume that  $f(t, x)$ ,  $g(x)$  are continuous functions and locally Lipschitz in  $x$ . The  $\omega$ -limit sets,  $\omega(t_0, x_0)$ , of forward bounded solutions  $x$  to (1.3), subject to  $x(t_0) = x_0$   $x \in \omega(t_0, x_0) \Leftrightarrow x = \lim_{j \rightarrow \infty} x(t_j)$  for some sequence  $t_j \rightarrow \infty$ .

**Theorem 1.1.** *Let  $n = 2$  and  $\omega$  be the  $\omega$ -limit set of a forward bounded solution  $x$  of the asymptotically autonomous system (1.3). Assume that there exists a neighborhood of  $\omega$  which contains at most finitely many equilibria of system (1.4). Then the following trichotomy holds:*

- $\omega$  consists of an equilibrium of system (1.4).
- $\omega$  is the union of periodic orbits of system (1.4) and possibly of centers of system (1.4) that are surrounded by periodic orbits of system (1.4) lying in  $\omega$ .
- $\omega$  contains equilibria of system (1.4) that are cyclically chained to each other in  $\omega$  orbits of system (1.4).

*It follows from the Thieme's Theorem and the subsequent study that, for the purpose of understanding the behavior of the original system (1.1), it suffices to investigate its limit system (1.2) [42].*

## 2. MODEL ANALYSIS

### 2.1. Equilibrium Point.

**Theorem 2.1.** Assume  $(\gamma_1 + \gamma_2 b) \left( \frac{\alpha}{\delta} - a \right) + \mu_2 b > (b + 1) (\beta + \gamma_2 a) + (\theta_1 + \eta + \delta_2)$ .

Then the system (1.1) has a unique equilibrium point  $\bar{E} = (S_*, E_*, I_*)$ , where

$$(2.1) \quad S_* = N_* - (E_* + I_*)$$

$$(2.2) \quad E_* = \frac{-m_1 - \sqrt{m_1^2 - 4m_0 m_2}}{2m_0}$$

$$(2.3) \quad I_* = \frac{\alpha_3 + \eta E_*}{\theta_2 + \mu + \delta_3}$$

$$(2.4) \quad a = \frac{\alpha_3}{\theta_2 + \mu + \delta_3}$$

$$(2.5) \quad b = \frac{\eta}{\theta_2 + \mu + \delta_3}$$

$$(2.6) \quad m_0 = -(b + 1) (\gamma_1 + \gamma_2 b)$$

$$(2.7) \quad m_1 = (\gamma_1 + \gamma_2 b) \left( \frac{\alpha}{\delta} - a \right) - (b + 1) (\beta + \gamma_2 a) - (\theta_1 + \eta + \delta_2) + \mu_2 b$$

$$(2.8) \quad m_2 = \alpha_2 + (\beta + \gamma_2 a) \left( \frac{\alpha}{\delta} - a \right) + \mu_2 a$$

Furthermore,  $S_* + E_* + I_* > 0$ .

**Proof. 1.** If we solve the system,

$$(2.9) \quad \begin{aligned} \alpha - \delta (S + E + I) &= 0 \\ \alpha_2 + (\beta + \gamma_1 E + \gamma_2 I) \left( \frac{\alpha}{\delta} - E - I \right) - (\theta_1 + \eta + \delta_2) E + \mu_2 I &= 0 \\ \alpha_3 + \eta E - (\theta_2 + \mu + \delta_3) I &= 0 \end{aligned}$$

From the third equation of system (2.9), we can get  $I_* = \frac{\alpha_3 + \eta E_*}{\theta_2 + \mu + \delta_3}$ . Substituting this equation into the second equation of system (2.9) and rearranging terms we have,

$$\begin{aligned} &[-(b + 1) (\gamma_1 + \gamma_2 b)] E_*^2 + \left[ (\gamma_1 + \gamma_2 b) \left( \frac{\alpha}{\delta} - a \right) - (b + 1) (\beta + \gamma_2 a) \right. \\ &\quad \left. - (\theta_1 + \eta + \delta_2) + \mu_2 b \right] E_* + \alpha_2 + (\beta + \gamma_2 a) \left( \frac{\alpha}{\delta} - a \right) + \mu_2 a = 0 \end{aligned}$$

where  $a = \frac{\alpha_3}{\theta_2 + \mu + \delta_3} > 0$ ,  $b = \frac{\eta}{\theta_2 + \mu + \delta_3} > 0$ . If we substituting equations (2.6), (2.7), (2.8) into this equation, we have

$$(2.10) \quad m_0 E_*^2 + m_1 E_* + m_2 = 0$$

where  $m_0 = -(b + 1) (\gamma_1 + \gamma_2 b) < 0$ , and we can get,

$$m_1 = (\gamma_1 + \gamma_2 b) \left( \frac{\alpha}{\delta} - a \right) - (b + 1) (\beta + \gamma_2 a) - (\theta_1 + \eta + \delta_2) + \mu_2 b > 0,$$

from our assuming

$$m_2 = \alpha_2 + (\beta + \gamma_2 a) \left( \frac{\alpha}{\delta} - a \right) + \mu_2 a > 0,$$

Let the discriminant of (2.10) be  $\Delta = m_1^2 - 4m_0 m_2 > 0$ . So,  $E_* = \frac{-m_1 - \sqrt{\Delta}}{2m_0} > 0$ .

That is shows us  $\bar{E} = (S_*, E_*, I_*)$  is the unique equilibrium of the system (1.1). It is trivial to verify that  $E_* + I_* > 0$ .

## 2.2. Stability Analysis.

**Lemma 1.** *The equilibrium point  $\bar{E}$  is locally asymptotically stable.*

**Proof. 2.** *According to Theorem (1.1),*

$$(2.11) \quad \begin{aligned} N' &= \alpha - \delta N, \\ E' &= \alpha_2 + (\beta + \gamma_1 E + \gamma_2 I) (N - E - I) - (\theta_1 + \eta + \delta_2) E + \mu_2 I \\ I' &= \alpha_3 + \eta E - (\theta_2 + \mu_1 + \mu_2 + \delta_3) I \end{aligned}$$

The dynamical system (2.11) has a unique equilibrium point  $\bar{E} = (N_*, E_*, I_*)$  and locally asymptotically stable at  $\bar{E}$ . The Jacobian matrix of the linearized system of system (2.11) evaluated at  $\bar{E}$  is

$$J = \begin{pmatrix} -\delta & 0 & 0 \\ j_{21} & j_{22} & j_{23} \\ 0 & -\eta & -(\theta_2 + \mu + \delta_3) \end{pmatrix}$$

where

$$(2.12) \quad \begin{aligned} j_{21} &= \beta + \gamma_1 E_* + \gamma_2 I_*, \\ j_{22} &= -(\beta + \gamma_1 E_* + \gamma_2 I_*) + \gamma_1 (N_* - E_* - I_*) - (\theta_1 + \eta + \delta_2), \\ j_{23} &= -(\beta + \gamma_1 E_* + \gamma_2 I_*) + \gamma_2 (N_* - E_* - I_*) + \mu_2, \end{aligned}$$

in order to determine the characteristic equation,

$$\det |\lambda I_n - J| = \begin{vmatrix} \lambda + \delta & 0 & 0 \\ j_{21} & \lambda - j_{22} & j_{23} \\ 0 & \eta & \lambda + (\theta_2 + \mu + \delta_3) \end{vmatrix},$$

$$(\lambda + \delta) [(\lambda - j_{22})(\lambda + \theta_2 + \mu + \delta_3) - \eta j_{23}] = 0$$

Thus the characteristic polynomial is

$$(2.13) \quad h(\lambda) = (\lambda + \delta) (n_2 \lambda^2 + n_1 \lambda + n_0),$$

where

$$\begin{aligned} n_2 &= 1, \\ n_1 &= \theta_2 + \mu + \delta_3 + (\beta + \gamma_1 E_* + \gamma_2 I_*) - \gamma_1 (N_* - E_* - I_*) + (\theta_1 + \eta + \delta_2), \\ n_0 &= (\theta_2 + \mu + \delta_3 + \eta) (\beta + \gamma_1 E_* + \gamma_2 I_*) - (\gamma_1 (\theta_2 + \mu + \delta_3) + \eta \gamma_2) (N_* - E_* - I_*) \\ &\quad + (\theta_2 + \mu + \delta_3) (\theta_1 + \eta + \delta_2) - \eta \mu_2, \end{aligned}$$

it is obvious that

$$\alpha_2 + (\beta + \gamma_1 E_* + \gamma_2 I_*) S_* - (\theta_1 + \eta + \delta_2) E_* + \mu_2 I_* = 0$$

hence

$$\alpha_2 + \beta S_* + \gamma_1 S_* E_* + \gamma_2 S_* \frac{\alpha_3 + \eta E_*}{\theta_2 + \mu + \delta_3} - (\theta_1 + \eta + \delta_2) E_* + \mu_2 \frac{\alpha_3 + \eta E_*}{\theta_2 + \mu + \delta_3} = 0,$$

we have

$$\gamma_1 S_* + \gamma_2 S_* \frac{\eta}{\theta_2 + \mu + \delta_3} < \theta_1 + \eta + \delta_2$$

and

$$\gamma_1 S_* + \gamma_2 S_* \frac{\eta}{\theta_2 + \mu + \delta_3} + \mu_2 \frac{\eta}{\theta_2 + \mu + \delta_3} < \theta_1 + \eta + \delta_2$$

it is,

$$S_* < \frac{(\theta_1 + \eta + \delta_2)(\theta_2 + \mu + \delta_3) - \eta\mu_2}{\gamma_1(\theta_2 + \mu + \delta_3) + \gamma_2\eta}$$

from here,

$$\begin{aligned} n_1 &= \theta_2 + \mu + \delta_3 + (\beta + \gamma_1 E_* + \gamma_2 I_*) - \gamma_1 S_* + (\theta_1 + \eta + \delta_2) \\ &> \theta_2 + \mu + \delta_3 + (\beta + \gamma_1 E_* + \gamma_2 I_*) - \frac{\theta_1 + \eta + \delta_2}{1 + \frac{\gamma_2 \eta}{\gamma_1(\theta_2 + \mu + \delta_3)}} + \eta\mu_2 + (\theta_1 + \eta + \delta_2) \\ &> \theta_2 + \mu + \delta_3 + (\beta + \gamma_1 E_* + \gamma_2 I_*) + \eta\mu_2 \\ &> 0 \\ n_0 &= (\theta_2 + \mu + \delta_3 + \eta)(\beta + \gamma_1 E_* + \gamma_2 I_*) - (\gamma_1(\theta_2 + \mu + \delta_3) + \eta\gamma_2) S_* \\ &\quad + (\theta_2 + \mu + \delta_3)(\theta_1 + \eta + \delta_2) - \eta\mu_2, \\ &> \eta \left( \beta + \gamma_1 E_* + \gamma_2 \frac{\alpha_3 + \eta E_*}{\theta_2 + \mu + \delta_3} \right) \\ &> 0 \end{aligned}$$

it follows from the Hurwitz criterion [43], that three roots of (2.13) have negative reel parts. So, the claimed result follows by the Lyapunov theorem [43].

As a consequence of Theorem (2.1) to prove the global stability of the endemic equilibrium point  $\bar{E}$  of system (1.1), it sufficies to prove the global stability of  $\bar{E} = (E_*, I_*)$  for system (1.2). For that purpose, let us establish two lemmas.

**Lemma 2.** (1.2) allows no periodic solution in the interior of  $\Delta$ .

**Proof. 3.** Let,

$$\begin{aligned} f_1(E, I) &= \alpha_2 + (\beta + \gamma_1 E + \gamma_2 I) \left( \frac{\alpha}{\delta} - E - I \right) - (\theta_1 + \eta + \delta_2) E + \mu_2 I \\ f_2(E, I) &= \alpha_3 + \eta E - (\theta_2 + \mu_1 + \mu_2 + \delta_3) I \\ D(E, I) &= \frac{1}{I}. \end{aligned}$$

Then

$$\begin{aligned} \frac{\partial(Df_1)}{\partial E} + \frac{\partial(Df_2)}{\partial I} &= -\gamma_1 \left( 1 + \frac{E}{I} - \frac{\alpha}{\delta I} \right) - (\beta + \gamma_1 E + \gamma_2 I) - (\theta_1 + \eta + \delta_2) \\ &\quad - \frac{\alpha_3 + \eta E}{I^2} \\ &< 0. \end{aligned}$$

The claimed result follows from the Bendixson-Dulac criterion [43].

**Lemma 3.** System (1.2) allows no periodic solution that passes through a point on the boundary of  $\Delta$ .

**Proof. 4.** If there is a periodic solution that passing through a non-corner point on  $\partial\Delta$ , then it must be tangent to  $\partial\Delta$  at this point. On the contrary, suppose there is a periodic solution  $\Gamma$  that passes through a non-corner point  $(E, I)$  on  $\partial\Delta$ . There are three cases to be considered.

**Case-1::**  $0 < E < \frac{\alpha}{\delta}, I = 0$ . Then,  $I|_{E,I} = \alpha_3 + \eta E > 0$ , implying that  $\Gamma$  is not tangent to  $\partial\Delta$  at this point, which leads to a contradiction.

**Case-2::**  $0 < I < \frac{\alpha}{\delta}, E = 0$ . Then,  $E|_{E,I} = \alpha_2 + (\beta + \gamma_2 I) \left(\frac{\alpha}{\delta} - I\right) + \mu_2 I > 0$ , implying that  $\Gamma$  is not tangent to  $\partial\Delta$  at this point, which leads to a contradiction.

**Case-3::**  $E + I = \frac{\alpha}{\delta}, E \neq 0$  and  $I \neq 0$ . Then,  $\frac{d(E+I)}{dt}|_{E,I} = -(\theta_1 + \delta_2) E - (\theta_2 + \mu_1 + \delta_3) I < 0$ , implying that  $\Gamma$  is not tangent to  $\partial\Delta$  at this point, also a contradiction.

The claimed result follows by combining the above discussions. Hence, the proof is complete.

On this basis, we present

**Theorem 2.2.** *The equilibrium point  $\bar{E}$  is globally asymptotically stable for system (1.2).*

**Proof.** 5. *The claimed result follows by combining the generalized Poincaré-Bendixson theorem [43] with lemmas 1-3.*

### 3. NUMERICAL EXAMPLES

This section provides numerical examples for illustrating main result and the effects of System Restore and antivirus software on virus spread. In what follows, observe the asymptotic behavior of system (1.2) with varying  $\alpha_1, \alpha_2, \alpha_3, \delta_1, \delta_2, \delta_3, \beta, \eta, \mu_1, \mu_2, \theta_1, \theta_2, \gamma_1$  and  $\gamma_2$ .

In Figs. 2, 3, 4, 5 Evolutions of  $S(t), E(t), I(t)$  are performed with constant  $\alpha_1 = 0.6, \alpha_2 = 0.2, \alpha_3 = 0.3, \beta = 0.32, \eta = 0.65, \delta_1 = 0.02, \delta_2 = 0.03, \delta_3 = 0.04, \gamma_1 = 0.28, \gamma_2 = 0.42$  and initial conditions  $(S(0), E(0), I(0)) = (10, 5, 2)$ , and varying  $\mu_1, \mu_2, \theta_1$  and  $\theta_2$  which are recover with antivirus software and system restore parameters.

Whether system restore and a scanning with an antivirus software will not be done, the number of infected computers will be equivalent to the number of the total computer in the system shown by Fig.2 with  $\mu_1, \mu_2, \theta_1, \theta_2 = 0$ . Also the system of computers equilibrium point  $\bar{E}_1 = (S_1, E_1, I_1)$  equivalent to  $\bar{E}_1 = (0.0574, 1.1753, 23.3037)$  in Fig.??.

If only a scanning with an antivirus software will be done, the system which is newly formed is shown in Fig.3 with  $\mu_1, \mu_2 = 0$  and  $\theta_1 = 0.65, \theta_2 = 0.35$ . Also just system restore will be done, the system which is newly formed by new parameters is shown in Fig.4 with  $\mu_1 = 0.7, \mu_2 = 0.2$  and  $\theta_1, \theta_2 = 0$ . And also the system of computers equilibrium point  $\bar{E}_2 = (S_2, E_2, I_2)$  equivalent to  $\bar{E}_2 = (1.2676, 9.1327, 15.7034)$  in Fig.3 and the system of computers equilibrium point  $\bar{E}_3 = (S_3, E_3, I_3)$  equivalent to  $\bar{E}_3 = (0.8915, 15.2111, 10.7503)$  in Fig.4.

If system restore and a scanning with an antivirus software will be done, one can easily see in the Fig.5 with  $\mu_1 = 0.7, \mu_2 = 0.2$  and  $\theta_1 = 0.65, \theta_2 = 0.35$  that




FIGURE 2. Evolutions of S, E, I, N with  $\mu_1, \mu_2, \theta_1, \theta_2 = 0$ .




FIGURE 3. Evolutions of S, E, I, N with  $\mu_1, \mu_2 = 0$  and  $\theta_1 = 0.65, \theta_2 = 0.35$ .

quite decreasing of the number of infected computers. And the system of computers equilibrium point  $\bar{E}_4 = (S_4, E_4, I_4)$  equivalent to  $\bar{E}_4 = (2.3740, 16.6080, 8.5448)$  in Fig.5.

Additionally one can see from Fig. 2, 3, 4, 5 that the equilibrium points  $\bar{E}_{1,2,3,4}$  are globally asymptotically stable. From the figures which exhibits solutions  $S(t), E(t), I(t)$  are converging to stable state, which is consistent with the main result.

If the System Restore will not done, infected computers will be dominate the system in time. Therefore, total number of computers will be equivalent to infected




FIGURE 4. Evolutions of S, E, I, N with  $\mu_1 = 0.7, \mu_2 = 0.2$  and  $\theta_1, \theta_2 = 0$ .




FIGURE 5. Evolutions of S, E, I, N with  $\mu_1 = 0.7, \mu_2 = 0.2$  and  $\theta_1 = 0.65, \theta_2 = 0.35$ .

computers. (Fig. 2).

If the System Restore will done, we will recover non-copy datas in infected computers also total number of exposed and susceptible computers will be increase while infected computers number decreasing. (Fig. 4).

If we use system restore after have a comprehensive virus scanning, we will have prevented the pretty much deterioration of our computer. (Fig. 5).



## 4. ACKNOWLEDGEMENTS

This work was supported by the Scientific and Technological Research Council of Turkey (TUBITAK) Grant No: Bideb-2211. Also this paper is part of the PhD Thesis of M.E. Erdogan, Selcuk University, 2016.

## 5. REFERENCE

- [1] Murray, W. H. (1988). The application of epidemiology to computer viruses. *Computers and Security*, 7(2), 139145. [http://doi.org/10.1016/0167-4048\(88\)90327-6](http://doi.org/10.1016/0167-4048(88)90327-6)
- [2] Mishra, B. K., & Jha, N. (2007). Fixed period of temporary immunity after run of anti-malicious software on computer nodes. *Applied Mathematics and Computation*, 190(2), 12071212. <http://doi.org/10.1016/j.amc.2007.02.004>
- [3] Cohen, F. (1989). Models of practical defenses against computer viruses. *Computers and Security*, 8(2), 149160. [http://doi.org/10.1016/0167-4048\(89\)90070-9](http://doi.org/10.1016/0167-4048(89)90070-9)
- [4] Aron, J.L., Gove, R.A., Azadegan, S., Schneider, M.C.: The Benefits of a Notification Process in Addressing the Worsening Computer Virus Problem: Results of a Survey and a Simulation Model. *Computers & Security* 20(8), 693714(2001). doi:10.1016/S0167-4048(01)00812-4
- [5] Capasso, V. (1993). *Mathematical Structures of Epidemic Systems* (Vol. 1993). Berlin, Heidelberg: Springer Berlin Heidelberg. <http://doi.org/10.1007/978-3-540-70514-7>
- [6] Mishra, B. K., & Saini, D. (2007). Mathematical models on computer viruses. *Applied Mathematics and Computation*, 187(2), 929936. <http://doi.org/10.1016/j.amc.2006.09.062>
- [7] Brauer, F., van den Driessche, P., & Wu, J. (Eds.). (2008). *Mathematical Epidemiology* (Vol. 1945). Berlin, Heidelberg: Springer Berlin Heidelberg. <http://doi.org/10.1007/978-3-540-78911-6>
- [8] Kermack, W. O., & McKendrick, A. G. (1927). A Contribution to the Mathematical Theory of Epidemics. *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*. <http://doi.org/10.1098/rspa.1927.0118>
- [9] Liu, B., Duan, Y., & Luan, S. (2012). Dynamics of an SI epidemic model with external effects in a polluted environment. *Nonlinear Analysis: Real World Applications*, 13(1), 2738. <http://doi.org/10.1016/j.nonrwa.2011.07.007>
- [10] Ren, J., Yang, X., Yang, L. X., Xu, Y., & Yang, F. (2012). A delayed computer virus propagation model and its dynamics. *Chaos, Solitons and Fractals*, 45(1), 7479. <http://doi.org/10.1016/j.chaos.2011.10.003>
- [11] Chen, L., & Sun, J. (2014). Global stability of an SI epidemic model with feedback controls. *Applied Mathematics Letters*, 28, 5355. <http://doi.org/10.1016/j.aml.2013.09.009>
- [12] Kephart, J. O., & White, S. R. (1991). Directed-graph epidemiological models of computer viruses. *Proceedings. 1991 IEEE Computer Society Symposium on Research in Security and Privacy*, 343359. <http://doi.org/10.1109/RISP.1991.130801>
- [13] Billings, L., Spears, W.M., Schwartz, I.B.: A unified prediction of computer virus spread in connected networks. *Physics Letters, Section A: General, Atomic and Solid State Physics* 297(3-4), 261266(2002). doi:10.1016/S0375-9601(02)00152-4

- [14] Wierman, J.C., Marchette, D.J.: Modeling computer virus prevalence with a susceptible - infected - susceptible model with reintroduction. *Computational Statistics and Data Analysis* 45(1), 323(2004). doi:10.1016/S0167-9473(03)00113-0
- [15] Song, L. P., Jin, Z., Sun, G. Q., Zhang, J., & Han, X. (2011). Influence of removable devices on computer worms: Dynamic analysis and control strategies. *Computers and Mathematics with Applications*, 61(7), 18231829. <http://doi.org/10.1016/j.camwa.2011.02.010>,
- [16] Ren, J., Yang, X., Yang, L. X., Xu, Y., & Yang, F. (2012). A delayed computer virus propagation model and its dynamics. *Chaos, Solitons and Fractals*, 45(1), 7479. <http://doi.org/10.1016/j.chaos.2011.10.003>
- [17] Zhu, Q., Yang, X., Yang, L. X., & Zhang, C. (2012). Optimal control of computer virus under a delayed model. *Applied Mathematics and Computation*, 218(23), 1161311619. <http://doi.org/10.1016/j.amc.2012.04.092>
- [18] Gan, C., Yang, X., Liu, W., Zhu, Q., Jin, J., & He, L. (2014). Propagation of computer virus both across the Internet and external computers: A complex-network approach. *Communications in Nonlinear Science and Numerical Simulation*, 19(8), 27852792. <http://doi.org/10.1016/j.cnsns.2013.12.026>
- [19] Gan, C., Yang, X., Liu, W., Zhu, Q., & Zhang, X. (2013). An epidemic model of computer viruses with vaccination and generalized nonlinear incidence rate. *Applied Mathematics and Computation*, 222, 265274. <http://doi.org/10.1016/j.amc.2013.07.055>
- [20] Clancy, D. (2014). SIR epidemic models with general infectious period distribution. *Statistics and Probability Letters*, 85(1), 15. <http://doi.org/10.1016/j.spl.2013.10.017>
- [21] Mena-Lorcat, J., & Hethcote, H. W. (1992). Dynamic models of infectious diseases as regulators of population sizes. *Journal of Mathematical Biology*, 30(7), 693716. <http://doi.org/10.1007/BF00173264>
- [22] Han, X., & Tan, Q. (2010). Dynamical behavior of computer virus on Internet. *Applied Mathematics and Computation*, 217(6), 25202526. <http://doi.org/10.1016/j.amc.2010.07.064>
- [23] Enatsu, Y., Nakata, Y., & Muroya, Y. (2012). Lyapunov functional techniques for the global stability analysis of a delayed SIRS epidemic model. *Nonlinear Analysis: Real World Applications*, 13(5), 21202133. <http://doi.org/10.1016/j.nonrwa.2012.01.007>
- [24] Amador, J., & Artalejo, J. R. (2013). Stochastic modeling of computer virus spreading with warning signals. *Journal of the Franklin Institute*, 350(5), 11121138. <http://doi.org/10.1016/j.jfranklin.2013.02.008>
- [25] Li, C. H., Tsai, C. C., & Yang, S. Y. (2014). Analysis of epidemic spreading of an SIRS model in complex heterogeneous networks. *Communications in Nonlinear Science and Numerical Simulation*, 19(4), 10421054. <http://doi.org/10.1016/j.cnsns.2013.08.033>
- [26] Muroya, Y., Li, H., & Kuniya, T. (2014). Complete global analysis of an SIRS epidemic model with graded cure and incomplete recovery rates. *Journal of Mathematical Analysis and Applications*, 410(2), 719732. <http://doi.org/10.1016/j.jmaa.2013.08.024>
- [27] Piqueira, J. R. C., de Vasconcelos, A. A., Gabriel, C. E. C. J., & Araujo, V. O. (2008). Dynamic models for computer viruses. *Computers and Security*, 27(7-8), 355359. <http://doi.org/10.1016/j.cose.2008.07.006>

- [28] Yuan, H., Chen, G., Wu, J., & Xiong, H. (2009). Towards controlling virus propagation in information systems with point-to-group information sharing. *Decision Support Systems*, 48(1), 5768. <http://doi.org/10.1016/j.dss.2009.05.014>
- [29] Pei, Y., Liu, S., Gao, S., Li, S., & Li, C. (2009). A delayed SEIQR epidemic model with pulse vaccination and the quarantine measure. *Computers and Mathematics with Applications*, 58(1), 135145. <http://doi.org/10.1016/j.camwa.2009.01.037>
- [30] Mishra, B. K., & Jha, N. (2010). SEIQRS model for the transmission of malicious objects in computer network. *Applied Mathematical Modelling*, 34(3), 710715. <http://doi.org/10.1016/j.apm.2009.06.011>
- [31] Wang, Z., Fan, X., & Han, Q. (2013). Global stability of deterministic and stochastic multigroup SEIQR models in computer network. *Applied Mathematical Modelling*, 37(20-21), 86738686. <http://doi.org/10.1016/j.apm.2013.07.037>
- [32] Huang, G. (2016). Artificial infectious disease optimization: A SEIQR epidemic dynamic model-based function optimization algorithm. *Swarm and Evolutionary Computation*, 27, 3167. <http://doi.org/10.1016/j.swevo.2015.09.007>
- [33] Yang, L. X., & Yang, X. (2012). The spread of computer viruses under the influence of removable storage devices. *Applied Mathematics and Computation*, 219(8), 39143922. <http://doi.org/10.1016/j.amc.2012.10.027>
- [34] Yang, L. X., & Yang, X. (2014). A new epidemic model of computer viruses. *Communications in Nonlinear Science and Numerical Simulation*, 19(6), 19351944. <http://doi.org/10.1016/j.cnsns.2013.09.038>
- [35] Chen, L., Hattaf, K., & Sun, J. (2015). Optimal control of a delayed SLBS computer virus model. *Physica A: Statistical Mechanics and Its Applications*, 427, 244250. <http://doi.org/10.1016/j.physa.2015.02.048>
- [36] Gan, C., & Yang, X. (2015). Theoretical and experimental analysis of the impacts of removable storage media and antivirus software on viral spread. *Communications in Nonlinear Science and Numerical Simulation*, 22(1-3), 167174. <http://doi.org/10.1016/j.cnsns.2014.10.009>
- [37] Gan, C., Yang, X., Liu, W., Zhu, Q., Jin, J., & He, L. (2014). Propagation of computer virus both across the Internet and external computers: A complex-network approach. *Communications in Nonlinear Science and Numerical Simulation*, 19(8), 27852792. <http://doi.org/10.1016/j.cnsns.2013.12.026>
- [38] Li, T., Wang, Y., & Guan, Z. H. (2014). Spreading dynamics of a SIQRS epidemic model on scale-free networks. *Communications in Nonlinear Science and Numerical Simulation*, 19(3), 686692. <http://doi.org/10.1016/j.cnsns.2013.07.010>
- [39] MUROYA, Y., LI, H., & KUNIYA, T. (2014). On global stability of a nonresident computer virus model. *Acta Mathematica Scientia*, 34(5), 14271445. [http://doi.org/10.1016/S0252-9602\(14\)60094-1](http://doi.org/10.1016/S0252-9602(14)60094-1)
- [40] Ahn, I., Oh, H. C., & Park, J. (2015). Investigation of the C-SEIRA model for controlling malicious code infection in computer networks. *Applied Mathematical Modelling*, 39(14), 41214133. <http://doi.org/10.1016/j.apm.2014.12.038>
- [41] Ren, J., Xu, Y., & Liu, J. (2015). Investigation of dynamics of a virus-antivirus model in complex network. *Physica A: Statistical Mechanics and Its Applications*, 421(1), 533540. <http://doi.org/10.1016/j.physa.2014.11.019>
- [42] Thieme, H. (1992). Convergence results and a Poincare-Bendixson trichotomy for asymptotically autonomous differential equations. *Journal of Mathematical Biology*, 30(7), 755763. <http://doi.org/10.1007/BF00173267>

[43] Robinson, R. C. (2004). An Introduction to Dynamical Systems: Continuous and Discrete. Pearson Prentice Hall. Retrieved from <https://books.google.ca/books?id=ZGUnAQAAIAAJ>

HUGLU VOCATIONAL HIGH SCHOOL, SELCUK UNIVERSITY  
*E-mail address:* `m.emre448@hotmail.com`

DEPARTMENT OF MATHEMATICS, SELCUK UNIVERSITY  
*E-mail address:* `kuslu@selcuk.edu.tr`

**BOUNDEDNESS OF THE SUBLINEAR OPERATORS WITH  
ROUGH KERNEL GENERATED BY CALDERÓN–ZYGMUND  
OPERATORS AND THEIR COMMUTATORS ON  
GENERALIZED VANISHING MORREY SPACES**

FERIT GURBUZ

ABSTRACT. In this paper, we are interested in the boundedness of sublinear operators with rough kernel generated by Calderón–Zygmund operators on generalized vanishing Morrey spaces and give bounded mean oscillation space estimates for their commutators on these spaces.

Received: 27–July–2016

Accepted: 29–August–2016

1. INTRODUCTION

The classical Morrey spaces  $M_{p,\lambda}$  have been introduced by Morrey in [32] to study the local behavior of solutions of second order elliptic partial differential equations(PDEs). Later, there are many applications of Morrey space to the Navier-Stokes equations (see [29]), the Schrödinger equations (see [40]) and the elliptic problems with discontinuous coefficients (see [3, 13, 35]).

Let  $B = B(x_0, r_B)$  denote the ball with the center  $x_0$  and radius  $r_B$ . For a given measurable set  $E$ , we also denote the Lebesgue measure of  $E$  by  $|E|$ . For any given  $\Omega_0 \subseteq \mathbb{R}^n$  and  $0 < p < \infty$ , denote by  $L_p(\Omega_0)$  the spaces of all functions  $f$  satisfying

$$\|f\|_{L_p(\Omega_0)} = \left( \int_{\Omega_0} |f(x)|^p dx \right)^{\frac{1}{p}} < \infty.$$

We recall the definition of classical Morrey spaces  $M_{p,\lambda}$  as

$$M_{p,\lambda}(\mathbb{R}^n) = \left\{ f : \|f\|_{M_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,r))} < \infty \right\},$$

where  $f \in L_p^{loc}(\mathbb{R}^n)$ ,  $0 \leq \lambda \leq n$  and  $1 \leq p < \infty$ .

Note that  $M_{p,0} = L_p(\mathbb{R}^n)$  and  $M_{p,n} = L_\infty(\mathbb{R}^n)$ . If  $\lambda < 0$  or  $\lambda > n$ , then  $M_{p,\lambda} = \Theta$ , where  $\Theta$  is the set of all functions equivalent to 0 on  $\mathbb{R}^n$ .

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 42B20, 42B25, 42B35.

*Key words and phrases.* Sublinear operator; Calderón–Zygmund operator; rough kernel; generalized Morrey space; generalized vanishing Morrey spaces; commutator; BMO.

We also denote by  $WM_{p,\lambda} \equiv WM_{p,\lambda}(\mathbb{R}^n)$  the weak Morrey space of all functions  $f \in WL_p^{loc}(\mathbb{R}^n)$  for which

$$\|f\|_{WM_{p,\lambda}} \equiv \|f\|_{WM_{p,\lambda}(\mathbb{R}^n)} = \sup_{x \in \mathbb{R}^n, r > 0} r^{-\frac{\lambda}{p}} \|f\|_{WL_p(B(x,r))} < \infty,$$

where  $WL_p(B(x,r))$  denotes the weak  $L_p$ -space of measurable functions  $f$  for which

$$\begin{aligned} \|f\|_{WL_p(B(x,r))} &\equiv \|f\chi_{B(x,r)}\|_{WL_p(\mathbb{R}^n)} \\ &= \sup_{t > 0} t |\{y \in B(x,r) : |f(y)| > t\}|^{1/p} \\ &= \sup_{0 < t \leq |B(x,r)|} t^{1/p} (f\chi_{B(x,r)})^*(t) < \infty, \end{aligned}$$

Where  $g^*$  denotes the non-increasing rearrangement of a function  $g$ .

Throughout the paper we assume that  $x \in \mathbb{R}^n$  and  $r > 0$  and also let  $B(x,r)$  denotes the open ball centered at  $x$  of radius  $r$ ,  $B^C(x,r)$  denotes its complement and  $|B(x,r)|$  is the Lebesgue measure of the ball  $B(x,r)$  and  $|B(x,r)| = v_n r^n$ , where  $v_n = |B(0,1)|$ .

Morrey has stated that many properties of solutions to PDEs can be attributed to the boundedness of some operators on Morrey spaces. For the boundedness of the Hardy–Littlewood maximal operator, the fractional integral operator and the Calderón–Zygmund singular integral operator on these spaces, we refer the readers to [1, 5, 37]. For the properties and applications of classical Morrey spaces, see [6, 7, 12, 13] and references therein.

The study of the operators of harmonic analysis in vanishing Morrey space, in fact has been almost not touched. A version of the classical Morrey space  $M_{p,\lambda}(\mathbb{R}^n)$  where it is possible to approximate by "nice" functions is the so called vanishing Morrey space  $VM_{p,\lambda}(\mathbb{R}^n)$  has been introduced by Vitanza in [51] and has been applied there to obtain a regularity result for elliptic PDEs. This is a subspace of functions in  $M_{p,\lambda}(\mathbb{R}^n)$ , which satisfies the condition

$$\lim_{r \rightarrow 0} \sup_{\substack{x \in \mathbb{R}^n \\ 0 < t < r}} t^{-\frac{\lambda}{p}} \|f\|_{L_p(B(x,t))} = 0.$$

Later in [52] Vitanza has proved an existence theorem for a Dirichlet problem, under weaker assumptions than in [30] and a  $W^{3,2}$  regularity result assuming that the partial derivatives of the coefficients of the highest and lower order terms belong to vanishing Morrey spaces depending on the dimension. Also Ragusa has proved a sufficient condition for commutators of fractional integral operators to belong to vanishing Morrey spaces  $VM_{p,\lambda}(\mathbb{R}^n)$  (see [38, 39]). For the properties and applications of vanishing Morrey spaces, see also [4]. It is known that, there is no research regarding boundedness of the sublinear operators with rough kernel on vanishing Morrey spaces.

Maximal functions and singular integrals play a key role in harmonic analysis since maximal functions could control crucial quantitative information concerning the given functions, despite their larger size, while singular integrals, Hilbert transform as it's prototype, recently intimately connected with PDEs, operator theory and other fields.

Let  $f \in L^{loc}(\mathbb{R}^n)$ . The Hardy-Littlewood(H-L) maximal operator  $M$  is defined by

$$Mf(x) = \sup_{t>0} |B(x,t)|^{-1} \int_{B(x,t)} |f(y)| dy.$$

Let  $\bar{T}$  be a standard Calderón-Zygmund(C-Z) singular integral operator, briefly a C-Z operator, i.e., a linear operator bounded from  $L_2(\mathbb{R}^n)$  to  $L_2(\mathbb{R}^n)$  taking all infinitely continuously differentiable functions  $f$  with compact support to the functions  $f \in L_1^{loc}(\mathbb{R}^n)$  represented by

$$\bar{T}f(x) = p.v. \int_{\mathbb{R}^n} k(x-y)f(y) dy \quad x \notin \text{supp}f.$$

Such operators have been introduced in [10]. Here  $k$  is a C-Z kernel [16]. Chiarenza and Frasca [5] have obtained the boundedness of H-L maximal operator  $M$  and C-Z operator  $\bar{T}$  on  $M_{p,\lambda}(\mathbb{R}^n)$ . It is also well known that H-L maximal operator  $M$  and C-Z operator  $\bar{T}$  play an important role in harmonic analysis (see [15, 27, 47, 48, 49]). Also, the theory of the C-Z operator is one of the important achievements of classical analysis in the last century, which has many important applications in Fourier analysis, complex analysis, operator theory and so on.

Suppose that  $S^{n-1}$  is the unit sphere in  $\mathbb{R}^n$  ( $n \geq 2$ ) equipped with the normalized Lebesgue measure  $d\sigma$ . Let  $\Omega \in L_q(S^{n-1})$  with  $1 < q \leq \infty$  be homogeneous of degree zero. Suppose that  $T_\Omega$  represents a linear or a sublinear operator, which satisfies that for any  $f \in L_1(\mathbb{R}^n)$  with compact support and  $x \notin \text{supp}f$

$$(1.1) \quad |T_\Omega f(x)| \leq c_0 \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^n} |f(y)| dy,$$

where  $c_0$  is independent of  $f$  and  $x$ .

For a locally integrable function  $b$  on  $\mathbb{R}^n$ , suppose that the commutator operator  $T_{\Omega,b}$  represents a linear or a sublinear operator, which satisfies that for any  $f \in L_1(\mathbb{R}^n)$  with compact support and  $x \notin \text{supp}f$

$$(1.2) \quad |T_{\Omega,b}f(x)| \leq c_0 \int_{\mathbb{R}^n} |b(x) - b(y)| \frac{|\Omega(x-y)|}{|x-y|^n} |f(y)| dy,$$

where  $c_0$  is independent of  $f$  and  $x$ .

We point out that condition (1.1) in the case of  $\Omega \equiv 1$  has been introduced by Soria and Weiss in [45]. Conditions (1.1) and (1.2) are satisfied by many interesting operators in harmonic analysis, such as Marcinkiewicz operator, the C-Z operators, Carleson's maximal operator, H-L maximal operator, C. Fefferman's singular multipliers, R. Fefferman's singular integrals, Ricci-Stein's oscillatory singular integrals, the Bochner-Riesz means and so on (see [25], [45] for details).

Let  $\Omega \in L_q(S^{n-1})$  with  $1 < q \leq \infty$  be homogeneous of degree zero and satisfies the cancellation condition

$$\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0,$$

where  $x' = \frac{x}{|x|}$  for any  $x \neq 0$ . The C-Z singular integral operator with rough kernel  $\bar{T}_\Omega$  is defined by

$$\bar{T}_\Omega f(x) = p.v. \int_{\mathbb{R}^n} \frac{\Omega(x-y)}{|x-y|^n} f(y) dy,$$

satisfies condition (1.1).

It is obvious that when  $\Omega \equiv 1$ ,  $\bar{T}_\Omega$  is the C-Z operator  $\bar{T}$ .

The case when  $\Omega$  is a smooth kernel and  $\bar{T}_\Omega$  a standard C-Z singular integral operator has been fully studied by many authors (see [16]).

In 1976, Coifman, Rocherberg and Weiss [8] introduced the commutator generated by  $\bar{T}_\Omega$  and a local integrable function  $b$  as follows:

$$(1.3) \quad [b, \bar{T}_\Omega]f(x) \equiv b(x)\bar{T}_\Omega f(x) - \bar{T}_\Omega(bf)(x) = p.v. \int_{\mathbb{R}^n} [b(x) - b(y)] \frac{\Omega(x-y)}{|x-y|^n} f(y) dy.$$

Sometimes, the commutator defined by (1.3) is also called the commutator in Coifman-Rocherberg-Weiss's sense, which has its root in the complex analysis and harmonic analysis (see [8]).

*Remark 1.1.* [43, 44] When  $\Omega$  satisfies the specified size conditions, the kernel of the operator  $\bar{T}_\Omega$  has no regularity, so the operator  $\bar{T}_\Omega$  is called a rough C-Z singular integral operator. In recent years, a variety of operators related to the C-Z singular integral operators, but lacking the smoothness required in the classical theory, have been studied. These include the operator  $[b, \bar{T}_\Omega]$ . For more results, we refer the reader to [2, 18, 19, 20, 26, 27].

In this paper, we prove the boundedness of certain sublinear operators with rough kernel  $T_\Omega$  satisfying condition (1.1), generated by C-Z singular integral operators on generalized vanishing Morrey spaces  $VM_{p,\varphi}$  for  $q' \leq p$ ,  $p \neq 1$  or  $p < q$ , where  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$  is a homogeneous of degree zero. The boundedness of the commutators of sublinear operators  $T_{\Omega,b}$  satisfying condition (1.2) on generalized vanishing Morrey spaces are also obtained. Provided that  $b \in BMO$  and  $T_{\Omega,b}$  is a sublinear operator, we obtain the sufficient conditions on the pair  $(\varphi_1, \varphi_2)$  which ensures the boundedness of the operators  $T_{\Omega,b}$ , from one vanishing generalized Morrey space  $VM_{p,\varphi_1}$  to another  $VM_{p,\varphi_2}$ , where  $1 < p < \infty$ . In all the cases the conditions for the boundedness of  $T_\Omega$  and  $T_{\Omega,b}$  are given in terms of Zygmund-type integral inequalities on  $(\varphi_1, \varphi_2)$ , where there is no assumption on monotonicity of  $\varphi_1, \varphi_2$  in  $r$ . As an example to the conditions of these theorems are satisfied, we can consider the Marcinkiewicz operator.

Finally, we present a relationship between essential supremum and essential infimum.

**Lemma 1.1.** (see [53] page 143) *Let  $f$  be a real-valued nonnegative function and measurable on  $E$ . Then*

$$(1.4) \quad \left( \operatorname{ess\,inf}_{x \in E} f(x) \right)^{-1} = \operatorname{ess\,sup}_{x \in E} \frac{1}{f(x)}.$$

By  $A \lesssim B$  we mean that  $A \leq CB$  with some positive constant  $C$  independent of appropriate quantities. If  $A \lesssim B$  and  $B \lesssim A$ , we write  $A \approx B$  and say that  $A$  and  $B$  are equivalent.



## 2. GENERALIZED VANISHING MORREY SPACES

After studying Morrey spaces in detail, researchers have passed to generalized Morrey spaces. Mizuhara [31] has given generalized Morrey spaces  $M_{p,\varphi}$  considering  $\varphi = \varphi(r)$  instead of  $r^\lambda$  in the above definition of the Morrey space. Later, Guliyev [14] has defined the generalized Morrey spaces  $M_{p,\varphi}$  with normalized norm as follows:

**Definition 2.1.** [14] Let  $\varphi(x, r)$  be a positive measurable function on  $\mathbb{R}^n \times (0, \infty)$  and  $1 \leq p < \infty$ . We denote by  $M_{p,\varphi} \equiv M_{p,\varphi}(\mathbb{R}^n)$  the generalized Morrey space, the space of all functions  $f \in L_p^{loc}(\mathbb{R}^n)$  with finite quasinorm

$$\|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))}.$$

Also by  $WM_{p,\varphi} \equiv WM_{p,\varphi}(\mathbb{R}^n)$  we denote the weak generalized Morrey space of all functions  $f \in WL_p^{loc}(\mathbb{R}^n)$  for which

$$\|f\|_{WM_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \varphi(x, r)^{-1} |B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x, r))} < \infty.$$

According to this definition, we recover the Morrey space  $M_{p,\lambda}$  and weak Morrey space  $WM_{p,\lambda}$  under the choice  $\varphi(x, r) = r^{\frac{\lambda-n}{p}}$ :

$$M_{p,\lambda} = M_{p,\varphi} \Big|_{\varphi(x,r)=r^{\frac{\lambda-n}{p}}}, \quad WM_{p,\lambda} = WM_{p,\varphi} \Big|_{\varphi(x,r)=r^{\frac{\lambda-n}{p}}}.$$

Everywhere in the sequel we assume that  $\inf_{x \in \mathbb{R}^n, r > 0} \varphi(x, r) > 0$  which makes the above spaces non-trivial, since the spaces of bounded functions are contained in these spaces.

In [14, 23, 24, 31, 34], the boundedness of the maximal operator and C–Z singular integral operator on the generalized Morrey spaces has been obtained. For generalized Morrey spaces with nondoubling measures see also [42].

For brevity, in the sequel we use the notations

$$\mathfrak{M}_{p,\varphi}(f; x, r) := \frac{|B(x, r)|^{-\frac{1}{p}} \|f\|_{L_p(B(x, r))}}{\varphi(x, r)}$$

and

$$\mathfrak{M}_{p,\varphi}^W(f; x, r) := \frac{|B(x, r)|^{-\frac{1}{p}} \|f\|_{WL_p(B(x, r))}}{\varphi(x, r)}.$$

In this paper, extending the definition of vanishing Morrey spaces [51], we introduce the generalized vanishing Morrey spaces  $VM_{p,\varphi}(\mathbb{R}^n)$ , including their weak versions and studies the boundedness of the sublinear operators with rough kernel generated by C–Z singular integral operators and their commutators in these spaces. Indeed, we find it convenient to define generalized vanishing Morrey spaces in the form as follows.

**Definition 2.2. (generalized vanishing Morrey space)** The generalized vanishing Morrey space  $VM_{p,\varphi}(\mathbb{R}^n)$  is defined as the spaces of functions  $f \in M_{p,\varphi}(\mathbb{R}^n)$  such that

$$(2.1) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi}(f; x, r) = 0.$$

**Definition 2.3. (weak generalized vanishing Morrey space)** The weak generalized vanishing Morrey space  $WVM_{p,\varphi}(\mathbb{R}^n)$  is defined as the spaces of functions  $f \in VM_{p,\varphi}(\mathbb{R}^n)$  such that

$$(2.2) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi}^W(f; x, r) = 0.$$

Everywhere in the sequel we assume that

$$(2.3) \quad \lim_{r \rightarrow 0} \frac{1}{\inf_{x \in \mathbb{R}^n} \varphi(x, r)} = 0,$$

and

$$(2.4) \quad \sup_{0 < r < \infty} \frac{1}{\inf_{x \in \mathbb{R}^n} \varphi(x, r)} < \infty,$$

which make the spaces  $VM_{p,\varphi}(\mathbb{R}^n)$  and  $WVM_{p,\varphi}(\mathbb{R}^n)$  non-trivial, because bounded functions with compact support belong to this space. The spaces  $VM_{p,\varphi}(\mathbb{R}^n)$  and  $WVM_{p,\varphi}(\mathbb{R}^n)$  are Banach spaces with respect to the norm

$$(2.5) \quad \|f\|_{VM_{p,\varphi}} \equiv \|f\|_{M_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \mathfrak{M}_{p,\varphi}(f; x, r),$$

$$(2.6) \quad \|f\|_{WVM_{p,\varphi}} = \|f\|_{WM_{p,\varphi}} = \sup_{x \in \mathbb{R}^n, r > 0} \mathfrak{M}_{p,\varphi}^W(f; x, r),$$

respectively.

### 3. SUBLINEAR OPERATORS WITH ROUGH KERNEL $T_\Omega$ ON THE SPACES $M_{p,\varphi}$ AND $VM_{p,\varphi}$

In this section, we will first prove the boundedness of the operator  $T_\Omega$  satisfying (1.1) on the generalized Morrey spaces  $M_{p,\varphi}$  by using Lemma 1.1 and the following Lemma 3.1. Then, We will also give the boundedness of  $T_\Omega$  satisfying (1.1) on generalized vanishing Morrey spaces  $VM_{p,\varphi}$ .

**Theorem 3.1.** [11, 33] *Suppose that  $1 \leq p < \infty$ ,  $\Omega \in L_q(S^{n-1})$ ,  $q > 1$ , is homogeneous of degree zero and has mean value zero on  $S^{n-1}$ . If  $q' \leq p$ ,  $p \neq 1$  or  $p < q$ , then the operator  $\bar{T}_\Omega$  is bounded on  $L_p(\mathbb{R}^n)$ . Also the operator  $\bar{T}_\Omega$  is bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$ . Moreover, we have for  $p > 1$*

$$\|\bar{T}_\Omega f\|_{L_p} \leq C \|f\|_{L_p},$$

and for  $p = 1$

$$\|\bar{T}_\Omega f\|_{WL_1} \leq C \|f\|_{L_1}.$$

**Lemma 3.1.** (Our main lemma) *Let  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , be homogeneous of degree zero, and  $1 \leq p < \infty$ . Let  $T_\Omega$  be a sublinear operator satisfying condition (1.1), bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$ .*

*If  $p > 1$  and  $q' \leq p$ , then the inequality*

$$(3.1) \quad \|T_\Omega f\|_{L_p(B(x_0, r))} \lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} t^{-\frac{n}{p}-1} \|f\|_{L_p(B(x_0, t))} dt$$

*holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .*

If  $p > 1$  and  $p < q$ , then the inequality

$$\|T_{\Omega}f\|_{L_p(B(x_0,r))} \lesssim r^{\frac{n}{p}-\frac{n}{q}} \int_{2r}^{\infty} t^{\frac{n}{q}-\frac{n}{p}-1} \|f\|_{L_p(B(x_0,t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .

Moreover, for  $q > 1$  the inequality

$$(3.2) \quad \|T_{\Omega}f\|_{WL_1(B(x_0,r))} \lesssim r^n \int_{2r}^{\infty} t^{-n-1} \|f\|_{L_1(B(x_0,t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_1^{loc}(\mathbb{R}^n)$ .

*Proof.* Let  $1 < p < \infty$  and  $q' \leq p$ . Set  $B = B(x_0, r)$  for the ball centered at  $x_0$  and of radius  $r$  and  $2B = B(x_0, 2r)$ . We represent  $f$  as

$$(3.3) \quad f = f_1 + f_2, \quad f_1(y) = f(y) \chi_{2B}(y), \quad f_2(y) = f(y) \chi_{(2B)^c}(y), \quad r > 0$$

and have

$$\|T_{\Omega}f\|_{L_p(B)} \leq \|T_{\Omega}f_1\|_{L_p(B)} + \|T_{\Omega}f_2\|_{L_p(B)}.$$

Since  $f_1 \in L_p(\mathbb{R}^n)$ ,  $T_{\Omega}f_1 \in L_p(\mathbb{R}^n)$  and from the boundedness of  $T_{\Omega}$  on  $L_p(\mathbb{R}^n)$  (see Theorem 3.1) it follows that:

$$\|T_{\Omega}f_1\|_{L_p(B)} \leq \|T_{\Omega}f_1\|_{L_p(\mathbb{R}^n)} \leq C \|f_1\|_{L_p(\mathbb{R}^n)} = C \|f\|_{L_p(2B)},$$

where constant  $C > 0$  is independent of  $f$ .

It is clear that  $x \in B$ ,  $y \in (2B)^c$  implies  $\frac{1}{2}|x_0 - y| \leq |x - y| \leq \frac{3}{2}|x_0 - y|$ . We get

$$|T_{\Omega}f_2(x)| \leq 2^n c_1 \int_{(2B)^c} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^n} dy.$$

By the Fubini's theorem, we have

$$\begin{aligned} \int_{(2B)^c} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^n} dy &\approx \int_{(2B)^c} |f(y)| |\Omega(x-y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1}} dy \\ &\approx \int_{2r}^{\infty} \int_{2r \leq |x_0-y| \leq t} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1}} \\ &\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1}}. \end{aligned}$$

Applying the Hölder's inequality, we get

$$(3.4) \quad \begin{aligned} &\int_{(2B)^c} \frac{|f(y)| |\Omega(x-y)|}{|x_0 - y|^n} dy \\ &\lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \|\Omega(x-\cdot)\|_{L_q(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{q}} \frac{dt}{t^{n+1}}. \end{aligned}$$

For  $x \in B(x_0, t)$ , notice that  $\Omega$  is homogenous of degree zero and  $\Omega \in L_q(S^{n-1})$ ,  $q > 1$ . Then, we obtain

$$\begin{aligned}
\left( \int_{B(x_0, t)} |\Omega(x-y)|^q dy \right)^{\frac{1}{q}} &= \left( \int_{B(x-x_0, t)} |\Omega(z)|^q dz \right)^{\frac{1}{q}} \\
&\leq \left( \int_{B(0, t+|x-x_0|)} |\Omega(z)|^q dz \right)^{\frac{1}{q}} \\
&\leq \left( \int_{B(0, 2t)} |\Omega(z)|^q dz \right)^{\frac{1}{q}} \\
&= \left( \int_0^{2t} \int_{S^{n-1}} |\Omega(z')|^q d\sigma(z') r^{n-1} dr \right)^{\frac{1}{q}} \\
(3.5) \quad &= C \|\Omega\|_{L_q(S^{n-1})} |B(x_0, 2t)|^{\frac{1}{q}}.
\end{aligned}$$

Thus, by (3.5), it follows that:

$$|T_\Omega f_2(x)| \lesssim \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{p}+1}}.$$

Moreover, for all  $p \in [1, \infty)$  the inequality

$$(3.6) \quad \|T_\Omega f_2\|_{L_p(B)} \lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{p}+1}}$$

holds. Thus

$$\|T_\Omega f\|_{L_p(B)} \lesssim \|f\|_{L_p(2B)} + r^{\frac{n}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{p}+1}}.$$

On the other hand, we have

$$\begin{aligned}
\|f\|_{L_p(2B)} &\approx r^{\frac{n}{p}} \|f\|_{L_p(B)} \int_{2r}^{\infty} \frac{dt}{t^{\frac{n}{p}+1}} \\
(3.7) \quad &\leq r^{\frac{n}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{p}+1}}.
\end{aligned}$$

By combining the above inequalities, we obtain

$$\|T_\Omega f\|_{L_p(B)} \lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0, t))} \frac{dt}{t^{\frac{n}{p}+1}}.$$

Let  $1 < p < q$ . Similarly to (3.5), when  $y \in B(x_0, t)$ , notice that

$$(3.8) \quad \left( \int_{B(x_0, r)} |\Omega(x-y)|^q dy \right)^{\frac{1}{q}} \leq C \|\Omega\|_{L_q(S^{n-1})} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{q}}.$$

By the Fubini's theorem, the Minkowski inequality and (3.8), we get

$$\begin{aligned} \|T_\Omega f_2\|_{L_p(B)} &\leq \left( \int_B \left| \int_{2r}^\infty \int_{B(x_0, t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1}} \right|^p dx \right)^{\frac{1}{p}} \\ &\leq \int_{2r}^\infty \int_{B(x_0, t)} |f(y)| \|\Omega(\cdot - y)\|_{L_p(B)} dy \frac{dt}{t^{n+1}} \\ &\leq |B(x_0, r)|^{\frac{1}{p} - \frac{1}{q}} \int_{2r}^\infty \int_{B(x_0, t)} |f(y)| \|\Omega(\cdot - y)\|_{L_q(B)} dy \frac{dt}{t^{n+1}} \\ &\lesssim r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^\infty \|f\|_{L_1(B(x_0, t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{q}} \frac{dt}{t^{n+1}} \\ &\lesssim r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^\infty \|f\|_{L_p(B(x_0, t))} t^{\frac{n}{q} - \frac{n}{p} - 1} dt. \end{aligned}$$

Let  $p = 1 < q \leq \infty$ . From the weak (1, 1) boundedness of  $T_\Omega$  and (3.7) it follows that:

$$(3.9) \quad \begin{aligned} \|T_\Omega f_1\|_{WL_1(B)} &\leq \|T_\Omega f_1\|_{WL_1(\mathbb{R}^n)} \lesssim \|f_1\|_{L_1(\mathbb{R}^n)} \\ &= \|f\|_{L_1(2B)} \lesssim r^n \int_{2r}^\infty \|f\|_{L_1(B(x_0, t))} \frac{dt}{t^{n+1}}. \end{aligned}$$

Then from (3.6) and (3.9) we get the inequality (3.2), which completes the proof.  $\square$

In the following theorem, we get the boundedness of the operator  $T_\Omega$  on the generalized Morrey spaces  $M_{p, \varphi}$ .

**Theorem 3.2.** (Our main result) Let  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , be homogeneous of degree zero, and  $1 \leq p < \infty$ . Let  $T_\Omega$  be a sublinear operator satisfying condition (1.1), bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$ . Let also, for  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(3.10) \quad \int_r^\infty \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{p} + 1}} dt \leq C \varphi_2(x, r),$$

and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(3.11) \quad \int_r^\infty \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{p} - \frac{n}{q} + 1}} dt \leq C \varphi_2(x, r) r^{\frac{n}{q}},$$

where  $C$  does not depend on  $x$  and  $r$ .

Then the operator  $T_\Omega$  is bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{1,\varphi_2}$ . Moreover, we have for  $p > 1$

$$(3.12) \quad \|T_\Omega f\|_{M_{p,\varphi_2}} \lesssim \|f\|_{M_{p,\varphi_1}},$$

and for  $p = 1$

$$(3.13) \quad \|T_\Omega f\|_{WM_{1,\varphi_2}} \lesssim \|f\|_{M_{1,\varphi_1}}.$$

*Proof.* Since  $f \in M_{p,\varphi_1}$ , by (2.6) and the non-decreasing, with respect to  $t$ , of the norm  $\|f\|_{L_p(B(x_0,t))}$ , we get

$$\begin{aligned} & \frac{\|f\|_{L_p(B(x_0,t))}}{\operatorname{ess\,inf}_{0 < t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \operatorname{ess\,sup}_{0 < t < \tau < \infty} \frac{\|f\|_{L_p(B(x_0,t))}}{\varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \operatorname{ess\,sup}_{0 < \tau < \infty} \frac{\|f\|_{L_p(B(x_0,\tau))}}{\varphi_1(x_0, \tau) \tau^{\frac{n}{p}}} \\ & \leq \|f\|_{M_{p,\varphi_1}}. \end{aligned}$$

For  $q' \leq p < \infty$ , since  $(\varphi_1, \varphi_2)$  satisfies (3.10), we have

$$\begin{aligned} & \int_r^\infty \|f\|_{L_p(B(x_0,t))} t^{-\frac{n}{p}} \frac{dt}{t} \\ & \leq \int_r^\infty \frac{\|f\|_{L_p(B(x_0,t))} \operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}}{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}} t^{\frac{n}{p}}} \frac{dt}{t} \\ & \leq C \|f\|_{M_{p,\varphi_1}} \int_r^\infty \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x_0, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{p}}} \frac{dt}{t} \\ & \leq C \|f\|_{M_{p,\varphi_1}} \varphi_2(x_0, r). \end{aligned}$$

Then by (3.1), we get

$$\begin{aligned} \|T_\Omega f\|_{M_{p,\varphi_2}} &= \sup_{x_0 \in \mathbb{R}^n, r > 0} \varphi_2(x_0, r)^{-1} |B(x_0, r)|^{-\frac{1}{p}} \|T_\Omega f\|_{L_p(B(x_0, r))} \\ &\leq C \sup_{x_0 \in \mathbb{R}^n, r > 0} \varphi_2(x_0, r)^{-1} \int_r^\infty \|f\|_{L_p(B(x_0, t))} t^{-\frac{n}{p}} \frac{dt}{t} \\ &\leq C \|f\|_{M_{p,\varphi_1}}. \end{aligned}$$

For the case of  $1 \leq p < q$ , we can also use the same method, so we omit the details. This completes the proof of Theorem 3.2.  $\square$

In the case of  $q = \infty$  by Theorem 3.2, we get

**Corollary 3.1.** *Let  $1 \leq p < \infty$  and the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10). Then the operators  $M$  and  $\bar{T}$  are bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{1,\varphi_2}$ .*

Let  $f \in L_1^{loc}(\mathbb{R}^n)$ . The rough H–L maximal operator  $M_\Omega$  is defined by

$$M_\Omega f(x) = \sup_{t>0} \frac{1}{|B(x,t)|} \int_{B(x,t)} |\Omega(x-y)| |f(y)| dy.$$

Then we can get the following corollary.

**Corollary 3.2.** *Let  $1 \leq p < \infty$ ,  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ . For  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10) and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.11). Then the operators  $M_\Omega$  and  $\bar{T}_\Omega$  are bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{1,\varphi_2}$ .*

Now using above results, we get the boundedness of the operator  $T_\Omega$  on the vanishing generalized Morrey spaces  $VM_{p,\varphi}$ .

**Theorem 3.3.** *(Our main result) Let  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , be homogeneous of degree zero, and  $1 \leq p < \infty$ . Let  $T_\Omega$  be a sublinear operator satisfying condition (1.1), bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$ . Let for  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and*

$$(3.14) \quad c_\delta := \int_\delta^\infty \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}+1}} dt < \infty$$

for every  $\delta > 0$ , and

$$(3.15) \quad \int_r^\infty \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}+1}} dt \leq C_0 \varphi_2(x, r),$$

and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and also

$$(3.16) \quad c_{\delta'} := \int_{\delta'}^\infty \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}-\frac{n}{q}+1}} dt < \infty$$

for every  $\delta' > 0$ , and

$$(3.17) \quad \int_r^\infty \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}-\frac{n}{q}+1}} dt \leq C_0 \varphi_2(x, r) r^{\frac{n}{q}},$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$ .

Then the operator  $T_\Omega$  is bounded from  $VM_{p,\varphi_1}$  to  $VM_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WVM_{1,\varphi_2}$ . Moreover, we have for  $p > 1$

$$(3.18) \quad \|T_\Omega f\|_{VM_{p,\varphi_2}} \lesssim \|f\|_{VM_{p,\varphi_1}},$$

and for  $p = 1$

$$(3.19) \quad \|T_\Omega f\|_{WVM_{1,\varphi_2}} \lesssim \|f\|_{VM_{1,\varphi_1}}.$$

*Proof.* The norm inequalities follow from Theorem 3.2. Thus we only have to prove that

$$(3.20) \quad \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi_1}(f; x, r) = 0 \text{ implies } \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p,\varphi_2}(T_\Omega f; x, r) = 0$$

and

$$(3.21) \quad \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p, \varphi_1}(f; x, r) = 0 \text{ implies } \lim_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \mathfrak{M}_{p, \varphi_2}^W(T_\Omega f; x, r) = 0.$$

To show that  $\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|T_\Omega f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} < \epsilon$  for small  $r$ , we split the right-hand side of (3.1):

$$(3.22) \quad \frac{r^{-\frac{n}{p}} \|T_\Omega f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} \leq C [I_{\delta_0}(x, r) + J_{\delta_0}(x, r)],$$

where  $\delta_0 > 0$  (we may take  $\delta_0 < 1$ ), and

$$I_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_r^{\delta_0} t^{-\frac{n}{p}-1} \|f\|_{L_p(B(x, t))} dt,$$

and

$$J_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_{\delta_0}^{\infty} t^{-\frac{n}{p}-1} \|f\|_{L_p(B(x, t))} dt$$

and  $r < \delta_0$ . Now we choose any fixed  $\delta_0 > 0$  such that

$$\sup_{x \in \mathbb{R}^n} \frac{t^{-\frac{n}{p}} \|f\|_{L_p(B(x, t))}}{\varphi_1(x, t)} < \frac{\epsilon}{2CC_0}, \quad t \leq \delta_0,$$

where  $C$  and  $C_0$  are constants from (3.15) and (3.22). This allows to estimate the first term uniformly in  $r \in (0, \delta_0)$ :

$$\sup_{x \in \mathbb{R}^n} CI_{\delta_0}(x, r) < \frac{\epsilon}{2}, \quad 0 < r < \delta_0.$$

The estimation of the second term may be obtained by choosing  $r$  sufficiently small. Indeed, we have

$$J_{\delta_0}(x, r) \leq c_{\delta_0} \frac{\|f\|_{M_{p, \varphi_1}}}{\varphi_2(x, r)},$$

where  $c_{\delta_0}$  is the constant from (3.14) with  $\delta = \delta_0$ . Then, by (2.3) it suffices to choose  $r$  small enough such that

$$\sup_{x \in \mathbb{R}^n} \frac{1}{\varphi_2(x, r)} \leq \frac{\epsilon}{2c_{\delta_0} \|f\|_{M_{p, \varphi_1}}},$$

which completes the proof of (3.20).

The proof of (3.21) is similar to the proof of (3.20). For the case of  $1 \leq p < q$ , we can also use the same method, so we omit the details.  $\square$

*Remark 3.1.* Conditions (3.14) and (3.16) are not needed in the case when  $\varphi(x, r)$  does not depend on  $x$ , since (3.14) follows from (3.15) and similarly, (3.16) follows from (3.17) in this case.

**Corollary 3.3.** *Let  $1 \leq p < \infty$ ,  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ . For  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.14)-(3.15) and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.16)-(3.17). Then the operators  $M_\Omega$  and  $\bar{T}_\Omega$  are bounded from  $VM_{p, \varphi_1}$  to  $VM_{p, \varphi_2}$  for  $p > 1$  and from  $VM_{1, \varphi_1}$  to  $WVM_{1, \varphi_2}$ .*



In the case of  $q = \infty$  by Theorem 3.3, we get

**Corollary 3.4.** *Let  $1 \leq p < \infty$  and the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and (3.14)-(3.15). Then the operators  $M$  and  $\bar{T}$  are bounded from  $VM_{p,\varphi_1}$  to  $VM_{p,\varphi_2}$  for  $p > 1$  and from  $VM_{1,\varphi_1}$  to  $WVM_{1,\varphi_2}$ .*

#### 4. COMMUTATORS OF THE SUBLINEAR OPERATORS WITH ROUGH KERNEL $T_\Omega$ ON THE SPACES $M_{p,\varphi}$ AND $VM_{p,\varphi}$

In this section, we will first prove the boundedness of the operator  $T_{\Omega,b}$  satisfying (1.2) with  $b \in BMO(\mathbb{R}^n)$  on the generalized Morrey spaces  $M_{p,\varphi}$  by using Lemma 1.1 and the following Lemma 4.1. Then, we will also obtain the boundedness of  $T_{\Omega,b}$  satisfying (1.2) with  $b \in BMO(\mathbb{R}^n)$  on generalized vanishing Morrey spaces  $VM_{p,\varphi}$ .

Let  $T$  be a linear operator. For a locally integrable function  $b$  on  $\mathbb{R}^n$ , we define the commutator  $[b, T]$  by

$$[b, T]f(x) = b(x)Tf(x) - T(bf)(x)$$

for any suitable function  $f$ .

The function  $b$  is also called the symbol function of  $[b, T]$ . The investigation of the operator  $[b, T]$  begins with Coifman-Rocherberg-Weiss pioneering study of the operator  $T$  (see [8]). Let  $\bar{T}$  be a C-Z operator. A well known result of Coifman et al. [8] states that when  $K(x) = \frac{\Omega(x')}{|x|^n}$  and  $\Omega$  is smooth, the commutator  $[b, \bar{T}]f = b\bar{T}f - \bar{T}(bf)$  is bounded on  $L_p(\mathbb{R}^n)$ ,  $1 < p < \infty$ , if and only if  $b \in BMO(\mathbb{R}^n)$ . There are two major reasons for considering the problem of commutators. The first one is that the boundedness of commutators can produce some characterizations of function spaces (see [2, 18, 19, 21, 36, 44]). The other one is that the theory of commutators plays an important role in the study of the regularity of solutions to elliptic and parabolic PDEs of the second order (see [6, 7, 12, 13]).

Many authors are interested in the study of commutators for which the symbol functions belong to  $BMO(\mathbb{R}^n)$  spaces (see [17, 21, 23, 28, 36] for example). The boundedness of the commutator has also been generalized to other contexts and important applications to some non-linear PDEs have been given by Coifman et al. [9].

Let us recall the definition of the space of  $BMO(\mathbb{R}^n)$  (bounded mean oscillation).

**Definition 4.1.** Suppose that  $b \in L_1^{loc}(\mathbb{R}^n)$ , let

$$\|b\|_* = \sup_{x \in \mathbb{R}^n, r > 0} \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}| dy < \infty,$$

where

$$b_{B(x, r)} = \frac{1}{|B(x, r)|} \int_{B(x, r)} b(y) dy.$$

Define

$$BMO(\mathbb{R}^n) = \{b \in L_1^{loc}(\mathbb{R}^n) : \|b\|_* < \infty\}.$$

If one regards two functions whose difference is a constant as one, then the space  $BMO(\mathbb{R}^n)$  is a Banach space with respect to norm  $\|\cdot\|_*$ .

*Remark 4.1.* [23] (1) The John-Nirenberg inequality [22]: there are constants  $C_1, C_2 > 0$ , such that for all  $b \in BMO(\mathbb{R}^n)$  and  $\beta > 0$

$$|\{x \in B : |b(x) - b_B| > \beta\}| \leq C_1 |B| e^{-C_2 \beta / \|b\|_*}, \quad \forall B \subset \mathbb{R}^n.$$

(2) The John-Nirenberg inequality implies that

$$(4.1) \quad \|b\|_* \approx \sup_{x \in \mathbb{R}^n, r > 0} \left( \frac{1}{|B(x, r)|} \int_{B(x, r)} |b(y) - b_{B(x, r)}|^p dy \right)^{\frac{1}{p}}$$

for  $1 < p < \infty$ .

(3) Let  $b \in BMO(\mathbb{R}^n)$ . Then there is a constant  $C > 0$  such that

$$(4.2) \quad |b_{B(x, r)} - b_{B(x, t)}| \leq C \|b\|_* \ln \frac{t}{r} \text{ for } 0 < 2r < t,$$

where  $C$  is independent of  $b, x, r$  and  $t$ .

**Theorem 4.1.** [11, 27] *Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $q > 1$ , is homogeneous of degree zero and has mean value zero on  $S^{n-1}$ . Let  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . If  $q' \leq p$  or  $p < q$ , then the commutator operator  $[b, \overline{T}_\Omega]$  is bounded on  $L_p(\mathbb{R}^n)$ .*

**Lemma 4.1.** (Our main lemma) *Let  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , be homogeneous of degree zero. Let  $1 < p < \infty$ ,  $b \in BMO(\mathbb{R}^n)$  and  $T_{\Omega, b}$  is a sublinear operator satisfying condition (1.2), bounded on  $L_p(\mathbb{R}^n)$ . Then, for  $q' \leq p$  the inequality*

$$(4.3) \quad \|T_{\Omega, b} f\|_{L_p(B(x_0, r))} \lesssim \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{p}-1} \|f\|_{L_p(B(x_0, t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .

Also, for  $p < q$  the inequality

$$\|T_{\Omega, b} f\|_{L_p(B(x_0, r))} \lesssim \|b\|_* r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{q} - \frac{n}{p} - 1} \|f\|_{L_p(B(x_0, t))} dt$$

holds for any ball  $B(x_0, r)$  and for all  $f \in L_p^{loc}(\mathbb{R}^n)$ .

*Proof.* Let  $1 < p < \infty$ . As in the proof of Lemma 3.1, we represent  $f$  in form (3.3) and have

$$\|T_{\Omega, b} f\|_{L_p(B)} \leq \|T_{\Omega, b} f_1\|_{L_p(B)} + \|T_{\Omega, b} f_2\|_{L_p(B)}.$$

From the boundedness of  $T_{\Omega, b}$  on  $L_p(\mathbb{R}^n)$  (see Theorem 4.1) it follows that:

$$\begin{aligned} \|T_{\Omega, b} f_1\|_{L_p(B)} &\leq \|T_{\Omega, b} f_1\|_{L_p(\mathbb{R}^n)} \\ &\lesssim \|b\|_* \|f_1\|_{L_p(\mathbb{R}^n)} = \|b\|_* \|f\|_{L_p(2B)}. \end{aligned}$$

It is known that  $x \in B, y \in (2B)^C$ , which implies  $\frac{1}{2}|x_0 - y| \leq |x - y| \leq \frac{3}{2}|x_0 - y|$ . Then for  $x \in B$ , we have

$$\begin{aligned} |T_{\Omega, b} f_2(x)| &\lesssim \int_{\mathbb{R}^n} \frac{|\Omega(x - y)|}{|x - y|^n} |b(y) - b(x)| |f(y)| dy \\ &\approx \int_{(2B)^C} \frac{|\Omega(x - y)|}{|x_0 - y|^n} |b(y) - b(x)| |f(y)| dy. \end{aligned}$$

Hence we get

$$\begin{aligned}
\|T_{\Omega,b}f_2\|_{L_p(B)} &\lesssim \left( \int_B \left( \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^n} |b(y)-b(x)| |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\
&\lesssim \left( \int_B \left( \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^n} |b(y)-b_B| |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\
&\quad + \left( \int_B \left( \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^n} |b(x)-b_B| |f(y)| dy \right)^p dx \right)^{\frac{1}{p}} \\
&= J_1 + J_2.
\end{aligned}$$

We have the following estimation of  $J_1$ . When  $s' \leq p$  and  $\frac{1}{\mu} + \frac{1}{p} + \frac{1}{q} = 1$ , by the Fubini's theorem

$$\begin{aligned}
J_1 &\approx r^{\frac{n}{p}} \int_{(2B)^c} \frac{|\Omega(x-y)|}{|x_0-y|^n} |b(y)-b_B| |f(y)| dy \\
&\approx r^{\frac{n}{p}} \int_{(2B)^c} |\Omega(x-y)| |b(y)-b_B| |f(y)| \int_{|x_0-y|}^{\infty} \frac{dt}{t^{n+1}} dy \\
&\approx r^{\frac{n}{p}} \int_{2r}^{\infty} \int_{2r < |x_0-y| < t} |\Omega(x-y)| |b(y)-b_B| |f(y)| dy \frac{dt}{t^{n+1}} \\
&\lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} \int_{B(x_0,t)} |\Omega(x-y)| |b(y)-b_B| |f(y)| dy \frac{dt}{t^{n+1}} \text{ holds.}
\end{aligned}$$

Applying the Hölder's inequality and by (3.8), (4.1), (4.2), we get

$$\begin{aligned}
J_1 &\lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} \int_{B(x_0,t)} |\Omega(x-y)| |b(y) - b_{B(x_0,t)}| |f(y)| dy \frac{dt}{t^{n+1}} \\
&+ r^{\frac{n}{p}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |\Omega(x-y)| |f(y)| dy \frac{dt}{t^{n+1}} \\
&\lesssim r^{\frac{n}{p}} \int_{2r}^{\infty} \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} \| (b(\cdot) - b_{B(x_0,t)}) \|_{L_\mu(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{n+1}} \\
&+ r^{\frac{n}{p}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} |B(x_0,t)|^{1-\frac{1}{p}-\frac{1}{q}} \frac{dt}{t^{n+1}} \\
&\lesssim \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{p}+1}}.
\end{aligned}$$

In order to estimate  $J_2$  note that

$$J_2 = \| (b(\cdot) - b_{B(x_0,t)}) \|_{L_p(B(x_0,t))} \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^n} |f(y)| dy.$$

By (4.1), we get

$$J_2 \lesssim \|b\|_* r^{\frac{n}{p}} \int_{(2B)^C} \frac{|\Omega(x-y)|}{|x_0-y|^n} |f(y)| dy.$$

Thus, by (3.4) and (3.5)

$$J_2 \lesssim \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{p}+1}}.$$

Summing up  $J_1$  and  $J_2$ , for all  $p \in (1, \infty)$  we get

$$(4.4) \quad \|T_{\Omega,b}f\|_{L_p(B)} \lesssim \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{p}+1}}.$$

Finally, we have the following

$$\|T_{\Omega,b}f\|_{L_p(B)} \lesssim \|b\|_* \|f\|_{L_p(2B)} + \|b\|_* r^{\frac{n}{p}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) \|f\|_{L_p(B(x_0,t))} \frac{dt}{t^{\frac{n}{p}+1}},$$

which completes the proof of first statement by (3.7).

On the other hand when  $p < q$ , by the Fubini's theorem and the Minkowski inequality, we get

$$\begin{aligned}
J_1 &\lesssim \left( \int_B \left| \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1}} \right|^p dx \right)^{\frac{1}{p}} \\
&+ \left( \int_B \left| \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| |\Omega(x-y)| dy \frac{dt}{t^{n+1}} \right|^p dx \right)^{\frac{1}{p}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| \|\Omega(\cdot - y)\|_{L_p(B(x_0,t))} dy \frac{dt}{t^{n+1}} \\
&+ \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_p(B(x_0,t))} dy \frac{dt}{t^{n+1}} \\
&\lesssim |B|^{\frac{1}{p} - \frac{1}{q}} \int_{2r}^{\infty} \int_{B(x_0,t)} |b(y) - b_{B(x_0,t)}| |f(y)| \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} dy \frac{dt}{t^{n+1}} \\
&+ |B|^{\frac{1}{p} - \frac{1}{q}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_q(B(x_0,t))} dy \frac{dt}{t^{n+1}}.
\end{aligned}$$

Applying the Hölder's inequality and by (3.8), (4.1), (4.2), we get

$$\begin{aligned}
J_1 &\lesssim r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \|(b(\cdot) - b_{B(x_0,t)}) f\|_{L_1(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{q}} \frac{dt}{t^{n+1}} \\
&+ r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|f\|_{L_p(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{q}} \frac{dt}{t^{\frac{n}{p}+1}} \\
&\lesssim r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \|(b(\cdot) - b_{B(x_0,t)})\|_{L_{p'}(B(x_0,t))} \|f\|_{L_p(B(x_0,t))} t^{\frac{n}{q}} \frac{dt}{t^{n+1}} \\
&+ r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} |b_{B(x_0,r)} - b_{B(x_0,t)}| \|f\|_{L_p(B(x_0,t))} t^{\frac{n}{q}} \frac{dt}{t^{\frac{n}{p}+1}} \\
&\lesssim \|b\|_* r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{q} - \frac{n}{p} - 1} \|f\|_{L_p(B(x_0,t))} dt.
\end{aligned}$$

Let  $\frac{1}{p} = \frac{1}{\nu} + \frac{1}{q}$ , then for  $J_2$ , by the Fubini's theorem, the Minkowski inequality, the Hölder's inequality and from (3.8), we get

$$\begin{aligned}
J_2 &\lesssim \left( \int_B \left| \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| |b(x) - b_B| |\Omega(x-y)| dy \frac{dt}{t^{n+1}} \right|^p dx \right)^{\frac{1}{p}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|(b(\cdot) - b_B)\Omega(\cdot - y)\|_{L_p(B)} dy \frac{dt}{t^{n+1}} \\
&\lesssim \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|b(\cdot) - b_B\|_{L_\nu(B)} \|\Omega(\cdot - y)\|_{L_q(B)} dy \frac{dt}{t^{n+1}} \\
&\lesssim \|b\|_* |B|^{\frac{1}{p} - \frac{1}{q}} \int_{2r}^{\infty} \int_{B(x_0,t)} |f(y)| \|\Omega(\cdot - y)\|_{L_q(B)} dy \frac{dt}{t^{n+1}} \\
&\lesssim \|b\|_* r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \|f\|_{L_1(B(x_0,t))} \left| B\left(x_0, \frac{3}{2}t\right) \right|^{\frac{1}{q}} \frac{dt}{t^{n+1}} \\
&\lesssim \|b\|_* r^{\frac{n}{p} - \frac{n}{q}} \int_{2r}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{\frac{n}{q} - \frac{n}{p} - 1} \|f\|_{L_p(B(x_0,t))} dt.
\end{aligned}$$

By combining the above estimates, we complete the proof of Lemma 4.1.  $\square$

Now we can give the following theorem (our main result).

**Theorem 4.2.** (Our main result) Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , is homogeneous of degree zero and  $T_{\Omega,b}$  is a sublinear operator satisfying condition (1.2), bounded on  $L_p(\mathbb{R}^n)$ . Let  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . Let also, for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(4.5) \quad \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{p} + 1}} dt \leq C \varphi_2(x, r),$$

and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies the condition

$$(4.6) \quad \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) \frac{\operatorname{ess\,inf}_{t < \tau < \infty} \varphi_1(x, \tau) \tau^{\frac{n}{p}}}{t^{\frac{n}{p} - \frac{n}{q} + 1}} dt \leq C \varphi_2(x, r) r^{\frac{n}{q}},$$

where  $C$  does not depend on  $x$  and  $r$ .

Then, the operator  $T_{\Omega,b}$  is bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$ . Moreover,

$$\|T_{\Omega,b}f\|_{M_{p,\varphi_2}} \lesssim \|b\|_* \|f\|_{M_{p,\varphi_1}}.$$

*Proof.* The statement of Theorem 4.2 follows by Lemma 1.1 and Lemma 4.1 in the same manner as in the proof of Theorem 3.2.  $\square$

For the sublinear commutator of the fractional maximal operator with rough kernel which is defined as follows

$$M_{\Omega,b}(f)(x) = \sup_{t>0} |B(x,t)|^{-1} \int_{B(x,t)} |b(x) - b(y)| |\Omega(x-y)| |f(y)| dy$$

and for the linear commutator of the singular integral  $[b, \overline{T}_\Omega]$  by Theorem 4.2 we get the following new results.

**Corollary 4.1.** *Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , is homogeneous of degree zero,  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . If for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.5) and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.6). Then, the operators  $M_{\Omega,b}$  and  $[b, \overline{T}_\Omega]$  are bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$ .*

We get the following new results for the sublinear commutator of the maximal operator

$$M_b(f)(x) = \sup_{t>0} |B(x,t)|^{-1} \int_{B(x,t)} |b(x) - b(y)| |f(y)| dy$$

and for the linear commutator of the singular integral  $[b, \overline{T}]$  by Theorem 4.2.

**Corollary 4.2.** *Let  $1 < p < \infty$ ,  $b \in BMO(\mathbb{R}^n)$  and the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.5). Then, the operators  $M_b$  and  $[b, \overline{T}]$  are bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$ .*

Now using above results, we also obtain the boundedness of the operator  $T_{\Omega,b}$  on the vanishing generalized Morrey spaces  $VM_{p,\varphi}$ .

**Theorem 4.3.** *(Our main result) Let  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , be homogeneous of degree zero. Let  $T_{\Omega,b}$  is a sublinear operator satisfying condition (1.2) bounded on  $L_p(\mathbb{R}^n)$ . Let  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . Let for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and*

$$(4.7) \quad \int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x,t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}+1}} dt \leq C_0 \varphi_2(x,r),$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$ ,

$$(4.8) \quad \lim_{r \rightarrow 0} \frac{\ln \frac{1}{r}}{\inf_{x \in \mathbb{R}^n} \varphi_2(x,r)} = 0$$

and

$$(4.9) \quad c_\delta := \int_\delta^\infty (1 + \ln |t|) \sup_{x \in \mathbb{R}^n} \varphi_1(x,t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}+1}} dt < \infty$$

for every  $\delta > 0$ , and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4) and also

$$(4.10) \quad \int_r^\infty \left(1 + \ln \frac{t}{r}\right) \varphi_1(x,t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p}-\frac{n}{q}+1}} dt \leq C_0 \varphi_2(x,r) r^{\frac{n}{q}},$$

where  $C_0$  does not depend on  $x \in \mathbb{R}^n$  and  $r > 0$

$$\lim_{r \rightarrow 0} \frac{\ln \frac{1}{r}}{\inf_{x \in \mathbb{R}^n} \varphi_2(x,r)} = 0$$

and

$$(4.11) \quad c_{\delta'} := \int_{\delta'}^{\infty} (1 + \ln |t|) \sup_{x \in \mathbb{R}^n} \varphi_1(x, t) \frac{t^{\frac{n}{p}}}{t^{\frac{n}{p} - \frac{n}{q} + 1}} dt < \infty$$

for every  $\delta' > 0$ .

Then the operator  $T_{\Omega, b}$  is bounded from  $VM_{p, \varphi_1}$  to  $VM_{p, \varphi_2}$ . Moreover,

$$(4.12) \quad \|T_{\Omega, b} f\|_{VM_{p, \varphi_2}} \lesssim \|b\|_* \|f\|_{VM_{p, \varphi_1}}.$$

*Proof.* The norm inequality having already been provided by Theorem 4.2, we only have to prove the implication

$$(4.13) \quad \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|f\|_{L_p(B(x, r))}}{\varphi_1(x, r)} = 0 \text{ implies } \limsup_{r \rightarrow 0} \sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|T_{\Omega, b} f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} = 0.$$

To show that

$$\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|T_{\Omega, b} f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} < \epsilon \text{ for small } r,$$

we use the estimate (4.3):

$$\sup_{x \in \mathbb{R}^n} \frac{r^{-\frac{n}{p}} \|T_{\Omega, b} f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} \lesssim \frac{\|b\|_*}{\varphi_2(x, r)} \int_r^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{p} - 1} \|f\|_{L_p(B(x, t))} dt.$$

We take  $r < \delta_0$ , where  $\delta_0$  will be chosen small enough and split the integration:

$$(4.14) \quad \frac{r^{-\frac{n}{p}} \|T_{\Omega, b} f\|_{L_p(B(x, r))}}{\varphi_2(x, r)} \leq C [I_{\delta_0}(x, r) + J_{\delta_0}(x, r)],$$

where  $\delta_0 > 0$  (we may take  $\delta_0 < 1$ ), and

$$I_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_r^{\delta_0} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{p} - 1} \|f\|_{L_p(B(x, t))} dt,$$

and

$$J_{\delta_0}(x, r) := \frac{1}{\varphi_2(x, r)} \int_{\delta_0}^{\infty} \left(1 + \ln \frac{t}{r}\right) t^{-\frac{n}{p} - 1} \|f\|_{L_p(B(x, t))} dt$$

Now we choose any fixed  $\delta_0 > 0$  such that

$$\sup_{x \in \mathbb{R}^n} \frac{t^{-\frac{n}{p}} \|f\|_{L_p(B(x, t))}}{\varphi_1(x, t)} < \frac{\epsilon}{2CC_0}, \quad t \leq \delta_0,$$

where  $C$  and  $C_0$  are constants from (4.7) and (4.14). This allows to estimate the first term uniformly in  $r \in (0, \delta_0)$ :

$$\sup_{x \in \mathbb{R}^n} CI_{\delta_0}(x, r) < \frac{\epsilon}{2}, \quad 0 < r < \delta_0.$$

For the second term, writing  $1 + \ln \frac{t}{r} \leq 1 + |\ln t| + \ln \frac{1}{r}$ , we obtain

$$J_{\delta_0}(x, r) \leq \frac{c_{\delta_0} + \widetilde{c}_{\delta_0} \ln \frac{1}{r}}{\varphi_2(x, r)} \|f\|_{M_{p, \varphi}},$$



where  $c_{\delta_0}$  is the constant from (4.9) with  $\delta = \delta_0$  and  $\widetilde{c_{\delta_0}}$  is a similar constant with omitted logarithmic factor in the integrand. Then, by (4.8) we can choose small enough  $r$  such that

$$\sup_{x \in \mathbb{R}^n} J_{\delta_0}(x, r) < \frac{\epsilon}{2},$$

which completes the proof of (4.13).

For the case of  $p < q$ , we can also use the same method, so we omit the details.  $\square$

*Remark 4.2.* Conditions (4.9) and (4.11) are not needed in the case when  $\varphi(x, r)$  does not depend on  $x$ , since (4.9) follows from (4.7) and similarly, (4.11) follows from (4.10) in this case.

**Corollary 4.3.** *Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , is homogeneous of degree zero,  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . If for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.9)-(4.7) and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.11)-(4.10). Then, the operators  $M_{\Omega, b}$  and  $[b, \overline{T}_{\Omega}]$  are bounded from  $VM_{p, \varphi_1}$  to  $VM_{p, \varphi_2}$ .*

In the case of  $q = \infty$  by Theorem 4.3, we get

**Corollary 4.4.** *Let  $1 < p < \infty$ ,  $b \in BMO(\mathbb{R}^n)$  and the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)-(2.4)-(4.8) and (4.9)-(4.7). Then the operators  $M_b$  and  $[b, \overline{T}]$  are bounded from  $VM_{p, \varphi_1}$  to  $VM_{p, \varphi_2}$ .*

### 5. SOME APPLICATIONS

In this section, we give the applications of Theorem 3.2, Theorem 3.3, Theorem 4.2, Theorem 4.3 for the Marcinkiewicz operator.

**5.1. Marcinkiewicz Operator.** Let  $S^{n-1} = \{x \in \mathbb{R}^n : |x| = 1\}$  be the unit sphere in  $\mathbb{R}^n$  equipped with the Lebesgue measure  $d\sigma$ . Suppose that  $\Omega$  satisfies the following conditions.

(a)  $\Omega$  is the homogeneous function of degree zero on  $\mathbb{R}^n \setminus \{0\}$ , that is,

$$\Omega(\mu x) = \Omega(x), \text{ for any } \mu > 0, x \in \mathbb{R}^n \setminus \{0\}.$$

(b)  $\Omega$  has mean zero on  $S^{n-1}$ , that is,

$$\int_{S^{n-1}} \Omega(x') d\sigma(x') = 0,$$

where  $x' = \frac{x}{|x|}$  for any  $x \neq 0$ .

(c)  $\Omega \in Lip_{\gamma}(S^{n-1})$ ,  $0 < \gamma \leq 1$ , that is there exists a constant  $M > 0$  such that,

$$|\Omega(x') - \Omega(y')| \leq M|x' - y'|^{\gamma} \text{ for any } x', y' \in S^{n-1}.$$

In 1958, Stein [46] defined the Marcinkiewicz integral of higher dimension  $\mu_{\Omega}$  as

$$\mu_{\Omega}(f)(x) = \left( \int_0^{\infty} |F_{\Omega, t}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega, t}(f)(x) = \int_{|x-y| \leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} f(y) dy.$$

Since Stein's work in 1958, the continuity of Marcinkiewicz integral has been extensively studied as a research topic and also provides useful tools in harmonic analysis [27, 47, 48, 49].

The sublinear commutator of the operator  $\mu_\Omega$  is defined by

$$[b, \mu_\Omega](f)(x) = \left( \int_0^\infty |F_{\Omega,t,b}(f)(x)|^2 \frac{dt}{t^3} \right)^{1/2},$$

where

$$F_{\Omega,t,b}(f)(x) = \int_{|x-y|\leq t} \frac{\Omega(x-y)}{|x-y|^{n-1}} [b(x) - b(y)] f(y) dy.$$

We consider the space  $H = \{h : \|h\| = (\int_0^\infty |h(t)|^2 \frac{dt}{t^3})^{1/2} < \infty\}$ . Then, it is clear that  $\mu_\Omega(f)(x) = \|F_{\Omega,t}(x)\|$ .

By the Minkowski inequality and the conditions on  $\Omega$ , we get

$$\mu_\Omega(f)(x) \leq \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^{n-1}} |f(y)| \left( \int_{|x-y|}^\infty \frac{dt}{t^3} \right)^{1/2} dy \leq C \int_{\mathbb{R}^n} \frac{|\Omega(x-y)|}{|x-y|^n} |f(y)| dy.$$

Thus,  $\mu_\Omega$  satisfies the condition (1.1). It is known that  $\mu_\Omega$  is bounded on  $L_p(\mathbb{R}^n)$  for  $p > 1$ , and bounded from  $L_1(\mathbb{R}^n)$  to  $WL_1(\mathbb{R}^n)$  for  $p = 1$  (see [50]), then from Theorems 3.2, 3.3, 4.2 and 4.3 we get

**Corollary 5.1.** *Let  $1 \leq p < \infty$ ,  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ . Let also, for  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.10) and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (3.11) and  $\Omega$  satisfies conditions (a)–(c). Then the operator  $\mu_\Omega$  is bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$  for  $p > 1$  and from  $M_{1,\varphi_1}$  to  $WM_{1,\varphi_2}$  for  $p = 1$ .*

**Corollary 5.2.** *Let  $1 \leq p < \infty$ ,  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ . Let also, for  $q' \leq p$ ,  $p \neq 1$ , the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4) and (3.14)–(3.15) and for  $1 < p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4) and (3.16)–(3.17) and  $\Omega$  satisfies conditions (a)–(c). Then the operator  $\mu_\Omega$  is bounded from  $VM_{p,\varphi_1}$  to  $VM_{p,\varphi_2}$  for  $p > 1$  and from  $VM_{1,\varphi_1}$  to  $WVM_{1,\varphi_2}$ .*

**Corollary 5.3.** *Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , is homogeneous of degree zero,  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . Let also, for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.5) and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies condition (4.6) and  $\Omega$  satisfies conditions (a)–(c). Then, the operator  $[b, \mu_\Omega]$  is bounded from  $M_{p,\varphi_1}$  to  $M_{p,\varphi_2}$ .*

**Corollary 5.4.** *Suppose that  $\Omega \in L_q(S^{n-1})$ ,  $1 < q \leq \infty$ , is homogeneous of degree zero,  $1 < p < \infty$  and  $b \in BMO(\mathbb{R}^n)$ . Let also, for  $q' \leq p$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4)–(4.8) and (4.9)–(4.7) and for  $p < q$  the pair  $(\varphi_1, \varphi_2)$  satisfies conditions (2.3)–(2.4)–(4.8) and (4.11)–(4.10) and  $\Omega$  satisfies conditions (a)–(c). Then, the operator  $[b, \mu_\Omega]$  is bounded from  $VM_{p,\varphi_1}$  to  $VM_{p,\varphi_2}$ .*

**Acknowledgement:** This study has been given as the plenary talk by the author at the “3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference (IFSCOM2016), August 29-September 1, 2016, Mersin, Turkey”.

## REFERENCES

- [1] D.R. Adams, A note on Riesz potentials, *Duke Math. J.*, 42 (1975), 765-778.
- [2] A.S. Balakishiyev, V.S. Guliyev, F. Gurbuz and A. Serbetci, Sublinear operators with rough kernel generated by Calderon-Zygmund operators and their commutators on generalized local Morrey spaces, *J. Inequal. Appl.* 2015, 2015:61. doi:10.1186/s13660-015-0582-y.
- [3] L. Caffarelli, Elliptic second order equations, *Rend. Semin. Math. Fis. Milano*, 58 (1990), 253-284.
- [4] X.N. Cao, D.X. Chen, The boundedness of Toeplitz-type operators on vanishing Morrey spaces. *Anal. Theory Appl.* 27 (2011), 309-319.
- [5] F. Chiarenza, M. Frasca, Morrey spaces and Hardy-Littlewood maximal function, *Rend. Mat.*, 7 (1987), 273-279.
- [6] F. Chiarenza, M. Frasca, P. Longo, Interior  $W^{2,p}$ -estimates for nondivergence elliptic equations with discontinuous coefficients, *Ricerche Mat.*, 40 (1991), 149-168.
- [7] F. Chiarenza, M. Frasca, P. Longo,  $W^{2,p}$ -solvability of Dirichlet problem for nondivergence elliptic equations with VMO coefficients, *Trans. Amer. Math. Soc.*, 336 (1993), 841-853.
- [8] R.R. Coifman, R. Rochberg, G. Weiss, Factorization theorems for Hardy spaces in several variables, *Ann. of Math.*, 103 (3) (1976), 611-635.
- [9] R.R. Coifman, P. Lions, Y. Meyer, S. Semmes, Compensated compactness and Hardy spaces. *J. Math. Pures Appl.* 72 (1993), 247-286.
- [10] R.R. Coifman, Y. Meyer, Au delà des Opérateurs Pseudo-Différentiels, *Astérisque* 57. Société Mathématique de France, Paris, 1978, 185 pp.
- [11] Y. Ding, D.C. Yang, Z. Zhou, Boundedness of sublinear operators and commutators on  $L^{p,\omega}(\mathbb{R}^n)$ , *Yokohama Math. J.*, 46 (1998), 15-27.
- [12] G. Di Fazio, M.A. Ragusa, Interior estimates in Morrey spaces for strong solutions to nondivergence form equations with discontinuous coefficients, *J. Funct. Anal.*, 112 (1993), 241-256.
- [13] G. Di Fazio, D.K. Palagachev and M.A. Ragusa, Global Morrey regularity of strong solutions to the Dirichlet problem for elliptic equations with discontinuous coefficients, *J. Funct. Anal.*, 166 (1999), 179-196.
- [14] V.S. Guliyev, Boundedness of the maximal, potential and singular operators in the generalized Morrey spaces, *J. Inequal. Appl.* 2009, Art. ID 503948, 20 pp.
- [15] J. Garcia-Cuerva and J.L. Rubio de Francia, *Weighted Norm Inequalities and Related Topics*, North-Holland Math. 16, Amsterdam, 1985.
- [16] L. Grafakos, *Classical and modern Fourier analysis*. Pearson Education. Inc. Upper Saddle River, New Jersey, 2004.
- [17] V.S. Guliyev, S.S. Aliyev, T. Karaman, P. S. Shukurov, Boundedness of sublinear operators and commutators on generalized Morrey Space, *Integ. Equ. Oper. Theory.*, 71 (3) (2011), 327-355.
- [18] F. Gurbuz, Boundedness of some potential type sublinear operators and their commutators with rough kernels on generalized local Morrey spaces [*Ph.D. thesis*], Ankara University, Ankara, Turkey, 2015 (in Turkish).
- [19] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic Calderón-Zygmund operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, *Open Math.*, 14 (2016), 300-323.
- [20] F. Gurbuz, Parabolic sublinear operators with rough kernel generated by parabolic fractional integral operators and parabolic local Campanato space estimates for their commutators on the parabolic generalized local Morrey spaces, *Advan. Math. (China)*, in press.
- [21] S. Janson, Mean oscillation and commutators of singular integral operators, *Ark. Mat.*, 16 (1978), 263-270.
- [22] F. John and L. Nirenberg, On functions of bounded mean oscillation. *Comm. Pure Appl. Math.*, 14 (1961), 415-426.
- [23] T. Karaman, Boundedness of some classes of sublinear operators on generalized weighted Morrey spaces and some applications [*Ph.D. thesis*], Ankara University, Ankara, Turkey, 2012 (in Turkish).
- [24] Y. Lin, Strongly singular Calderón-Zygmund operator and commutator on Morrey type spaces, *Acta Math. Sin. (Engl. Ser.)*, 23 (11) (2007), 2097-2110.
- [25] G. Lu, S.Z. Lu, D.C. Yang, Singular integrals and commutators on homogeneous groups, *Analysis Mathematica*, 28 (2002), 103-134.

- [26] S.Z. Lu, D.C. Yang, Z. Zhou, Sublinear operators with rough kernel on generalized Morrey spaces. *Hokkaido Math. J.* 27 (1998), 219-232.
- [27] S.Z. Lu, Y. Ding, D.Y. Yan, *Singular integrals and related topics*, World Scientific Publishing, Singapore, 2006.
- [28] S.Z. Lu, Some results and problems on commutators, *Front. Math. China*, 6 (2011), 821-833.
- [29] A. Mazzucato, Besov-Morrey spaces: functions space theory and applications to non-linear PDE., *Trans. Amer. Math. Soc.*, 355 (2002), 1297-1364.
- [30] C. Miranda, Sulle equazioni ellittiche del secondo ordine di tipo non variazionale, a coefficienti discontinui. *Ann. Math. Pura E Appl.* 63 (4) (1963), 353-386.
- [31] T. Mizuhara, Boundedness of some classical operators on generalized Morrey spaces, *Harmonic Analysis* (S. Igari, Editor), ICM 90 Satellite Proceedings, Springer - Verlag, Tokyo (1991), 183-189.
- [32] C.B. Morrey, On the solutions of quasi-linear elliptic partial differential equations, *Trans. Amer. Math. Soc.*, 43 (1938), 126-166.
- [33] B. Muckenhoupt, R.L. Wheeden, Weighted norm inequalities for singular and fractional integrals. *Trans. Amer. Math. Soc.*, 161 (1971), 249-258.
- [34] E. Nakai, Hardy-Littlewood maximal operator, singular integral operators and Riesz potentials on generalized Morrey spaces, *Math. Nachr.*, 166 (1994), 95-103.
- [35] D.K. Palagachev, L.G. Softova, Singular integral operators, Morrey spaces and fine regularity of solutions to PDE's, *Potential Anal.*, 20 (2004), 237-263.
- [36] M. Paluszynski, Characterization of the Besov spaces via the commutator operator of Coifman, Rochberg and Weiss, *Indiana Univ. Math. J.*, 44 (1995), 1-17.
- [37] J. Peetre, On the theory of  $M_{p,\lambda}$ , *J. Funct. Anal.*, 4 (1969), 71-87.
- [38] L.E. Persson, M.A. Ragusa, N. Samko, P. Wall, Commutators of Hardy operators in vanishing Morrey spaces. *AIP Conf. Proc.* 1493, 859 (2012); <http://dx.doi.org/10.1063/1.4765588>.
- [39] M.A. Ragusa, Commutators of fractional integral operators on vanishing-Morrey spaces. *J. Global Optim.* 368 (40) (2008), 1-3.
- [40] A. Ruiz, L. Vega, On local regularity of Schrödinger equations, *Int. Math. Res. Not.*, 1 (1993), 13-27.
- [41] N. Samko, Maximal, Potential and Singular Operators in vanishing generalized Morrey Spaces. *J. Global Optim.* 2013, 1-15. DOI 10.1007/s10898-012-9997-x.
- [42] Y. Sawano, Generalized Morrey Space for Non-doubling Measures, *Nonlinear Differ. Equ. Appl.* 15 (2008), 413-425.
- [43] S.G. Shi, Z.W. Fu, F.Y. Zhao, Estimates for operators on weighted Morrey spaces and their applications to nondivergence elliptic equations, *J. Inequal. Appl.* 2013, 2013: 390.
- [44] S.G. Shi, S.Z. Lu, A characterization of Campanato space via commutator of fractional integral, *J. Math. Anal. Appl.*, **419** (2014), 123-137.
- [45] F. Soria, G. Weiss, A remark on singular integrals and power weights, *Indiana Univ. Math. J.*, 43 (1994) 187-204.
- [46] E.M. Stein, On the functions of Littlewood-Paley, Lusin and Marcinkiewicz, *Trans. Amer. Math. Soc.* 88 (1958) 430-466.
- [47] E.M. Stein, *Singular integrals and differentiability of functions*, Princeton University Press, Princeton, NJ, 1970.
- [48] E.M. Stein, *Harmonic Analysis: Real Variable Methods, Orthogonality and Oscillatory Integrals*, Princeton Univ. Press, Princeton NJ, 1993.
- [49] A. Torchinsky, *Real Variable Methods in Harmonic Analysis*, Pure and Applied Math. 123, Academic Press, New York, 1986.
- [50] A. Torchinsky and S. Wang, A note on the Marcinkiewicz integral, *Colloq. Math.* 60/61 (1990), 235-243.
- [51] C. Vitanza, Functions with vanishing Morrey norm and elliptic partial differential equations. In: *Proceedings of Methods of Real Analysis and Partial Differential Equations*, Capri, pp. 147-150. Springer (1990).
- [52] C. Vitanza, Regularity results for a class of elliptic equations with coefficients in Morrey spaces. *Ricerche di Matematica* 42 (2) (1993), 265-281.
- [53] R.L. Wheeden and A. Zygmund, *Measure and Integral: An Introduction to Real Analysis*, vol. 43 of Pure and Applied Mathematics, Marcel Dekker, New York, NY, USA, 1977.

ANKARA UNIVERSITY, FACULTY OF SCIENCE, DEPARTMENT OF MATHEMATICS, TANDOĞAN 06100,  
ANKARA, TURKEY

*E-mail address:* feritgurbuz84@hotmail.com

**DISTANCE MEASURE, SIMILARITY MEASURE, ENTROPY  
AND INCLUSION MEASURE FOR TEMPORAL  
INTUITIONISTIC FUZZY SETS**

FATIH KUTLU, ÖZKAN ATAN, AND TUNAY BILGIN

**ABSTRACT.** In this study, we proposed distance measures similarity measure, entropy and inclusion measure for temporal intuitionistic fuzzy sets and investigated some properties of these measures. We defined these concepts in two different ways, namely temporal and overall and we examined the relationship between these definitions. Also we gave numerical examples for TIFS. We compared these measures defined with two and three parameters in terms of reliability and applicability.

Received: 30–July–2016

Accepted: 29–August–2016

1. INTRODUCTION

The concept of temporal intuitionistic fuzzy sets is defined by Atanassov in 1991. In this theory membership and non-membership degrees are defined depending on the time-moment and element. This idea leads to a rich field to be used in applications on dynamic fields such as weather, economy, image and video processing. On the other hand, the similarity and distance measures on fuzzy and intuitionistic fuzzy sets, as seen in present studies, are used in many different areas and obtained effective results. Temporal intuitionistic fuzzy measures which achieved by combining these two ideas are still not defined in the literature. This is one of major shortcoming of temporal intuitionistic fuzzy set theory. Temporal intuitionistic fuzzy measurement is a natural consequence of idea that making dynamic measurements used in the dynamic areas.

In this study, firstly we give definitions of temporal intuitionistic fuzzy distance and similarity measures. Then, we investigate some major properties of these measures. Also we investigate how to define these measurements. With more clearly, these measures will be examined by defining which parameters need to adhere to. Additionally, the concept of entropy and inclusion which are closely related to aforementioned measures are defined to the temporal intuitionistic fuzzy sets. Finally, some other basic concepts needed in this context will be defined in temporal space intuitionistic fuzzy sets.

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference

*Key words and phrases.* temporal intuitionistic fuzzy sets, distance measure, similarity measure, entropy, inclusion measure.

## 2. PRELIMINARIES

**Definition 2.1.** [1] An intuitionistic fuzzy set on a non-empty set  $X$  given by a set of ordered triples  $A = \{(x, \mu_A(x), \eta_A(x)) : x \in X\}$  where  $\mu_A(x) : X \rightarrow I = [0, 1]$ ,  $\eta_A(x) : X \rightarrow I$ , are functions such that  $0 \leq \mu(x) + \eta(x) \leq 1$  for all  $x \in X$ . For  $x \in X$ ,  $\mu_A(x)$  and  $\eta_A(x)$  represent the degree of membership and degree of non-membership of  $x$  to  $A$  respectively. For each  $x \in X$ ; intuitionistic fuzzy index of  $x$  in  $A$  is defined as follows  $\pi_A(x) = 1 - \mu_A(x) - \eta_A(x)$ .  $\pi_A$  is the called degree of hesitation or indeterminacy.

**Definition 2.2.** [1] Let  $A, B \in IFS(X)$ . Then,

- (i)  $A \subseteq B \Leftrightarrow \mu_A(x) \leq \mu_B(x)$  and  $\eta_A(x) \geq \eta_B(x)$  for  $\forall x \in X$ ,
- (ii)  $A = B \Leftrightarrow A \subseteq B$  and  $B \subseteq A$ ,
- (iii)  $\bar{A} = \{(x, \eta_A(x), \mu_A(x)) : x \in X\}$ ,
- (iv)  $\bigcap A_i = \{(x, \wedge \mu_{A_i}(x), \vee \eta_{A_i}(x)) : x \in X\}$ ,
- (v)  $\bigcup A_i = \{(x, \vee \mu_{A_i}(x), \wedge \eta_{A_i}(x)) : x \in X\}$ ,
- (vi)  $\tilde{0} = \{(x, 0, 1) : x \in X\}$  and  $\tilde{1} = \{(x, 1, 0) : x \in X\}$ .

**Definition 2.3.** [2] Let  $X$  be an universe and  $T$  be a non-empty time set. We call the elements of  $T$  as "time moments". Based on the definition of IFS, a temporal intuitionistic fuzzy set (TIFS) is defined as the following:

$$A(T) = \{(x, \mu_A(x, t), \eta_A(x, t)) : X \times T\}$$

where:

- a.  $A \subseteq X$  is a fixed set
- b.  $\mu_A(x, t) + \eta_A(x, t) \leq 1$  for every  $(x, t) \in X \times T$
- c.  $\mu_A(x, t)$  and  $\eta_A(x, t)$  are the degrees of membership and non-membership, respectively, of the element  $x \in X$  at the time moment  $t \in T$

For brevity, we write  $A$  instead of  $A(T)$ . The hesitation degree of an TIFS is defined as  $\pi_A(x, t) = 1 - \mu_A(x, t) - \eta_A(x, t)$ . Obviously, every ordinary IFS can be regarded as TIFS for which  $T$  is a singleton set. All operations and operators on IFS can be defined for TIFSs.

By  $TIFS^{(X, T)}$ , we denote to the set of all temporal intuitionistic fuzzy sets defined on  $X$  and time set  $T$ . Obviously, each intuitionistic fuzzy sets can be expressed as temporal intuitionistic fuzzy set via a singular time set. In additionally, all operations and operators defined for intuitionistic fuzzy sets can be defined for temporal intuitionistic fuzzy sets.

**Definition 2.4.** [2] Let

$$A(T') = \{(x, \mu_A(x, t), \eta_A(x, t)) : X \times T'\}$$

and

$$B(T'') = \{(x, \mu_B(x, t), \eta_B(x, t)) : X \times T''\}$$

where  $T'$  and  $T''$  have finite number of distinct time-elements or they are time intervals. Then;

$$A(T') \cap B(T'') =$$

$$\{(x, \min(\bar{\mu}_A(x, t), \bar{\mu}_B(x, t)), \max(\bar{\eta}_A(x, t), \bar{\eta}_B(x, t))) : (x, t) \in X \times (T' \cup T'')\}$$

and

$$A(T') \cup B(T'') = \{(x, \max(\bar{\mu}_A(x, t), \bar{\mu}_B(x, t)), \min(\bar{\eta}_A(x, t), \bar{\eta}_B(x, t))) : (x, t) \in X \times (T' \cup T'')\}$$

Also from definition of subset in IFS theory, Subsets of TIFS can be defined as the following:

$$A(T') \subseteq B(T'') \Leftrightarrow \bar{\mu}_A(x, t) \geq \bar{\mu}_B(x, t) \text{ and } \bar{\eta}_A(x, t) \leq \bar{\eta}_B(x, t) \text{ for every } [(x, t) \in X \times (T' \cup T'')] \text{ where}$$

$$\begin{aligned} \bar{\mu}_A(x, t) &= \begin{cases} \mu_A(x, t) & , \text{ if } t \in T' \\ 0 & , \text{ if } t \in T'' - T' \end{cases} \\ \bar{\mu}_B(x, t) &= \begin{cases} \mu_B(x, t) & , \text{ if } t \in T'' \\ 0 & , \text{ if } t \in T' - T'' \end{cases} \\ \bar{\eta}_A(x, t) &= \begin{cases} \eta_A(x, t) & , \text{ if } t \in T' \\ 1 & , \text{ if } t \in T'' - T' \end{cases} \\ \bar{\eta}_B(x, t) &= \begin{cases} \eta_B(x, t) & , \text{ if } t \in T'' \\ 1 & , \text{ if } t \in T' - T'' \end{cases} \end{aligned}$$

It is obviously seen that if  $T' = T''$ ;  $\bar{\mu}_A(x, t) = \mu_A(x, t)$ ,  $\bar{\mu}_B(x, t) = \mu_B(x, t)$ ,  $\bar{\eta}_A(x, t) = \eta_A(x, t)$ ,  $\bar{\eta}_B(x, t) = \eta_B(x, t)$ . [2]

Let  $J$  be an index set and  $T_i$  is a time set for each  $i \in J$ . Let define that  $T = \bigcup_{i \in J} T_i$ . Now we extend union and intersection of temporal intuitionistic fuzzy sets to the family  $F = \{A_i(T_i) = (x, \mu_{A_i}(x, t), \eta_{A_i}(x, t)) : x \in X \times T_i, i \in J\}$  as:

$$\begin{aligned} \bigcup_{i \in J} A(T_i) &= \left\{ \left( x, \max_{i \in J} (\bar{\mu}_{A_i}(x, t)), \min_{i \in J} (\bar{\eta}_{A_i}(x, t)) : (x, t) \in X \times T \right) \right\}, \\ \bigcap_{i \in J} A(T_i) &= \left\{ \left( x, \min_{i \in J} (\bar{\mu}_{A_i}(x, t)), \max_{i \in J} (\bar{\eta}_{A_i}(x, t)) : (x, t) \in X \times T \right) \right\} \end{aligned}$$

where

$$\begin{aligned} \bar{\mu}_{A_i}(x, t) &= \begin{cases} \mu_{A_i}(x, t) & , \text{ if } t \in T_i \\ 0 & , \text{ if } t \in T - T_i \end{cases} \\ \bar{\eta}_{A_i}(x, t) &= \begin{cases} \eta_{A_i}(x, t) & , \text{ if } t \in T_i \\ 1 & , \text{ if } t \in T - T_i \end{cases} \end{aligned}$$

### 3. DISTANCE MEASURE, SIMILARITY MEASURE, ENTROPY AND INCLUSION MEASURE FOR TEMPORAL INTUITIONISTIC FUZZY SETS

Let  $X$  be a universe and  $T$  be a non-empty time-moment set. The definition of distance measure defined in [31] can be extended to  $TIFS^{(X, T)}$  such as:

**Definition 3.1.** Let  $X$  be a universe,  $T$  be a non-empty time-moment set and  $d^t : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow R^+ \cup \{0\}$  be a mapping for fixed  $t \in T$ . If  $d^t$  satisfies following properties for all  $A, B \in TIFS^{(X, T)}$  and fixed time-moment  $t \in T$ , then  $d^t$  is called a temporal intuitionistic fuzzy distance measure on  $TIFS^{(X, T)}$  at time-moment  $t$ :

- D1.  $d^t(A, B) = 0 \Leftrightarrow A = B$ ,
- D2.  $d^t(A, B) = d^t(B, A)$ ,
- D3. If  $A$  is crisp set,  $d^t(A, \bar{A}) = \max_{B, C \in TIFS^{(X, T)}} d^t(B, C)$



D4. If  $A \subseteq B \subseteq C$  for  $A, B, C \in TIFS^{(X,T)}$ ,  $d^t(A, C) \geq d^t(A, B)$  and  $d^t(A, C) \geq d^t(B, C)$ .

If there exist a mapping  $d : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow R^+ \cup \{0\}$  which satisfies these conditions for every time moment  $t \in T$ , then  $d$  is called a overall intuitionistic fuzzy distance measure on  $TIFS^{(X,T)}$ . In order to distinguish between these two concepts from each other, the first measure is referred as temporal intuitionistic fuzzy distance measure at just only time moment  $t$ . But the second one measures overall distance between temporal intuitionistic fuzzy sets in full time range.

Distance measure between temporal intuitionistic fuzzy sets in terms of being discrete time set or interval time set can be defined in the following way (see for more information: [4], [13], [16], [18], [20], [22], [31], [33], [34], [37], [43]):

**Theorem 3.1.** Let  $X$  be non-empty set and  $T = \{t_1, t_2, t_3, \dots, t_n\}$  be finite and distinct time set. Let define  $A, B \in TIFS^{(X,T)}$  such as

$$A(T) = \{(x, \mu_A(x, t), \eta_A(x, t)) : (x, t) \in X \times T\}$$

and

$$B(T) = \{(x, \mu_B(x, t), \eta_B(x, t)) : (x, t) \in X \times T\}$$

, respectively. Then we define  $d^i_t(A, B)$  mappings for  $t \in T$  and  $i = 1, 2, 3, 4$  as following:

1.  $d^1_{t_0}(A, B) = \sqrt{\sum_{x \in X} \left( (\mu_A(x, t_0) - \mu_B(x, t_0))^2 + (\eta_A(x, t_0) - \eta_B(x, t_0))^2 \right)}$ ,
2.  $d^2_{t_0}(A, B) = \sum_{x \in X} (|\mu_A(x, t_0) - \mu_B(x, t_0)| + |\eta_A(x, t_0) - \eta_B(x, t_0)|)$ ,
3.  $d^3_{t_0}(A, B) = \sum_{x \in E} (\max\{|\mu_A(x, t_0) - \mu_B(x, t_0)|, |\eta_A(x, t_0) - \eta_B(x, t_0)|\})$ ,
4.  $d^4_{t_0}(A, B) = \sqrt{\sum_{x \in E} \max\left\{ (\mu_A(x, t_0) - \mu_B(x, t_0))^2, (\eta_A(x, t_0) - \eta_B(x, t_0))^2 \right\}}$ .

It is clear that each mapping  $d^i_t(A, B)$  is a temporal intuitionistic fuzzy distance measures for  $t \in T$ . On the other hand, the following temporal distance measures are obtained by adding the degree of uncertainty to former ones.

5.  $D^1_{t_0}(A, B) =$

$$\sqrt{\sum_{x \in X} \left( (\mu_A(x, t_0) - \mu_B(x, t_0))^2 + (\eta_A(x, t_0) - \eta_B(x, t_0))^2 + (\pi_A(x, t_0) - \pi_B(x, t_0))^2 \right)}$$

6.  $D^2_{t_0}(A, B) =$

$$\sum_{x \in X} (|\mu_A(x, t_0) - \mu_B(x, t_0)| + |\eta_A(x, t_0) - \eta_B(x, t_0)| + |\pi_A(x, t_0) - \pi_B(x, t_0)|)$$

7.  $D^3_{t_0}(A, B) =$

$$\sum_{x \in E} \max\{|\mu_A(x, t_0) - \mu_B(x, t_0)|, |\eta_A(x, t_0) - \eta_B(x, t_0)|, |\pi_A(x, t_0) - \pi_B(x, t_0)|\}$$

8.  $D^4_{t_0}(A, B) =$

$$\sqrt{\sum_{x \in E} \max\left\{ (\mu_A(x, t_0) - \mu_B(x, t_0))^2, (\eta_A(x, t_0) - \eta_B(x, t_0))^2, (\pi_A(x, t_0) - \pi_B(x, t_0))^2 \right\}}$$

Additionally, the mappings  $d_i$  (or  $D_i$ ) defined as  $d_i(A, B) = \max_{t \in T} (d_t^i(A, B))$  (or  $D_i(A, B) = \max_{t \in T} (D_t^i(A, B))$ ) are overall temporal intuitionistic fuzzy distance measures for  $i = 1, 2, 3, 4$ . It is clear that these temporal and overall intuitionistic fuzzy distance measures are intuitionistic fuzzy distance measure for a singleton time set.

*Proof.* As it can be seen in studies ([4], [13], [16], [18], [20], [22], [31], [33], [34], [37], [43], etc.) temporal intuitionistic fuzzy distance measures defined above is obtained by the addition of time parameters to the intuitionistic fuzzy distance measures. As stated previously, these measures are also intuitionistic fuzzy distance measure for each individual time moment. Now we prove that  $d_1(A, B) = \max_{t \in T} (d_t^1(A, B))$  is an overall intuitionistic fuzzy distance measure.

**D1:** Since  $d_t^1(A, A) = 0$  for all  $t \in T$  and  $A \in TIFS^{(X, T)}$ , it is clearly obtained that  $d_1(A, A) = 0$ .

**D2:** Since  $d_t^1(A, B) = d_t^1(B, A)$  for all  $t \in T$  and  $A, B \in TIFS^{(X, T)}$ , Then it is clearly obtained that  $d_1(A, B) = \max_{t \in T} (d_t^1(A, B)) = \max_{t \in T} (d_t^1(B, A)) = d_1(B, A)$ .

**D3:** Since  $d_t^1(A, \bar{A}) = \max_{B, C \in TIFS^{(X, T)}} d_t^1(B, C)$  for all  $t \in T$  and  $A$  crisp set.

Then, we can easily get  $d^1(A, \bar{A}) = \max_{t \in T} (d_t^1(A, \bar{A})) = \max_{t \in T} \left\{ \max_{B, C \in TIFS^{(X, T)}} d_t^1(B, C) \right\} = \max_{B, C \in TIFS^{(X, T)}} \left\{ \max_{t \in T} d_t(B, C) \right\} = \max_{B, C \in TIFS^{(X, T)}} d^1(B, C)$

**D4:** From the definition of being subset in concept of TIFS, When  $A \subseteq B \subseteq C$  for  $A, B, C \in TIFS^{(X, T)}$ , the inequalities  $\mu_A(x, t) \geq \mu_B(x, t) \geq \mu_C(x, t)$  and  $\eta_A(x, t) \leq \eta_B(x, t) \leq \eta_C(x, t)$  are hold for each  $(x, t) \in X \times T$ . From the last two inequalities the inequalities  $d_t^1(A, C) \geq d_t^1(A, B)$  and  $d_t^1(A, C) \geq d_t^1(B, C)$  are obtained for  $t \in T$ . Then the inequalities  $d_1(A, C) \geq d_1(A, B)$  and  $d_1(A, C) \geq d_1(B, C)$  are clearly obtained from definition of  $d_1$ . The other situations can be proved similarly.  $\square$

Each distance measures  $d_t^i$  indicates temporal distance between the temporal intuitionistic fuzzy sets at time moment  $t \in T$ . On the other hand,  $d_i$  measurements which are obtained from the maximum of  $d_t^i$  gives a overall distance measurement between temporal intuitionistic fuzzy sets. These two approaches gain different importance degrees depending on the applications. With more open expression, overall distance measure expresses inferential distance in the total situation, while since the temporal distance between the temporal intuitionistic fuzzy sets are sensitive to sudden changes at distance, temporal distance is a measure of instant change between cases represented by the temporal intuitionistic fuzzy set. This situation offers multiple ways to ensure expected benefits from the application.

We give definition of temporal intuitionistic distance measures on infinite and interval time set as follow:

**Proposition 3.1.** *Let  $X$  be a infinite set and  $T = \{t_1, t_2, \dots, t_i, \dots\}$  be a infinite time set (or time interval). Let suppose that  $t_k \leq t_{k+1}$  for each  $t_k, t_{k+1} \in T$ . On the other hand, let define  $A, B \in TIFS^{(X, T)}$  TIFSs as follows:*

$$A(T) = \{(x, \mu_A(x, t), \eta_A(x, t)) : (x, t) \in X \times T\}$$

and

$$B(T) = \{(x, \mu_B(x, t), \eta_B(x, t)) : (x, t) \in X \times T\},$$

respectively. With these definitions, the following statements are overall intuitionistic fuzzy distance measure on  $TIFS^{(X,T)}$

1.  $d^*_1(A, B) =$

$$\sup_{t \in T} \left\{ \max \left\{ \sup_{x \in X} |\mu_A(x, t) - \mu_B(x, t)|, \sup_{x \in X} |\eta_A(x, t) - \eta_B(x, t)| \right\} \right\}$$

2.  $d^*_2(A, B) =$

$$\sup_{t \in T} \left\{ \sup_{x \in X} (|\mu_A(x, t) - \mu_B(x, t)| + |\eta_A(x, t) - \eta_B(x, t)|) \right\}$$

3.  $d^*_3(A, B) =$

$$\sup_{t \in T} \sqrt{\sup_{x \in X} \left( (\mu_A(x, t) - \mu_B(x, t))^2 + (\eta_A(x, t) - \eta_B(x, t))^2 \right)}$$

4.  $D^*_1(A, B) =$

$$\sup_{t \in T} \left\{ \max \left\{ \sup_{x \in X} |\mu_A(x, t) - \mu_B(x, t)|, \sup_{x \in X} |\eta_A(x, t) - \eta_B(x, t)|, \sup_{x \in X} |\pi_A(x, t) - \pi_B(x, t)| \right\} \right\}$$

5.  $D^*_2(A, B) =$

$$\sup_{t \in T} \left( \sup_{x \in X} (|\mu_A(x, t) - \mu_B(x, t)| + |\eta_A(x, t) - \eta_B(x, t)| + |\pi_A(x, t) - \pi_B(x, t)|) \right)$$

6.  $D^*_3(A, B) =$

$$\sup_{t \in T} \sqrt{\sup_{x \in X} \left( (\mu_A(x, t) - \mu_B(x, t))^2 + (\eta_A(x, t) - \eta_B(x, t))^2 + (\pi_A(x, t) - \pi_B(x, t))^2 \right)}$$

(see [4], [13], [16], [18], [20], [22], [31], [33], [34], [37], [43], etc.)

**Proposition 3.2.** Let  $X$  be an infinite set and  $T = [t_1, t_2]$  for  $t_1, t_2 \in R^+$  and  $t_1 < t_2$ . Then the following statements are overall intuitionistic c fuzzy distance measures on  $TIFS^{(X,T)}$

1.  $D^{***}_1(A, B) = \sup_{t \in T} \left\{ \int_X (\mu_A(x, t) - \mu_B(x, t)) .dx + \left| \int_X (\eta_A(x, t) - \eta_B(x, t)) .dx \right| \right\}$

2.  $D^{***}_2(A, B) =$

$$\sup_{t \in T} \left\{ \left| \int_X (\mu_A(x, t) - \mu_B(x, t)) .dx \right| + \left| \int_X (\eta_A(x, t) - \eta_B(x, t)) .dx \right| + \left| \int_X (\pi_A(x, t) - \pi_B(x, t)) .dt \right| \right\}$$

for  $A, B \in TIFS^{(X,T)}$  and  $t \in T = [t_1, t_2]$ . (see [4], [13], [16], [18], [20], [22], [31], [33], [34], [37], [43], etc.)

**Example 3.1.** Let suppose that  $X = [0, 4]$  and  $T = [0, 5]$ . Let define that

$$A(T) = \{(x, t), \mu_A(x, t), \eta_A(x, t)\}; (x, t) \in X \times T\}$$

and

$$B(T) = \{(x, t), \mu_B(x, t), \eta_B(x, t)\}; (x, t) \in X \times T\}$$




FIGURE 1




FIGURE 2

where degrees of membership and non-membership are defined as follows respectively  $\mu_A(x, t) = e^{-10\frac{(x-1)^2}{t+1}}$  and  $\eta_A(x, t) = 1 - e^{-\frac{(x-1)^2}{t+1}}$  ;  $\mu_B(x, t) = \frac{1}{1+e^{-t.(x-2)^2}}$  and  $\eta_B(x, t) = \frac{e^{-t.(x-2)^2}}{2+e^{-t.(x-2)^2}}$  for all  $(x, t) \in X \times T$ . 3D- graphics of  $\mu_A, \eta_A, \mu_B$  and  $\eta_B$  are given in Fig.1,2,3,4, respectively.

Fig 1. Graphic of temporal intuitionistic fuzzy membership function  $\mu_A$

Fig 2. Graphic of temporal intuitionistic fuzzy non-membership function  $\eta_A$

Fig 3. Graphic of temporal intuitionistic fuzzy membership function  $\mu_B$

Fig 4. Graphic of temporal intuitionistic fuzzy non-membership function  $\eta_B$

In the following figures, we give changing of distance between  $A$  and  $B$  obtained by  $D_1^{***}$  and  $D_2^{***}$  over time in Fig 5. and Fig. 6.

Fig. 5. Changing of distance between  $A$  and  $B$  obtained by  $D_1^{***}$




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6

Fig. 6. Changing of distance between  $A$  and  $B$  obtained by  $D_2^{***}$ 

From these figures, the following overall distances are obtained

$$d_{D_1^{***}}(A, B) = \sup_{t \in T} \{D_1^{***}(A, B)\} = 3,391$$

and

$$d_{D_2^{***}}(A, B) = \sup_{t \in T} \{D_2^{***}(A, B)\} = 5,415$$

respectively. A key issue at this point is  $D_2^{***}$  temporal distance measure is more durable and more reliable than  $D_1^{***}$  in high degrees of uncertainty. As previously stated in various studies of Szmidt and Kacprzyk for intuitionistic fuzzy sets, in the cases which contains data with high degree of uncertainty, it is obvious that temporal (or overall) distance which obtained with three parameters between temporal intuitionistic fuzzy sets is more accurate than temporal (or overall) distance which obtained with two parameters.

We use aggregation function to generalize the correlation between temporal and overall intuitionistic fuzzy distance as used in [11].

**Theorem 3.2.** *Let  $X$  be a non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be a finite time set. Let suppose that  $d_t : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  is a normal temporal intuitionistic fuzzy sets for  $t \in T$  and  $f$  is a  $n$ -aggregation function without zero divisor. Then, the mapping  $d : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  which defined as:*

$$d(A, B) = f(d_{t_1}(A, B), d_{t_2}(A, B), \dots, d_{t_n}(A, B))$$

for  $A, B \in TIFS^{(X,T)}$  is a overall intuitionistic fuzzy distance measure.

*Proof.* Since  $d_t : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  is a intuitionistic fuzzy distance measure on TIFSs for each  $t \in T$ .  $d(A, B) = f(d_{t_1}(A, B), d_{t_2}(A, B), \dots, d_{t_n}(A, B))$  is a intuitionistic fuzzy distance measure (see [11]). Due to  $d_t$  expresses temporal

distance between  $A$  and  $B$  for each  $t \in T$ , which is obtained by  $d_t$  expresses overall distance between  $A$  and  $B$ .  $\square$

Now we give definitions of temporal and overall intuitionistic fuzzy similarity measure in sense of [31]

**Definition 3.2.** Let  $X$  be a non-empty set,  $T$  be time set and  $s^t : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  be a mapping. If  $s^t$  satisfies following conditions for all  $A, B, C \in TIFS^{(X,T)}$  and fixed time moment  $t \in T$ ,  $s_t$  is called temporal intuitionistic fuzzy similarity measure on  $TIFS^{(X,T)}$  at time moment  $t$ .

- S1. If  $A$  is a crisp set,  $s^t(A, \bar{A}) = 0$
- S2.  $A = B \Leftrightarrow S(A, B) = 1$  for all  $A, B \in TIFS^{(X,T)}$
- S3.  $s^t(A, B) = s^t(B, A)$  for all  $A, B \in TIFS^{(X,T)}$
- S4. The inequalities  $s^t(A, C) \leq s^t(A, B)$  and  $s^t(A, C) \leq s^t(B, C)$  are satisfied for all  $A, B, C \in TIFS^{(X,T)}$  which are satisfied  $A \subseteq B \subseteq C$

As in the concept of temporal distance measure, similarity measure can be examined in two parts as named temporal and overall. Let us give examples of temporal intuitionistic fuzzy similarity measure can be defined in accordance with this approach. It is clear that these measures are obtained by changing domain set of intuitionistic fuzzy similarity measure which are defined in the literature (see [4, 5, 6, 8, 10, 13, 23, 26, 27, 28, 29, 30, 31, 33, 34, 36, 37, 41, 44, 45]) as  $TIFS^{(X,T)}$  :

$$1. S_1^{t_0}(A, B) =$$

$$1 - \frac{\sum_{i=1}^n \left( |\mu_A(x_i, t_0) - \mu_B(x_i, t_0)|^{t_0} + |\eta_A(x_i, t_0) - \eta_B(x_i, t_0)|^{t_0} \right)^{\frac{1}{t_0}}}{2n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$  and  $A, B \in TIFS^{(X,T)}$ ,

$$2. S_2^{t_{j_0}}(A, B) =$$

$$1 - \frac{\sum_{i=1}^n \omega_{(i, j_0)} (|\mu_A(x_i, t_{j_0}) - \mu_B(x_i, t_{j_0})| + |\eta_A(x_i, t_{j_0}) - \eta_B(x_i, t_{j_0})|)}{2n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_{j_0} \in T$ ,  $A, B \in TIFS^{(X,T)}$  and  $\sum_{i=1}^n \omega_{(i, j_0)} = 1$  for each  $j_0 \in \{1, 2, \dots, m\}$  where  $\omega_{(i, j)} \in [0, 1]$  for each  $(i, j) \in \{1, 2, \dots, n\} \times \{1, 2, \dots, m\}$ .

$$3. S_3^{t_{j_0}}(A, B) =$$

$$1 - \frac{\sum_{i=1}^n \omega_{(i, j_0)} (|\mu_A(x_i, t_{j_0}) - \mu_B(x_i, t_{j_0})| + |\eta_A(x_i, t_{j_0}) - \eta_B(x_i, t_{j_0})|)}{2n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_{j_0} \in T$ ,  $A, B \in TIFS^{(X,T)}$  and  $\sum_{i=1}^n \omega_{(i, j_0)} = 1$  for each  $j_0 \in \{1, 2, \dots, m\}$  where  $\omega_{(i, j)} \in [0, 1]$  for each  $(i, j) \in \{1, 2, \dots, n\} \times \{1, 2, \dots, m\}$ .

$$4. S_4^{t_{j_0}}(A, B) = \frac{\sum_{i=1}^n \omega_{(i, j_0)} (|\mu_A(x_i, t_{j_0}) - \mu_B(x_i, t_{j_0})| + |\eta_A(x_i, t_{j_0}) - \eta_B(x_i, t_{j_0})| + |\pi_A(x_i, t_{j_0}) - \pi_B(x_i, t_{j_0})|)}{2n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_{j_0} \in T$ ,  $A, B \in TIFS^{(X, T)}$  and  $\sum_{i=1}^n \omega_{(i, j_0)} = 1$  for each  $j_0 \in \{1, 2, \dots, m\}$  where  $\omega_{(i, j)} \in [0, 1]$  for each  $(i, j) \in \{1, 2, \dots, n\} \times \{1, 2, \dots, m\}$ .

$$5. S_5^{t_0}(A, B) = \frac{\frac{1}{n} \sum_{i=1}^n \frac{\min\{\mu_A(x_i, t_0), \mu_B(x_i, t_0)\} + \min\{\eta_A(x_i, t_0), \eta_B(x_i, t_0)\}}{\max\{\mu_A(x_i, t_0), \mu_B(x_i, t_0)\} + \max\{\eta_A(x_i, t_0), \eta_B(x_i, t_0)\}}}{n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$ ,  $A, B \in TIFS^{(X, T)}$

$$6. S_6^{t_0}(A, B) = \frac{\sum_{i=1}^n 1 - \frac{1}{2} (|\mu_A(x_i, t_0) - \mu_B(x_i, t_0)| + |\eta_A(x_i, t_0) - \eta_B(x_i, t_0)|)}{n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$ ,  $A, B \in TIFS^{(X, T)}$

$$7. S_7^{t_0}(A, B) = \frac{\frac{1}{n} \sum_{i=1}^n \frac{\min\{\mu_A(x_i, t_0), \mu_B(x_i, t_0)\} + \min\{\eta_A(x_i, t_0), \eta_B(x_i, t_0)\}}{\max\{\mu_A(x_i, t_0), \mu_B(x_i, t_0)\} + \max\{\eta_A(x_i, t_0), \eta_B(x_i, t_0)\}}}{n}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$ ,  $A, B \in TIFS^{(X, T)}$

$$8. S_8^{t_0}(A, B) = 1 - \frac{1}{2} (\max |\mu_A(x_i, t_0) - \mu_B(x_i, t_0)| + \max |\eta_A(x_i, t_0) - \eta_B(x_i, t_0)|)$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$ ,  $A, B \in TIFS^{(X, T)}$

$$9. S_9^{t_0}(A, B) = 1 - \frac{\sum_{i=1}^n (|\mu_A(x_i, t_0) - \mu_B(x_i, t_0)| + |\eta_A(x_i, t_0) - \eta_B(x_i, t_0)|)}{\sum_{i=1}^n (|\mu_A(x_i, t_0) + \mu_B(x_i, t_0)| + |\eta_A(x_i, t_0) + \eta_B(x_i, t_0)|)}$$

where  $X = \{x_1, x_2, \dots, x_n\}$ ,  $T = \{t_1, t_2, \dots, t_m\}$ ,  $t_0 \in T$ ,  $A, B \in TIFS^{(X, T)}$

In these definitions, it is seen that the temporal similarity measures are dependent on  $t$  the moment with selected temporal intuitionistic fuzzy sets. Since these TIFSs change over time, similarity measures on  $TIFS^{(X, T)}$  inevitably change over time. This approach elicits a more spacious work area for applications changed in comparison mechanism. there are many different methods to achieve the overall similarity measure defined from a temporal intuitionistic fuzzy similarity measures. Among these the most significant are defined as follows:

**Theorem 3.3.** *Let  $X$  be non-empty set,  $T$  be time set and  $s^t$  be a temporal intuitionistic fuzzy similarity measure for each  $t \in T$ . Then the mapping  $s$  defined as  $s(A, B) = \max_{t \in T} \{s^t(A, B)\}$  is a overall intuitionistic fuzzy similarity measure.*

More general version of this theorem with Du and Xu's approach [11] can be given as follows:

**Theorem 3.4.** *: Let  $X$  be non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be a time set. Let suppose that the mappings  $s^t : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow [0, 1]$  are temporal intuitionistic fuzzy similarity measure for each  $t \in T$  and  $f$  is a  $n$ -aggregation function without zero divisor. Then, the mapping  $s : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow [0, 1]$  defined as:*

$$s(A, B) = f(s^{t_1}(A, B), s^{t_2}(A, B), \dots, s^{t_n}(A, B))$$

for all  $A, B \in TIFS^{(X, T)}$  is a overall intuitionistic fuzzy similarity measure.



*Proof.* It can be proven as Theorem 1. □

As noted for temporal intuitionistic fuzzy distance measure, all similarity measures defined for intuitionistic fuzzy sets can be accepted as temporal intuitionistic fuzzy similarity measure for singular time set and they also can be converted into temporal intuitionistic fuzzy similarity measure by selecting the domain set  $TIFS^{(X,T)}$ . After this process, overall intuitionistic fuzzy similarity measurements indicating the general jurisdiction can be obtained by using some methods such as aggregation function. The most common feature of data used in real-world applications is that they can change over time. In this context, the most remarkable feature is that the uncertainty of the situation will may increase sometimes. As in fuzzy and intuitionistic fuzzy sets, temporal (or overall) intuitionistic fuzzy distance and similarity measures are dual concepts. So, there are several ways to obtain other one from another. In [11], Du and Xu have generalized this relationship by fuzzy negation and aggregation function for intuitionistic fuzzy distance and similarity measures. Their feature is also available in the temporal intuitionistic fuzzy sets. Having not generalized some basic concepts for temporal intuitionistic fuzzy sets is shortcoming in the literature. Some of these concepts which given in [11] are generalized for the temporal intuitionistic fuzzy sets as follows:

**Definition 3.3.** Let  $T$  be a time set. If the mapping  $N_t : [0, 1] \rightarrow [0, 1]$  is satisfied following condition for  $t \in T$ , it is called temporal fuzzy negation at time moment  $t$  :

N1.  $N_t(0) = 1, N_t(1) = 0$

N2.  $N_t(b) \leq N_t(a)$  for all  $a \leq b$

if  $N_t$  is satisfied

a.  $N_t(N_t(a)) = a$  for  $t \in T$  and all  $a \in [0, 1]$  , it is called temporal fuzzy strong negation at time moment  $t$ ,

b.  $x = 0 \Leftrightarrow N_t(x) = 1$  for  $t \in T$  and all  $a \in [0, 1]$  , it is called temporal fuzzy non-filling negation at time moment  $t$ ,

c.  $x = 1 \Leftrightarrow N_t(x) = 0$  for  $t \in T$  and all  $a \in [0, 1]$  , it is called temporal fuzzy non-vanishing negation at time moment  $t$ .

The novelty of this definition is that negation will change over time. Adding the time parameters and changing temporal fuzzy negation over time offers unlimited options for obtaining similarity measure from distance measure (or conversely). the temporal fuzzy strong negations can be obtained by adding time parameters to fuzzy negations as follows:

1.  $N_{1,\lambda_t}(x) = \frac{1-x}{1+\lambda_t x}$  for  $\lambda_t \in (-1, +\infty)$  and  $t \in T$ ,

2.  $N_{2,\delta_t}(x) = \sqrt[\delta_t]{1-x^{\delta_t}}$  for  $\delta_t \in (0, +\infty)$  and  $t \in T$ ,

3.  $N_{3,\varphi_t}(x) = \begin{cases} 1, & x \leq \varphi_t \\ 0, & otherwise \end{cases}$  for  $\varphi_t \in (0, 1)$  and  $t \in T$ .

Following theorems are obtained by adding time parameter to theorems which given by Du and Xu [11].

**Theorem 3.5.** Let  $X$  be non-empty set,  $T$  be time set and  $A, B \in TIFS^{(X,T)}$ . Let suppose that  $d^t$  is a temporal distance measure and  $N_t$  is a temporal fuzzy

non-filling negation. Then, the mapping  $s_{N_t} : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $s_{N_t}(A, B) = N_t(d^t(A, B))$  is a temporal intuitionistic fuzzy similarity measure. Conversely, Let suppose that  $s^t$  is a temporal intuitionistic fuzzy similarity measure and  $N_t$  is a temporal fuzzy non-vanishing negation, then the mapping  $d_{N_t} : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $d_{N_t}(A, B) = N_t(s^t(A, B))$  is a temporal intuitionistic fuzzy distance measure. If  $N_t$  is a temporal fuzzy strong negation, the equations  $d^t(A, B) = N^t(s_{N_t}(A, B))$  and  $s^t(A, B) = N_t(d_{N_t}(A, B))$  are satisfied.

This theorem is preserved with Du and Xu's approach [11] for overall temporal measures as follows.

**Theorem 3.6.** *Let  $X$  be non-empty set,  $T$  be time set and  $A, B \in TIFS^{(X,T)}$ . Let suppose that  $d$  is overall intuitionistic fuzzy distance measure and  $N$  is a fuzzy non-filling negation. Then, the mapping  $s_N : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $s_N(A, B) = N(d(A, B))$  is an overall intuitionistic fuzzy similarity measure. Conversely, let suppose that  $s$  is an overall intuitionistic fuzzy similarity measure and  $N$  is a fuzzy non-vanishing negation, then the mapping  $d_N : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $d_N(A, B) = N(s(A, B))$  is a overall intuitionistic fuzzy similarity measure. If  $N$  is a fuzzy strong negation, the equations  $d(A, B) = N(s_N(A, B))$  and  $s(A, B) = N(d_N(A, B))$  are satisfied.*

Another theorem which is given in [11] can be generalized to temporal intuitionistic fuzzy sets as below.

**Theorem 3.7.** *Let  $X$  be a non empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be a finite time set. Let suppose that  $d^t : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  is a temporal intuitionistic fuzzy distance measure for each  $t \in T$ ,  $f$  is a aggregation function without zero divisor and  $N_t$  is a temporal fuzzy non-filling negation, then the mapping  $s : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as*

$$s(A, B) = f(s_{N_{t_1}}(A, B), s_{N_{t_2}}(A, B), \dots, s_{N_{t_n}}(A, B))$$

for all  $A, B \in TIFS^{(X,T)}$  is a overall intuitionistic fuzzy similarity measure.

*Proof.* It is clear that each  $s_{N_t}$  is a temporal intuitionistic similarity measure for  $t \in T$  from Theorem 5. Then, it is obtained that  $s$  is a overall intuitionistic fuzzy similarity measure from Theorem 4.  $\square$

Now, entropy which is a measure of difference between intuitionistic fuzzy set (or fuzzy sets) and crisp set will be defined for temporal intuitionistic fuzzy sets. the temporal variability encountered in real-world problems changes this difference in a continuous manner. The definition of entropy (as temporal entropy and overall entropy) with Szmidt and Kacprzyk's approach [35] is defined for temporal intuitionistic fuzzy sets as follow:

**Definition 3.4.** Let  $X$  be a non-empty set and  $T$  be a time set. If the mapping  $e^t : TIFS^{(X,T)} \rightarrow [0, 1]$  is satisfied the following conditions for  $A \in TIFS^{(X,T)}$  and fixed  $t \in T$ ,  $e^t$  is called temporal intuitionistic fuzzy entropy on  $TIFS^{(X,T)}$ .

- E1.  $A$  is a crisp set for  $\Leftrightarrow e^t(A) = 0$ ,
- E2.  $e^t(A) = 1 \Leftrightarrow \mu_A(x, t) = \eta_A(x, t)$  for all  $x \in X$  and fixed  $t \in T$

E3.  $e^t(A) \leq e^t(B) \Leftrightarrow$

- i.  $\mu_A(x, t) \geq \mu_B(x, t)$  and  $\eta_A(x, t) \leq \eta_B(x, t)$  for  $\mu_B(x, t) \leq \eta_B(x, t)$
  - ii.  $\mu_A(x, t) \leq \mu_B(x, t)$  and  $\eta_A(x, t) \geq \eta_B(x, t)$  for  $\mu_B(x, t) \geq \eta_B(x, t)$
- for all  $x \in X$  and fixed  $t \in T$

E4.  $e^t(A) = e^t(\overline{A})$ .

With this definition, the value  $e^t(A)$  represents how much is far from being crisp sets at time moment  $t$ . If we change the condition E1 with

E1\*.  $A$  is a fuzzy set  $\Leftrightarrow e^t(A) = 0$

, the value  $e^t(A)$  represents how much is far from being fuzzy sets at time moment  $t$ . Let's call this second measure as type-2 temporal entropy. On the other hand, If the mapping  $e : TIFS^{(X,T)} \rightarrow [0, 1]$  is satisfied the conditions (E1, E2, E3, E4) for each  $t \in T$ ,  $e$  is called overall entropy on  $TIFS^{(X,T)}$ . The mapping  $e : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $e(A) = \sup_{t \in T} \{e^t(A)\}$  is an overall intuitionistic fuzzy entropy.

**Theorem 3.8.** *Let  $X$  be a non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be a finite time set. Let suppose that  $e^t : TIFS^{(X,T)} \rightarrow [0, 1]$  is temporal intuitionistic fuzzy entropy for  $t \in T$  and  $f$  is a aggregation function without zero divisor. Then the mapping  $e : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as*

$$e(A) = f(e^{t_1}(A), e^{t_2}(A), \dots, e^{t_n}(A))$$

for  $A \in TIFS^{(X,T)}$  is an overall intuitionistic fuzzy entropy.

*Proof.* It can be proven as Theorem 1. □

The intuitionistic fuzzy entropies which are defined in [12, 14, 19, 21, 24, 25, 28, 29, 30, 31, 35, 36, 38, 41, 42, 43, 45] can be converted into the temporal intuitionistic fuzzy entropy with some minor modifications as follows:

**Proposition 3.3.** *Let  $X = \{x_1, x_2, \dots, x_n\}$  be a non-empty set and  $T$  is a time set, Then the following mappings are temporal intuitionistic fuzzy entropy on  $TIFS^{(X,T)}$*

1.  $e_{SK}^t(A) = 1 - \frac{1}{n} \sum_{i=1}^n |\mu_A(x_i, t) - \eta_A(x_i, t)|$
2.  $e_{VS-1}^t = \frac{\sum_{i=1}^n \min\{\mu_A(x_i, t), \eta_A(x_i, t)\} + \min\{1 - \mu_A(x_i, t), 1 - \eta_A(x_i, t)\}}{\sum_{i=1}^n \max\{\mu_A(x_i, t), \eta_A(x_i, t)\} + \max\{1 - \mu_A(x_i, t), 1 - \eta_A(x_i, t)\}}$
3.  $e_{VS-2}^t(A) = \sum_{i=1}^n \frac{2\mu_A(x_i, t)\eta_A(x_i, t) + \pi_A(x_i, t)^2}{\mu_A(x_i, t)^2 + \eta_A(x_i, t)^2 + \pi_A(x_i, t)^2}$
4.  $e_{Li-1}^t(A) = 1 - \frac{1}{2n} \sum_{i=1}^n \left( |\mu_A(x_i, t) - \eta_A(x_i, t)|^3 + |\mu_A(x_i, t) - \eta_A(x_i, t)| \right)$
5.  $e_{GS-1}^t(A) = \frac{1}{n} \sum_{i=1}^n (1 - |\mu_A(x_i, t) - \eta_A(x_i, t)|)^{\frac{1 + \pi_A(x_i, t)}{2}}$
6.  $e_{Huang-1}^t(A) = 1 - \frac{1}{n} \sum_{i=1}^n \left| \mu_A(x_i, t)^2 - \eta_A(x_i, t)^2 \right|$
7.  $e_{Huang-2}^t(A) = 1 - \sqrt[p]{\frac{1}{n} \sum_{i=1}^n \left| \mu_A(x_i, t)^2 - \eta_A(x_i, t)^2 \right|^p}$
8.  $e_{Huang-3}^t(A) = 1 - \sqrt[p]{\frac{1}{n} \sum_{i=1}^n |\mu_A(x_i, t) - \eta_A(x_i, t)|^p}$

$$9. e^{t_{Huang-4}}(A) = \frac{1}{n} \sum_{i=1}^n \frac{1 - |\mu_A(x_i, t) - \eta_A(x_i, t)| + \pi_A(x_i, t)}{1 + |\mu_A(x_i, t) - \eta_A(x_i, t)| + \pi_A(x_i, t)}$$

$$10. e^{t_{Huang-5}}(A) =$$

$$\frac{1}{n} \sum_{i=1}^n 1 - |\mu_A(x_i, t) - \eta_A(x_i, t)| + \pi_A(x_i, t) |\mu_A(x_i, t) - \eta_A(x_i, t)|$$

$$11. e^{t_{Hung-1}}(A) = 1 - \sum_{i=1}^n \frac{1}{n} |\mu_A(x_i, t) - \eta_A(x_i, t)|$$

$$12. e^{t_{Hung-2}}(A) = 1 - \sqrt{\sum_{i=1}^n \frac{1}{n} |\mu_A(x_i, t) - \eta_A(x_i, t)|^2}$$

$$13. e^{t_{Ye-1}}(A) =$$

$$\frac{1}{n} \sum_{i=1}^n \left( \sin \left( \frac{\pi [1 + \mu_A(x_i, t) - \eta_A(x_i, t)]}{4} \right) + \sin \left( \frac{\pi [1 - \mu_A(x_i, t) + \eta_A(x_i, t)]}{4} - 1 \right) \frac{1}{\sqrt{2} - 1} \right)$$

$$14. e^{t_{Ye-2}}(A) =$$

$$\frac{1}{n} \sum_{i=1}^n \left( \cos \left( \frac{\pi [1 + \mu_A(x_i, t) - \eta_A(x_i, t)]}{4} \right) + \cos \left( \frac{\pi [1 - \mu_A(x_i, t) + \eta_A(x_i, t)]}{4} - 1 \right) \frac{1}{\sqrt{2} - 1} \right)$$

$$15. e^{t_{ZL-1}} = 1 - \frac{1}{b-a} \int_a^b |\mu_A(x, t) - \eta_A(x, t)| dx \text{ for } X = [a, b]$$

$$16. e^{t_{ZL-1}} = \frac{\int_a^b \mu_A(x, t) \wedge \eta_A(x, t) dx}{\int_a^b \mu_A(x, t) \vee \eta_A(x, t) dx} \text{ for } X = [a, b]$$

The other measure which is closely related with distance measure, similarity measure and entropy in intuitionistic fuzzy sets (and fuzzy set) is inclusion measure (named subsethood measure in some studies). Inclusion measure has been defined for intuitionistic fuzzy sets by Cornelis and Kerre in [9]. This concept is defined for temporal intuitionistic fuzzy sets as follows:

**Definition 3.5.** Let  $X$  be a non-empty set and  $T$  be a time set. If the mapping  $I_t : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow [0, 1]$  is satisfied following conditions for fixed  $t \in T$ , it is named temporal intuitionistic fuzzy inclusion measure on  $t \in T$ .

$$I1. I_t(\tilde{1}, \tilde{0}) = 0,$$

$$I2. I_t(A, B) = 1 \Leftrightarrow A \subseteq B$$

$$I3. I_t(C, A) \leq I_t(C, B) \text{ and } I_t(B, C) \leq I_t(A, C) \text{ when } A \subseteq B \text{ and } C \in TIFS^{(X, T)}$$

On the other hand, if the mapping  $I : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow [0, 1]$  is satisfied the conditions (I1, I2, I3) for each  $t \in T$ ,  $I$  is called overall intuitionistic fuzzy inclusion measure on  $TIFS^{(X, T)}$ . The mapping  $I : TIFS^{(X, T)} \times TIFS^{(X, T)} \rightarrow [0, 1]$  defined as  $I(A, B) = \sup_{t \in T} \{I_t(A, B)\}$  for all  $A, B \in TIFS^{(X, T)}$  and  $t \in T$  is an overall intuitionistic fuzzy inclusion measure.

The intuitionistic fuzzy inclusion measures which are defined in [44], [45] can be converted into the temporal intuitionistic fuzzy inclusion measures with some minor modifications as well as for other measures.

**Proposition 3.4.** *Let  $X = \{x_1, x_2, \dots, x_n\}$  be a non-empty set and  $T$  be a time set. Then the following mappings are temporal intuitionistic fuzzy inclusion measure on  $TIFS^{(X,T)}$ .*

$$1. I_{ZHXL-1}^t(A, B) = 1 - \frac{1}{2n} \sum_{i=1}^n \{ |\mu_A(x_i, t) - \min\{\mu_A(x_i, t), \mu_B(x_i, t)\}| + |\max\{\eta_A(x_i, t), \eta_B(x_i, t)\} - \eta_A(x_i, t)| \}$$

$$2. I_{ZHXL-2}^t(A, B) =$$

$$1 - \sqrt{\frac{1}{2n} \sum_{i=1}^n \{ |\mu_A(x_i, t) - \min\{\mu_A(x_i, t), \mu_B(x_i, t)\}|^2 + |\max\{\eta_A(x_i, t), \eta_B(x_i, t)\} - \eta_A(x_i, t)|^2 \}}$$

$$3. I_{ZDZS-3}^t(A, B) =$$

$$\begin{cases} 1 & , \quad \begin{matrix} \mu_B(x_i, t) = \mu_A(x_i, t) \\ \eta_B(x_i, t) = \eta_A(x_i, t) \end{matrix} \\ \sum_{i=1}^n \frac{1}{2} \left[ \frac{\mu_B(x_i, t) - \mu_A(x_i, t) + \eta_B(x_i, t) - \eta_A(x_i, t)}{|\mu_B(x_i, t) - \mu_A(x_i, t)| + |\eta_B(x_i, t) - \eta_A(x_i, t)|} \right] + 1 & , \quad \text{otherwise} \end{cases}$$

The following theorem which is given for the others measures can be given for temporal intuitionistic fuzzy inclusion measure and overall intuitionistic fuzzy inclusion measure as follows:

**Theorem 3.9.** *Let  $X$  be a non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be a finite time set. Let suppose that the mappings  $I_t : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  is a temporal intuitionistic fuzzy inclusion measure for each  $t \in T$  and  $f$  is a aggregation function without zero divisor. Then the mapping  $I : TIFS^{(X,T)} \times TIFS^{(X,T)} \rightarrow [0, 1]$  defined as*

$$I(A, B) = f(I_{t_1}(A), I_{t_2}(A), \dots, I_{t_n}(A))$$

for all  $A, B \in TIFS^{(X,T)}$  is a overall intuitionistic fuzzy inclusion measure.

The relationship between these measures are protected as described for the fuzzy and intuitionistic fuzzy sets. Some of these relationships can be generalized for the temporal intuitionistic fuzzy as follows. In this context, these relationships can be proved as in the studies which are given in the reference. Therefore, we will give the following theorems without proof (see more information: [3, 4, 5, 6, 8, 9, 10, 12, 13, 14, 15, 16, 17, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 33, 34, 35, 37, 38, 40, 41, 42, 43, 44, 45, etc.] ).

**Theorem 3.10.** *Let  $X$  be a non-empty set and  $T$  be time set. Let suppose that  $s^t$  is a temporal intuitionistic fuzzy similarity measure for  $t \in T$ . Then the mapping  $e^t : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $e^t(A) = s^t(A, \bar{A})$  for all  $A \in TIFS^{(X,T)}$  is a temporal intuitionistic fuzzy entropy*

**Theorem 3.11.** *Let  $X$  be a non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be finite time set. Let suppose that  $s^t$  is a temporal intuitionistic fuzzy similarity measure for*

$t \in T$  ve  $f$  is a aggregation function without zero divisor. Then the mapping  $e : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $e(A) = f(s^{t_1}(A, \bar{A}), s^{t_2}(A, \bar{A}), \dots, s^{t_n}(A, \bar{A}))$  for all  $A \in TIFS^{(X,T)}$  is a overall intuitionistic fuzzy entropy.

**Theorem 3.12.** Let  $X$  be a non-empty set and  $T$  be time set. Let suppose that  $d^t$  is a temporal intuitionistic fuzzy distance measure for  $t \in T$  and  $N_t$  is a temporal fuzzy non-filling negation. Then the mapping  $e^t : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $e^t(A) = N_t(d^t(A, \bar{A}))$  for all  $A \in TIFS^{(X,T)}$  is a temporal intuitionistic fuzzy entropy.

**Theorem 3.13.** Let  $X$  be a non-empty set and  $T = \{t_1, t_2, \dots, t_n\}$  be finite time set. Let suppose that  $s^t$  is a temporal intuitionistic fuzzy similarity measure for  $t \in T$ ,  $f$  is a aggregation function without zero divisor and  $N_t$  is a temporal fuzzy non-filling negation. The mapping  $e : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as

$$e(A) = f(N_{t_1}(d^{t_1}(A, \bar{A})), N_{t_2}(d^{t_2}(A, \bar{A})), \dots, N_{t_n}(d^{t_n}(A, \bar{A})))$$

for all  $A \in TIFS^{(X,T)}$  is a overall intuitionistic fuzzy entropy.

**Theorem 3.14.** Let  $X$  be a non-empty set and  $T$  be time set. Let suppose that  $I_t$  is a temporal intuitionistic fuzzy inclusion measure. Then the mapping  $E_I^t : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $E_I^t(A) = (I_t(A \cup \bar{A}, A \cap \bar{A}))$  for all  $A \in TIFS^{(X,T)}$  is a temporal intuitionistic fuzzy entropy.

**Theorem 3.15.** Let  $X$  be a non-empty set and  $T$  be time set. . Let suppose that  $*$  is a  $t$ - norm için  $I^t$  is a temporal intuitionistic fuzzy inclusion measure for  $t \in T$ . Then the mapping  $E_I^t : TIFS^{(X,T)} \rightarrow [0, 1]$  defined as  $S_I^t(A, B) = *(I_t(A, B), I_t(B, A))$  for all  $A, B \in TIFS^{(X,T)}$  is a temporal intuitionistic fuzzy similarity measure.

As seen from the above theorem, the basic relationship between distance measure, similarity measure, entropy and coverage measure are protected for temporal intuitionistic fuzzy sets as provided in for fuzzy and intuitionistic fuzzy sets. Furthermore, these measures can be separate two groups which are named as temporal and overall for temporal intuitionistic fuzzy sets. It can be done the first instant evaluation and later general evaluation for all data with time-varying nature by this method. Thus, the optimal results can be obtained by various way in the application.

## REFERENCES

- [1] Atanassov K. T., 1986. Intuitionistic fuzzy sets, Fuzzy Sets and Systems, 20 , 87-96.
- [2] Atanassov K. T., 1991. Temporal intuitionistic fuzzy sets, Comptes Rendus de l'Academie Bulgare, Vol. 7, 5-7.
- [3] Atanassov K. T., 2012. On Intuitionistic Fuzzy Sets Theory, Springer, Berlin.
- [4] B.Y. Cao, et al, 2010. Relationships among fuzzy entropy, similarity measure and distance measure of intuitionistic fuzzy sets. Fuzzy Information and Engineering, 78: 539-548.
- [5] Baccour, L., Alimi A. M., John, R. I., 2011. Relationship between intuitionistic fuzzy similarity measures. IEEE Int. Conf. on Fuzzy Syst.: 971-975
- [6] Beg I., Ashraf S., 2009. Similarity measures for fuzzy sets. Appl. And Comput. Math., 8(2): 192-202.
- [7] Chen L., Tu C., 2015. Time-validating-based Atanassov's intuitionistic fuzzy decision-making, IEEE Transactions on Fuzzy Systems, 23(4), 743 - 756.
- [8] Chen, S.M., 1997. Similarity measure between vague sets and between elements. IEEE Trans. Syst. Man Cybern., 27 (1): 153-158.

- [9] Cornelis C., Kerre, E., 2003. Inclusion measure in intuitionistic fuzzy set theory. *Sym. and Quant. Appr. to Reas. with Unc.* Springer Berlin Heidelberg: 345-356.
- [10] Deng, G., Jiang Y., Fu, J., 2015. Monotonic similarity measures between intuitionistic fuzzy sets and their relationship with entropy and inclusion measure. *Information Sciences*, 316: 348-369.
- [11] Du, W. S., Hu, B. Q., 2015. Aggregation distance measure and its induced similarity measure between intuitionistic fuzzy sets. *Pattern Recog. Lett.*, 60: 65-71.
- [12] Fan, J. L., Ma, Y. L., 2002. Some new fuzzy entropy formulas, *Fuzzy sets and Systems*, 128: 277-284
- [13] Fan, J., Xie, W., 1999. Some notes on similarity measure and proximity measure. *Fuzzy Sets and Systems*, 101: 403-412.
- [14] Fan, J., Xie, W.X., 1999. Distance measure and induced fuzzy entropy. *Fuzzy Sets and Systems*, 104: 305-314.
- [15] Fan, J., Xie, W.X., Pei, J., 1999. Subsethood measure: new definitions. *Fuzzy Sets and Systems*, 106: 201-209.
- [16] Grzegorzewski P., 2004. Distances between intuitionistic fuzzy sets and/or interval-valued fuzzy sets based on Hausdorff metric, *Fuzzy Sets and Systems*, 148, , 319-328.
- [17] Grzegorzewski, P., Mrowka, E., 2004. Subsethood measure for intuitionistic fuzzy sets. *IEEE International Conference on Fuzzy Systems*, 1: 139 - 142.
- [18] Guha, D., Chakraborty, D., 2010. A theoretical development of distance measure for intuitionistic fuzzy numbers, *Int. J. of Math. and Math. Sci.*: 1-25.
- [19] Guo, K., Song, Q., 2014. On the entropy for Atanassov's intuitionistic fuzzy sets: an interpretation from the perspective of amount of knowledge. *Applied Soft Computing*, 24: 328-340.
- [20] Huang G., Liu Y., Wang X., 2005. New Distances between intuitionistic fuzzy sets, in *Proceedings of fourth international conference on machine learning and cybernetics*, Guangzhou, 18-21.
- [21] Huang, G. S., 2007. A new fuzzy entropy for intuitionistic fuzzy sets. . *Fourth International Conference on Fuzzy Systems and Knowledge Discovery*, 2007. *FSKD 2007*: 57 - 61.
- [22] Huang, G., Liu, Y., Wang, X., 2005. New Distances between intuitionistic fuzzy sets. *Proceedings of fourth international conference on machine learning and cybernetics*, Guangzhou. 18-21.
- [23] Hung, W. L., Yang, M. S., 2004. Similarity measures of intuitionistic fuzzy sets based on Hausdorff distance, *Pattern Recognition Letters*, 23: 1603-1611.
- [24] Hung, W. L., Yang, M. S., 2006. Fuzzy entropy on intuitionistic fuzzy sets. *Int. Jour. Of Intell. Syst.* 21: 443-451.
- [25] Hwang, C.H., Yang, M.S., 2008. On entropy of fuzzy sets. *International Journal of Uncertainty*, 16(4): 519-527.
- [26] Intrapai boon P., 2014. New similarity measures for intuitionistic fuzzy sets, *Applied Math. Sci.*, 8(45): 2239-2250.
- [27] Julian, P., Hung K. C., Lin, S. J., 2012. On the Mitchell similarity measure and its application to pattern recognition. *Pattern Recog. Lett.*, 33: 1219-1223.
- [28] Lee, S. H., Kim, J. M., Choi, Y. K., 2006. Similarity measure construction using fuzzy entropy and distance measure. *Computational Intelligence*, Springer Berlin Heidelberg: 952-958.
- [29] Li, J., Deng, G., Li, H., Zeng, W., 2012. The relationship between similarity measure and entropy of intuitionistic fuzzy sets. *Information Sciences*, 188: 314-321.
- [30] Li, Y., Qin, K., He, X., 2013. Relations among similarity measures, subsethood measure and fuzzy entropy. *International Journal of Computational Intelligence Systems*, 6(3): 411-422.
- [31] Liu, X., 1992. Entropy, distance measure and similarity measure of fuzzy sets and their relations. *Fuzzy and Systems*, 52: 305-318.
- [32] Parvathi, R., S. P. Geetha, 2009. A note on properties of temporal intuitionistic fuzzy sets, *Notes on Intuitionistic Fuzzy Sets*, 15(1), 42-48.
- [33] Szmidt E. , Kacprzyk J., 2000. Distances between intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 114, 505-518.
- [34] Szmidt E., 2014. Distances and similarities in intuitionistic fuzzy sets, Springer, Switzerland,
- [35] Szmidt, E., Kacprzyk, J., 2006. New measures of entropy for intuitionistic fuzzy sets. *Notes on IFS*, 11 (2): 12-20.
- [36] Vlachos, K., Sergiadis, G.D., 2007. Intuitionistic fuzzy image processing, *StudFuzz*, 210: 383-414.

- [37] Yang Y., Chiclana F., 2012. Consistency of 2D and 3D distances of intuitionistic fuzzy sets, Expert System with Applications, 39, 8665-8670.
- [38] Ye, J., 2010. Two effective measures of intuitionistic fuzzy entropy. Computing, 87(1): 55-62.
- [39] Yılmaz S., Çuvalcıoğlu G., On level operators for temporal intuitionistic fuzzy sets, Notes on Intuitionistic Fuzzy Sets 20(2), 2014, 6-15
- [40] Young, V. R., 1996. Fuzzy subsethood. Fuzzy Sets and Systems, 77: 371-384.
- [41] Zeng W., Guo, P., 2008. Normalized distance, similarity measure, inclusion measure and entropy of interval-valued fuzzy sets and their relationship. Information Sci., 178: 1334-1342.
- [42] Zeng, W., Feng, Q., Li, H. X., 2006. Relationship between inclusion measure and entropy of fuzzy sets. Rough Sets and Knowledge Technology: 333-340.
- [43] Zhang Q., Huang, Y., Xing, H., Liu, F., 2012. Distance measure, information entropy and inclusion measure of intuitionistic fuzzy sets and their relations. Int. Jour. of Adv. in Comp. Tech., 4(15): 480-487.
- [44] Zhang, H., Dong, M., Zhang, W., Song, X., 2007. Inclusion measure and similarity measure of intuitionistic and interval-valued fuzzy sets. Proceedings of the 2007 International Conference on Intelligent Systems and Knowledge Engineering (ISKE2007).
- [45] Zhang, Q., Liu F., Wu, L., Luo, S., 2012. Information entropy, similarity measures and inclusion measure of intuitionistic fuzzy sets. Inf. Comp. and App., 307: 392-398.

YUZUNCU YIL UNIVERSITY, DEPT. OF ELECTRONIC AND COMMUNICATION TECHNOLOGIES, ER-CIS 65080, VAN, TURKEY.

*E-mail address:* fatihkutlu@yyu.edu.tr

YUZUNCU YIL UNIVERSITY, DEPARTMENT OF ELECTRICAL-ELECTRONIC ENGINEERING, TUSBA, 65090, VAN, TURKEY.

*E-mail address:* oatan@yyu.edu.tr

YUZUNCU YIL UNIVERSITY, FACULTY OF EDUCATION, TUSBA, 65090, VAN, TURKEY.

*E-mail address:* tbilgin@yyu.edu.tr



## ON A SOCIAL ECONOMIC MODEL

OLGUN CABRI AND KHANLAR R. MAMEDOV

ABSTRACT. Distribution of saving for a family sets on a region satisfies Kolmogorov equation given by

$$\frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} ((c + F)u) + \frac{1}{2} \frac{\partial^2}{\partial x^2} (bu) + f$$

where  $u = u(x, t)$  is density distribution of family saving. Boundary condition defined by distribution of minimum saving and total family saving are considered for the model. By the separation of variables method, eigenvalues and eigenfunctions of problem are obtained and solution is written. In addition, numerical methods are applied to problem and errors of numerical methods are presented.

Received: 08–August–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Nonlocal boundary conditions are dealt with some wave, diffusion and any other physical equations [Cannon, Van der Hoek, Ionkin, Kamynin, etc...] Generally these type of problems are solved by numeric methods or reducing point boundary conditions. In this study we consider a family saving model used in economy. This problem is expressed with diffusion equation with integral boundary conditions.

Suppose that  $x(t)$  shows saving of a family at time  $t$  and satisfy the differential equation

$$(1.1) \quad dx = F(x, t) dt + G(x, t) dX, \quad G \geq 0$$

where  $X$  is Markov process,  $F(x, t)$  denotes rate of change of the family saving and  $G(x, t) dX$  denotes random change of family income.

For a family set let us assume that equation (1.1) describes the saving of all families by ignoring the dynamic of individual family saving. The density distribution of the saving of families  $u(x, t)$  satisfies

$$(1.2) \quad \frac{\partial u}{\partial t} = -\frac{\partial}{\partial x} ((c(x, t) + F(x, t)) u) + \frac{1}{2} \frac{\partial^2}{\partial x^2} (b(x, t)u) + f(x, t)$$

with initial condition

$$(1.3) \quad u(x, 0) = \varphi(x), \quad 0 \leq x \leq l$$

and boundary conditions

$$(1.4) \quad u(0, t) = 0$$

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
*Key words and phrases.* Nonlocal Boundary Condition, Family Saving Model, Method of Lines Method, Crank Nicolson Method.

$$(1.5) \quad \int_0^l x u(x, t) dx = K(t), \quad t \geq 0$$

where  $c(x, t)$ ,  $b(x, t)$ ,  $K(t)$  are continuously differentiable functions.  $K(t)$  in (1.5) describes total amount of family saving in  $[0, l]$  [5].

We will consider special case of problem (1.1)-(1.5) on region  $D = (0 < t < \infty) \times (0 < x < l)$

$$(1.6) \quad \frac{\partial u}{\partial t} = a^2 \frac{\partial^2 u}{\partial x^2} + f(x, t)$$

$$(1.7) \quad u(x, 0) = \varphi(x), \quad 0 \leq x \leq l$$

$$(1.8) \quad u(0, t) = 0$$

$$(1.9) \quad \int_0^l x u(x, t) dx = K(t), \quad 0 \leq x \leq l$$

where  $f(x, t)$ ,  $K(t)$ ,  $\varphi(x)$  are continuously differentiable function on region  $D$  and  $a$  is given constant. Compatibility conditions of this problem is  $\int_0^l \varphi(x) dx = K(0)$ .

In order to obtain classical solution of problem (1.6)-(1.9), we transform boundary conditions into homegenous ones by the transformation

$$(1.10) \quad u(x, t) = v(x, t) + 3K(t)x$$

Carrying out substitution (1.10) in (1.1)-(1.5) problem gives

$$(1.11) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2} + F(x, t)$$

$$(1.12) \quad v(x, 0) = \psi(x)$$

$$(1.13) \quad v(0, t) = 0$$

$$(1.14) \quad \int_0^l x v(x, t) dx = 0$$

where  $F(x, t) = f(x, t) - 3K'(t)x$  and  $\psi(x) = \varphi(x) - 3K(0)x$

This problem has homegenous boundary conditions. Due to linearity, problem can split into two auxilary problem:

i)

$$(1.15) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2}$$

$$(1.16) \quad v(x, 0) = \psi(x)$$

$$(1.17) \quad v(0, t) = 0$$

$$(1.18) \quad \int_0^1 xv(x,t)dx = 0$$

ii)

$$(1.19) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2} + F(x,t)$$

$$(1.20) \quad v(x,0) = 0$$

$$(1.21) \quad v(x,t) = 0$$

$$(1.22) \quad \int_0^1 xv(x,t)dx = 0$$

Integrating both sides of (1.15) with respect to  $x$  from 0 to 1 and using integration by parts, integral boundary condition turns into Neumann boundary condition

$$v_x(1,t) - v(1,t) = 0$$

Thus problem (1.15)-(1.18) becomes

$$(1.23) \quad \frac{\partial v}{\partial t} = a^2 \frac{\partial^2 v}{\partial x^2}$$

$$(1.24) \quad v(x,0) = \psi(x)$$

$$(1.25) \quad v(0,t) = 0$$

$$(1.26) \quad v_x(1,t) - v(1,t) = 0$$

By the Fourier method, Sturm Liouville problem and ODE are ,respectively, obtained as

$$(1.27) \quad X''(x) + \lambda X(x) = 0$$

$$(1.28) \quad X(0) = 0$$

$$(1.29) \quad X'(1) - X(1) = 0$$

and

$$(1.30) \quad T'(t) + \lambda a^2 T(t) = 0$$

Sturm Liouville problem (1.27)-(1.29)is self adjoint and boundary conditions are regular, moreover strongly regular. Then eigenfunctions of Sturm Liouville problem are Riesz basis on  $L^2[0, 1]$ .

Charecteristic equation of Sturm Liouville problem is

$$\tan k = k$$

Thus the problem has the eigenvalues  $\lambda_n$  ,  $n = 0, 1, 2..$  such that  $\lambda_0 = 0$  and by using Langrange-Burmman formula

$$k_n = \frac{(2n+1)\pi}{2} - \left(\frac{(2n+1)\pi}{2}\right)^{-1} - \frac{2}{3} \left(\frac{(2n+1)\pi}{2}\right)^{-3} - \frac{13}{15} \left(\frac{(2n+1)\pi}{2}\right)^{-5} - \frac{146}{105} \left(\frac{(2n+1)\pi}{2}\right)^{-7} + O\left(\frac{1}{n^9}\right)$$

where  $\sqrt{\lambda_n} = k_n$ .

Corresponding eigenfunctions are obtained by

$$\begin{aligned} X_0(x) &= cx \\ X_n(x) &= \sin(k_n x), n = 1, 2, \dots \end{aligned}$$

Hence solution of problem (1.23)-(1.26) is

$$v_1(x, t) = A_0 x + \sum_{n=1}^{\infty} A_n e^{-a^2 k_n^2 t} \sin(k_n x)$$

where

$$\begin{aligned} A_0 &= 3 \int_0^1 x \psi(x) dx \\ A_n &= \frac{1}{\frac{1}{2} - \frac{\sin(2k_n)}{4k_n}} \int_0^1 \psi(x) \sin(k_n x) dx, n = 1, 2, \dots \end{aligned}$$

Solution of problem (1.19)-(1.22) can easily obtained by

$$v_2(x, t) = \left[ \int_0^t F_0(\tau) d\tau \right] x + \sum_{n=1}^{\infty} \left[ \int_0^t F_n(\tau) e^{-a^2 k_n^2 (t-\tau)} d\tau \right] \sin(k_n(x))$$

where

$$\begin{aligned} F_n(\tau) &= \int_0^1 F(x, \tau) \sin(k_n x) dx \\ F_0(\tau) &= \int_0^1 F(x, \tau) x dx \end{aligned}$$

## 2. NUMERICAL SOLUTION

For the numerical solution of problem, we will use the Method of Lines method and Crank-Nicolson method presented in [8],[9] respectively. In both methods, Simpson's rule is used to approximate the integral in (1.18) numerically. We display here a few of numerical results.

Example 1..

$$\begin{aligned} \frac{\partial u}{\partial t} &= \frac{\partial^2 u}{\partial x^2} + (x^2 - 2)e^t \\ u(x, 0) &= x^2 \\ u(0, t) &= 0 \\ \int_0^1 x u(x, t) dx &= \frac{e^t}{4} \end{aligned}$$

Exact solution is  $u(x, t) = x^2 e^t$ . The computed results at various spatial lengths are shown in Table 2. This table exhibits the absolute relative error results for  $u(0.5, 0.5)$ .

Example 2:

$$\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{-2(x^2 + t + 1)}{(t + 1)^3}$$

TABLE 1. Relative Error at  $u(0.5,0.5)$  in Example 1

Spatial Length	MOL Method	Crank-Nicolson Method		
h=0.1	1.6291E-5	4.8730E-5		
h=0.05	2.553E-6	1.2398E-5		
h=0.025	3.6733E-7	3.1067E-6		
h=0.0125	4.8533E-8	7.7695E-7		

TABLE 2. Relative Error at  $u(0.5,0.5)$  in Example 2

Spatial Length	MOL Method	Crank-Nicolson Method		
h=0.1	1.4572E-4	3.7936E-4		
h=0.05	2.2868E-5	1.0927E-4		
h=0.025	3.3245E-6	2.7647E-5		
h=0.0125	4.468E-7	6.9193E-6		

$$u(x, 0) = x^2$$

$$u(0, t) = 0$$

$$\int_0^1 xu(x, t)dx = \frac{1}{4(t+1)^2}$$

Exact solution is  $u(x, t) = \frac{x^2}{(t+1)^2}$ . The computed results at various spatial lengths are shown in Table 2. This table exhibits the absolute relative error results for  $u(0.5, 0.5)$ .

We studied a special case of family saving model which is diffusion equation with nonlocal boundary condition. Analytic solution of this problem is obtained. Moreover by applying the Method of Lines method [8] and Crank Nicolson [9], numerical solution of problem is found.

#### REFERENCES

- [1] Cannon, J.R., *The solution of the heat equation subject to specification of energy*, Quarterly of Applied Mathematics, Vol. **21**, 155-160, 1963.
- [2] Cannon, J. R. ve Van Der Hoek, J., *An implicit finite difference scheme for the diffusion of mass in a portion of the domain*, Numerical Solutions of Partial Differential Equations (J. Noye, Ed), 527-539. 1982
- [3] Dehghan, M. , *On the numerical solution of the diffusion equation with a nonlocal boundary condition*, Mathematical Problems in Engineering, 2003(2), 81-92, 2003
- [4] Ekolin, G., *Finite difference methods for a nonlocal boundary value problem for the heat equation*, BIT Numerical Mathematics, 1991, 31.2: 245-261.
- [5] Erofeenko, V.T. ve Kozlovski, I.Y., *Equation Partial Differential Equation and Mathematical Models in Economic*, URRS, 2011
- [6] Ionkin N. I, *Solution of a boundary-value problem in heat conduction with a nonclassical boundary condition*, Diff. Eqs., Vol. 13, No. 2,1977, pp. 294-304.
- [7] Kamynin, L. I., *A boundary value problem in the theory of heat conduction with a nonclassical boundary condition*, USSR Computational Mathematics and Mathematical Physics, 1964, 4.6: 33-59.
- [8] Rehmana, M. A. ve TaJ, M. S., *A Numerical technique for heat equation subject to integral specification*, Sci.Ä°nt(Lahore),24(1),21-26,2011
- [9] Shingmin W. ve Lin, Y. *A numerical method for the diffusion equation with nonlocal boundary specifications*, International Journal of Engineering Science, 1990, 28.6: 543-546.

ARTVIN CORUH UNIVERSITY  
*E-mail address:* `olguncabri@artvin.edu.tr`

MERSIN UNIVERSITY  
*E-mail address:* `hanlarm@mersin.edu.tr`

## SOME RESULTS ON $S_{\alpha,\beta}$ AND $T_{\alpha,\beta}$ INTUITIONISTIC FUZZY MODAL OPERATORS

GÖKHAN ÇUVALCIOĞLU, KRASSIMIR T. ATANASSOV,  
AND SINEM TARSUSLU(YILMAZ)

ABSTRACT. In 1999, first Intuitionistic Fuzzy Modal Operators introduced in[2]. Expansion of these operators and new operators defined by different authors[3, 5, 6, 7, 8, 9]. Characteristics of these operators has been studied by several researchers.

In this study, we obtained new results on modal operators which are called  $S_{\alpha,\beta}$  and  $T_{\alpha,\beta}$ .

Received: 25–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

The concept of Intuitionistic fuzzy sets was introduced by Atanassov in 1986 [1], form an extension of fuzzy sets[10] by expanding the truth value set to the lattice  $[0, 1] \times [0, 1]$ .

Intuitionistic fuzzy modal operators defined firstly by Atanassov[1, 2]. Then several extensions of these operators introduced by different authors[2, 8, 5, 6]. Some algebraic and characteristic properties of these operators were studied by several authors.

**Definition 1.1.** [1] An intuitionistic fuzzy set (shortly IFS) on a set  $X$  is an object of the form

$$A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle : x \in X \}$$

where  $\mu_A(x), (\mu_A : X \rightarrow [0, 1])$  is called the “degree of membership of  $x$  in  $A$ ”,  $\nu_A(x), (\nu_A : X \rightarrow [0, 1])$  is called the “degree of non- membership of  $x$  in  $A$ ”, and where  $\mu_A$  and  $\nu_A$  satisfy the following condition:

$$\mu_A(x) + \nu_A(x) \leq 1, \text{ for all } x \in X.$$

The class of intuitionistic fuzzy sets on  $X$  is denoted by  $IFS(X)$ .

The hesitation degree of  $x$  is defined by  $\pi_A(x) = 1 - \mu_A(x) - \nu_A(x)$

**Definition 1.2.** [1] An IFS  $A$  is said to be contained in an IFS  $B$  (notation  $A \sqsubseteq B$ ) if and only if, for all  $x \in X : \mu_A(x) \leq \mu_B(x)$  and  $\nu_A(x) \geq \nu_B(x)$ .

It is clear that  $A = B$  if and only if  $A \sqsubseteq B$  and  $B \sqsubseteq A$ .

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 03E72,47S40.

*Key words and phrases.* Intuitionistic fuzzy sets, Intuitionistic fuzzy modal operators.

**Definition 1.3.** [1] Let  $A \in IFS$  and let  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\}$  then the above set is called the complement of  $A$

$$A^c = \{\langle x, \nu_A(x), \mu_A(x) \rangle : x \in X\}$$

The intersection and the union of two IFSs  $A$  and  $B$  on  $X$  is defined by

$$A \sqcap B = \{\langle x, \mu_A(x) \wedge \mu_B(x), \nu_A(x) \vee \nu_B(x) \rangle : x \in X\}$$

$$A \sqcup B = \{\langle x, \mu_A(x) \vee \mu_B(x), \nu_A(x) \wedge \nu_B(x) \rangle : x \in X\}$$

The notion of Second Type Intuitionistic Fuzzy Modal Operators was firstly introduced by Atanassov as following:

**Definition 1.4.** [1] Let  $X$  be universal and  $A \in IFS(X)$  then

$$(1) \square(A) = \{\langle x, \mu_A(x), 1 - \mu_A(x) \rangle : x \in X\}$$

$$(2) \diamond(A) = \{\langle x, 1 - \nu_A(x), \nu_A(x) \rangle : x \in X\}$$

**Definition 1.5.** [2] Let  $X$  be universal and  $A \in IFS(X)$ ,  $\alpha \in [0, 1]$  then

$$D_\alpha(A) = \{\langle x, \mu_A(x) + \alpha\pi_A(x), \nu_A(x) + (1 - \alpha)\pi_A(x) \rangle : x \in X\}$$

**Definition 1.6.** [2] Let  $X$  be universal and  $A \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$  and  $\alpha + \beta \leq 1$  then

$$F_{\alpha, \beta}(A) = \{\langle x, \mu_A(x) + \alpha\pi_A(x), \nu_A(x) + \beta\pi_A(x) \rangle : x \in X\}$$

**Definition 1.7.** [2] Let  $X$  be universal and  $A \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$  then

$$G_{\alpha, \beta}(A) = \{\langle x, \alpha\mu_A(x), \beta\nu_A(x) \rangle : x \in X\}$$

**Definition 1.8.** [2] Let  $X$  be universal and  $A \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$  then

$$(1) H_{\alpha, \beta}(A) = \{\langle x, \alpha\mu_A(x), \nu_A(x) + \beta\pi_A(x) \rangle : x \in X\}$$

$$(2) H_{\alpha, \beta}^*(A) = \{\langle x, \alpha\mu_A(x), \nu_A(x) + \beta(1 - \alpha\mu_A(x) - \nu_A(x)) \rangle : x \in X\}$$

$$(3) J_{\alpha, \beta}(A) = \{\langle x, \mu_A(x) + \alpha\pi_A(x), \beta\nu_A(x) \rangle : x \in X\}$$

$$(4) J_{\alpha, \beta}^*(A) = \{\langle x, \mu_A(x) + \alpha(1 - \mu_A(x) - \beta\nu_A(x)), \beta\nu_A(x) \rangle : x \in X\}$$

The simplest One Type Intuitionistic Fuzzy Modal Operators defined in 1999.

**Definition 1.9.** [2] Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$ .

$$(1) \boxplus A = \left\{ \left\langle x, \frac{\mu_A(x)}{2}, \frac{\nu_A(x)+1}{2} \right\rangle : x \in X \right\}$$

$$(2) \boxtimes A = \left\{ \left\langle x, \frac{\mu_A(x)+1}{2}, \frac{\nu_A(x)}{2} \right\rangle : x \in X \right\}$$

After this definition, in 2001, the extension of these operators were defined as following:

**Definition 1.10.** [3] Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$ .

$$(1) \boxplus_\alpha A = \{\langle x, \alpha\mu_A(x), \alpha\nu_A(x) + 1 - \alpha \rangle : x \in X\}$$

$$(2) \boxtimes_\alpha A = \{\langle x, \alpha\mu_A(x) + 1 - \alpha, \alpha\nu_A(x) \rangle : x \in X\}$$

The operators  $\boxplus_\alpha$  and  $\boxtimes_\alpha$  are the extensions of the operators  $\boxplus$ ,  $\boxtimes$ , resp.

In 2004, Dencheva introduced the second extension of  $\boxplus_\alpha$  and  $\boxtimes_\alpha$ .

**Definition 1.11.** [8] Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$ .



- (1)  $\boxplus_{\alpha,\beta}A = \{\langle x, \alpha\mu_A(x), \alpha\nu_A(x) + \beta \rangle : x \in X\}$  where  $\alpha + \beta \in [0, 1]$ .
- (2)  $\boxtimes_{\alpha,\beta}A = \{\langle x, \alpha\mu_A(x) + \beta, \alpha\nu_A(x) \rangle : x \in X\}$  where  $\alpha + \beta \in [0, 1]$ .

In 2006, the third extension of the above operators was studied by author . He defined the following operators;

**Definition 1.12.** [3]Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ .

- (1)  $\boxplus_{\alpha,\beta,\gamma}(A) = \{\langle x, \alpha\mu_A(x), \beta\nu_A(x) + \gamma \rangle : x \in X\}$   
where  $\alpha, \beta, \gamma \in [0, 1], \max\{\alpha, \beta\} + \gamma \leq 1$ .
- (2)  $\boxtimes_{\alpha,\beta,\gamma}(A) = \{\langle x, \alpha\mu_A(x) + \gamma, \beta\nu_A(x) \rangle : x \in X\}$   
where  $\alpha, \beta, \gamma \in [0, 1], \max\{\alpha, \beta\} + \gamma \leq 1$ .

In 2007, author[5] defined a new operator named  $E_{\alpha,\beta}$  and studied some of its properties. This operator as following:

**Definition 1.13.** [5]Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta \in [0, 1]$ . We define the following operator:

$$E_{\alpha,\beta}(A) = \{\langle x, \beta(\alpha\mu_A(x) + 1 - \alpha), \alpha(\beta\nu_A(x) + 1 - \beta) \rangle : x \in X\}$$

At the same year, Atanassov introduced the operator  $\boxminus_{\alpha,\beta,\gamma,\delta}$  which is a natural extension of all these operators in [3].

**Definition 1.14.** [3]Let  $X$  be a set,  $A \in IFS(X)$ ,  $\alpha, \beta, \gamma, \delta \in [0, 1]$  such that

$$\max(\alpha, \beta) + \gamma + \delta \leq 1$$

then the operator  $\boxminus_{\alpha,\beta,\gamma,\delta}$  defined by

$$\boxminus_{\alpha,\beta,\gamma,\delta}(A) = \{\langle x, \alpha\mu_A(x) + \gamma, \beta\nu_A(x) + \delta \rangle : x \in X\}$$

In 2008, most general operator  $\odot_{\alpha,\beta,\gamma,\delta,\varepsilon,\zeta}$  defined as following:

**Definition 1.15.** [3]Let  $X$  be a set,  $A \in IFS(X)$ ,  $\alpha, \beta, \gamma, \delta, \varepsilon, \zeta \in [0, 1]$  such that

$$\max(\alpha - \zeta, \beta - \varepsilon) + \gamma + \delta \leq 1$$

and

$$\min(\alpha - \zeta, \beta - \varepsilon) + \gamma + \delta \geq 0$$

then the operator  $\odot_{\alpha,\beta,\gamma,\delta,\varepsilon,\zeta}$  defined by

$$\odot_{\alpha,\beta,\gamma,\delta,\varepsilon,\zeta}(A) = \{\langle x, \alpha\mu_A(x) - \varepsilon\nu_A(x) + \gamma, \beta\nu_A(x) - \zeta\mu_A(x) + \delta \rangle : x \in X\}$$

In 2010, Çuvalcıođlu[6] defined a new operator which is a generalization of  $E_{\alpha,\beta}$ .

**Definition 1.16.** [6]Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta, \omega \in [0, 1]$  then

$$Z_{\alpha,\beta}^{\omega}(A) = \{\langle x, \beta(\alpha\mu_A(x) + \omega - \omega.\alpha), \alpha(\beta\nu_A(x) + \omega - \omega.\beta) \rangle : x \in X\}$$

**Definition 1.17.** [6]Let  $X$  be a set and  $A = \{\langle x, \mu_A(x), \nu_A(x) \rangle : x \in X\} \in IFS(X)$ ,  $\alpha, \beta, \omega, \theta \in [0, 1]$  then

$$Z_{\alpha,\beta}^{\omega,\theta}(A) = \{\langle x, \beta(\alpha\mu_A(x) + \omega - \omega.\alpha), \alpha(\beta\nu_A(x) + \theta - \theta.\beta) \rangle : x \in X\}$$

The operator  $Z_{\alpha,\beta}^{\omega,\theta}$  is a generalization of  $Z_{\alpha,\beta}^{\omega}$ , and also,  $E_{\alpha,\beta}, \boxplus_{\alpha,\beta}, \boxtimes_{\alpha,\beta}$ .

Uni-type intuitionistic fuzzy modal operators introduced by author as following;

**Definition 1.18.** [7]Let  $X$  be a universal,  $A \in IFS(X)$  and  $\alpha, \beta, \omega \in [0, 1]$  then

- (1)  $\boxplus_{\alpha,\beta}^{\omega}(A) = \{ \langle x, \beta(\mu_A(x) + (1 - \alpha)\nu_A(x)), \alpha(\beta\nu_A(x) + \omega - \omega\beta) \rangle : x \in X \}$
- (2)  $\boxtimes_{\alpha,\beta}^{\omega}(A) = \{ \langle x, \beta(\alpha\mu_A(x) + \omega - \omega\alpha), \alpha((1 - \beta)\mu_A(x) + \nu_A(x)) \rangle : x \in X \}$

**Definition 1.19.** [7] Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \omega, \theta \in [0, 1]$  then

$$E_{\alpha,\beta}^{\omega,\theta}(A) = \left\{ \left\langle \begin{array}{l} x, \beta((1 - (1 - \alpha)(1 - \theta))\mu_A(x) + (1 - \alpha)\theta\nu_A(x) + (1 - \alpha)(1 - \theta)\omega), \\ \alpha((1 - \beta)\theta\mu_A(x) + (1 - (1 - \beta)(1 - \theta))\nu_A(x) + (1 - \beta)(1 - \theta)\omega) \end{array} \right\rangle : x \in X \right\}$$

**Definition 1.20.** [7] Let  $X$  be a set,  $A \in IFS(X)$  and  $\alpha, \beta \in [0, 1]$  then

- (1)  $B_{\alpha,\beta}(A) = \{ \langle x, \beta(\mu_A(x) + (1 - \alpha)\nu_A(x)), \alpha((1 - \beta)\mu_A(x) + \nu_A(x)) \rangle : x \in X \}$
- (2)  $\boxminus_{\alpha,\beta}(A) = \{ \langle x, \beta(\mu_A(x) + (1 - \beta)\nu_A(x)), \alpha((1 - \alpha)\mu_A(x) + \nu_A(x)) \rangle : x \in X \}$

In 2014, new one type intuitionistic fuzzy modal operators were defined in [9].

**Definition 1.21.** [9] Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \omega \in [0, 1]$  and  $\alpha + \beta \leq 1$

- (1)  $L_{\alpha,\beta}^{\omega}(A) = \{ \langle x, \alpha\mu_A(x) + \omega(1 - \alpha), \alpha(1 - \beta)\nu_A(x) + \alpha\beta(1 - \omega) \rangle : x \in X \}$
- (2)  $K_{\alpha,\beta}^{\omega}(A) = \{ \langle x, \alpha(1 - \beta)\mu_A(x) + \alpha\beta(1 - \omega), \alpha\nu_A(x) + \omega(1 - \alpha) \rangle : x \in X \}$

As above, we get the following diagram;




FIGURE 1

The intuitionistic fuzzy modal operator, represented by  $\otimes_{\alpha,\beta,\gamma,\delta}$ , introduced in 2014 as following;

**Definition 1.22.** [4] Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \gamma, \delta \in [0, 1]$  and  $\alpha + \beta \leq 1, \gamma + \delta \leq 1$  then

$$\otimes_{\alpha,\beta,\gamma,\delta}(A) = \{ \langle x, \alpha\mu_A(x) + \gamma\nu_A(x), \beta\mu_A(x) + \delta\nu_A(x) \rangle \}$$

2. SOME PROPERTIES OF NEW INTUITIONISTIC FUZZY MODAL OPERATORS

**Definition 2.1.** Let  $X$  be a set and  $A \in IFS(X)$ ,  $\alpha, \beta, \alpha + \beta \in [0, 1]$ .

- (1)  $T_{\alpha,\beta}(A) = \{ \langle x, \beta(\mu_A(x) + (1 - \alpha)\nu_A(x) + \alpha), \alpha(\nu_A(x) + (1 - \beta)\mu_A(x)) \rangle : x \in X \}$  where  $\alpha + \beta \in [0, 1]$ .
- (2)  $S_{\alpha,\beta}(A) = \{ \langle x, \alpha(\mu_A(x) + (1 - \beta)\nu_A(x)), \beta(\nu_A(x) + (1 - \alpha)\mu_A(x) + \alpha) \rangle : x \in X \}$  where  $\alpha + \beta \in [0, 1]$ .

It is clear that;

$$\begin{aligned} \beta(\mu_A(x) + (1 - \alpha)\nu_A(x) + \alpha) + \alpha(\nu_A(x) + (1 - \beta)\mu_A(x)) &= (\mu_A(x) + \nu_A(x))(\alpha + \beta - \alpha\beta) + \alpha\beta \\ &\leq \alpha + \beta - \alpha\beta + \alpha\beta \leq 1 \end{aligned}$$

**Theorem 2.1.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $\alpha, \beta, \alpha + \beta \in [0, 1]$  then  $T_{\alpha,\beta}(A)^c = S_{\alpha,\beta}(A^c)$ .

*Proof.* It is clear from definition. □

**Proposition 2.1.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $\alpha, \beta, \alpha + \beta \in [0, 1]$  then

- (1)  $T_{\beta,\alpha}(A)^c \sqsubseteq T_{\alpha,\beta}(A^c)$




FIGURE 2

$$(2) S_{\alpha,\beta}(A^c) \sqsubseteq S_{\beta,\alpha}(A)^c$$

*Proof.* (1) From definition of this operators and complement of an intuitionistic fuzzy set we get that,

$$\beta(\nu_A(x) + (1 - \alpha)\mu_A(x)) \leq \beta(\nu_A(x) + (1 - \alpha)\mu_A(x) + \alpha)$$

and

$$\alpha(\mu_A(x) + (1 - \beta)\nu_A(x) + \beta) \geq \alpha(\mu_A(x) + (1 - \beta)\nu_A(x))$$

So, we can say  $T_{\beta,\alpha}(A)^c \sqsubseteq T_{\alpha,\beta}(A^c)$ .

(2) We can show this inclusion same way. □

**Theorem 2.2.** Let  $X$  be a set and  $A \in IFS(X)$ . If  $\alpha, \beta, \alpha + \beta \in [0, 1]$  and  $\beta \leq \alpha$  then

- (1)  $T_{\alpha,\beta}(A) \sqsubseteq T_{\beta,\alpha}(A)$
- (2)  $S_{\beta,\alpha}(A) \sqsubseteq S_{\alpha,\beta}(A)$

*Proof.* It is clear. □

**Theorem 2.3.** Let  $X$  be a set and  $A, B \in IFS(X)$ . If  $\alpha, \beta, \alpha + \beta \in [0, 1]$  then

- (1)  $T_{\alpha,\beta}(A) \sqcap T_{\alpha,\beta}(B) \sqsubseteq T_{\alpha,\beta}(A \sqcap B)$
- (2)  $T_{\alpha,\beta}(A \sqcup B) \sqsubseteq T_{\alpha,\beta}(A) \sqcup T_{\alpha,\beta}(B)$

*Proof.* (1) Let  $\alpha, \beta \in [0, 1]$ ,

$$\begin{aligned} \beta(1 - \alpha) \min(\nu_A(x), \nu_B(x)) &\leq \beta(1 - \alpha) \max(\nu_A(x), \nu_B(x)) \\ &\Rightarrow \beta(\min(\mu_A(x), \mu_B(x)) + (1 - \alpha) \min(\nu_A(x), \nu_B(x)) + \alpha) \\ &\leq \beta(\min(\mu_A(x), \mu_B(x)) + (1 - \alpha) \max(\nu_A(x), \nu_B(x)) + \alpha) \end{aligned}$$

and

$$\begin{aligned} \alpha(1 - \beta) \max(\mu_A(x), \mu_B(x)) &\geq \alpha(1 - \beta) \min(\mu_A(x), \mu_B(x)) \\ &\Rightarrow \alpha (\max(\nu_A(x), \nu_B(x)) + (1 - \beta) \max(\mu_A(x), \mu_B(x))) \\ &\geq \alpha (\max(\nu_A(x), \nu_B(x)) + (1 - \beta) \min(\mu_A(x), \mu_B(x))) \end{aligned}$$

It is appear from here that  $T_{\alpha,\beta}(A) \sqcap T_{\alpha,\beta}(B) \sqsubseteq T_{\alpha,\beta}(A \sqcap B)$ .

(2) It can be shown easily.  $\square$

**Theorem 2.4.** *Let  $X$  be a set and  $A, B \in IFS(X)$ . If  $\alpha, \beta, \alpha + \beta \in [0, 1]$  then*

- (1)  $S_{\alpha,\beta}(A \sqcup B) \sqsubseteq S_{\alpha,\beta}(A) \sqcup S_{\alpha,\beta}(B)$
- (2)  $S_{\alpha,\beta}(A) \sqcap S_{\alpha,\beta}(B) \sqsubseteq S_{\alpha,\beta}(A \sqcap B)$

*Proof.* (1) Let  $\alpha, \beta \in [0, 1]$ ,

$$\begin{aligned} \alpha(1 - \beta) \min(\nu_A(x), \nu_B(x)) &\leq \alpha(1 - \beta) \max(\nu_A(x), \nu_B(x)) \\ &\Rightarrow \alpha (\max(\mu_A(x), \mu_B(x)) + (1 - \beta) \min(\nu_A(x), \nu_B(x))) \\ &\leq \alpha (\max(\mu_A(x), \mu_B(x)) + (1 - \beta) \max(\nu_A(x), \nu_B(x))) \end{aligned}$$

and

$$\begin{aligned} \beta(1 - \alpha) \max(\mu_A(x), \mu_B(x)) &\geq \beta(1 - \alpha) \min(\mu_A(x), \mu_B(x)) \\ &\Rightarrow \beta (\min(\nu_A(x), \nu_B(x)) + (1 - \alpha) \max(\mu_A(x), \mu_B(x)) + \alpha) \\ &\geq \beta (\min(\nu_A(x), \nu_B(x)) + (1 - \alpha) \min(\mu_A(x), \mu_B(x)) + \alpha) \end{aligned}$$

Thus,  $S_{\alpha,\beta}(A \sqcup B) \sqsubseteq S_{\alpha,\beta}(A) \sqcup S_{\alpha,\beta}(B)$ .

(2) Can be proved similarly.  $\square$

#### REFERENCES

- [1] Atanassov K.T., Intuitionistic Fuzzy Sets, VII ITKR's Session, Sofia, June (1983).
- [2] Atanassov K.T., Intuitionistic Fuzzy Sets, Physica-Verlag, Heidelberg, NewYork, (1999).
- [3] Atanassov K.T., Studies in Fuzziness and Soft Computing-On Intuitionistic Fuzzy Sets Theory, ISBN 978-3-642-29126-5, Springer Heidelberg, New York, 2012.
- [4] Atanassov K.T., Çuvalcıođlu, G., Atanassova V. K., A new modal operator over intuitionistic fuzzy sets, Notes on IFS, 20(5), 2014, 1-8.
- [5] Çuvalcıođlu, G., On the diagram of One Type Modal Operators on Intuitionistic Fuzzy Sets: Last Expanding with  $Z_{\alpha,\beta}^{\omega,\theta}$ , Iranian Journal of Fuzzy Systems Vol. 10, No. 1, (2013) pp. 89-106.
- [6] Çuvalcıođlu G., New Tools Based on Intuitionistic Fuzzy Sets and Generalized Nets, ISBN 978-3-319-26301-4, Springer International Publishing Switzerland, 2016, 55-71.
- [7] Çuvalcıođlu G., Yılmaz S. On New Intuitionistic Fuzzy Operators:  $S_{\alpha,\beta}$  and  $T_{\alpha,\beta}$ , KAMERA, 43(2), 2015, 317-327.
- [8] Dencheva K., Extension of intuitionistic fuzzy modal operators  $\boxplus$  and  $\boxtimes$ , Proc. of the Second Int. IEEE Symp. Intelligent systems, Varna, June 22-24, (2004), Vol. 3, 21-22.
- [9] Yılmaz, S., Bal, A., Extensıon of Intuitionistic Fuzzy Modal Operators Diagram with New Operators, "Notes on IFS", Vol. 20, 2014, Number 5, pp. 26-35.
- [10] Zadeh L.A., Fuzzy Sets, Information and Control, 8, (1965) , p. 338-353.

UNIVERSITY OF MERSIN DEPARTMENT OF MATHEMATICS MERSIN, TURKEY.

*E-mail address:* [gcuvalcioglu@mersin.edu.tr](mailto:gcuvalcioglu@mersin.edu.tr)

DEPT. OF BIOINFORMATICS AND MATHEMATICAL MODELLING INSTITUTE OF BIOPHYSICS AND BIOMEDICAL ENGINEERING BULGARIAN ACADEMY OF SCIENCES

*E-mail address:* [krat@bas.bg](mailto:krat@bas.bg)

UNIVERSITY OF MERSIN DEPARTMENT OF MATHEMATICS MERSIN, TURKEY.

*E-mail address:* [sinemyilmaz@mersin.edu.tr](mailto:sinemyilmaz@mersin.edu.tr)

ON HERMITE-HADAMARD INEQUALITIES FOR  
GEOMETRIC-ARITHMETICALLY  $\varphi$ - $s$ -CONVEX FUNCTIONS  
VIA FRACTIONAL INTEGRALS

SÜMEYYE ERMEYDAN AND HÜSEYİN YILDIRIM

ABSTRACT. In this paper, we established integral inequalities of Hermite-Hadamard type involving Riemann-Liouville fractional integrals motivated by the definition of geometric-arithmetically  $s$ -convex and  $v$ -convex functions. We give equality for fractional integral of  $\varphi$ -convex functions on twice differentiable mapping.

Received: 26–August–2016

Accepted: 29–August–2016

1. INTRODUCTION

Fractional calculus was born in 1695. In the past three hundred years, fractional calculus developed in diverse fields from physical sciences and engineering to biological sciences and economics[1 – 8]. Fractional Hermite-Hadamard inequalities involving all kinds of fractional integrals have attracted by many researches. In [12], Shuang et al. introduced a new concept of geometric-arithmetically  $s$ -convex functions and presented interesting Hermite-Hadamard type inequalities for integer integrals of such functions. In [15], Youness introduced a new concept of  $\varphi$ -convex functions. In [16 – 17], YuMei Liao and colleagues gave Riemann-Liouville Hermite-Hadamard integral inequalities for once and twice differentiable geometric-arithmetically  $s$ -convex functions. We establish on Hermite-Hadamard inequalities for twice differentiable geometric-arithmetically  $\varphi - s$ -convex functions via fractional integrals.

2. PRELIMINARIES

In this section, we will give some definitions, lemmas and notations which we use later in this work.

**Definition 2.1.** (see [3]) Let  $f \in L[a, b]$ . The Riemann- Liouville fractional integral  $J_{a+}^{\alpha} f$  and  $J_{b-}^{\alpha} f$  of order  $\alpha > 0$  with  $\alpha \geq 0$  are defined by

$$J_{a+}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_a^x (x-t)^{\alpha-1} f(t) dt, 0 \leq a < x \leq b$$

and

$$(2.1) \quad J_{b-}^{\alpha} f(x) = \frac{1}{\Gamma(\alpha)} \int_x^b (t-x)^{\alpha-1} f(t) dt, 0 \leq a < x \leq b$$

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
2010 *Mathematics Subject Classification.* 26A33, 26D15, 41A55.

*Key words and phrases.* Fractional Hermite-Hadamard inequalities, geometric-arithmetically  $s$ -convex,  $\varphi$ -convex functions, Riemann-Liouville Fractional Integral.

Here  $\Gamma$  is the gamma function.

**Definition 2.2.** (see [12 – 16]) Let  $f : I \subseteq \mathbb{R}^+ \rightarrow \mathbb{R}^+$  and  $s \in (0, 1]$ . A function  $f(x)$  is said to be geometric-arithmetically  $s$ -convex on  $I$  if for every  $x, y \in I$  and  $t \in [0, 1]$ , we have:

$$(2.2) \quad f(x^t y^{1-t}) \leq t^s f(x) + (1-t)^s f(y)$$

**Definition 2.3.** (see [14]) The incomplete beta function is defined as follows:

$$(2.3) \quad B_x(a, b) = \int_0^x t^{a-1} (1-t)^{b-1} dt.$$

Where  $x \in [0, 1]$ ,  $a, b > 0$ .

**Definition 2.4.** (see [15]) Let  $\varphi : [a, b] \subset \mathbb{R} \rightarrow [a, b]$  A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be  $\varphi$ -convex on  $[a, b]$  if, for every  $x, y \in [a, b]$  and  $t \in [0, 1]$  the following inequality holds:

$$(2.4) \quad f(t\varphi(x) + (1-t)\varphi(y)) \leq t f(\varphi(x)) + (1-t) f(\varphi(y))$$

**Definition 2.5.** Let  $\varphi : [a, b] \subset \mathbb{R} \rightarrow [a, b]$  and  $s \in (0, 1]$ . A function  $f : [a, b] \rightarrow \mathbb{R}$  is said to be  $\varphi$ - $s$ -convex on  $[a, b]$  if, for every  $x, y \in [a, b]$  and  $t \in [0, 1]$  the following inequality holds:

$$(2.5) \quad f(\varphi(x)^t \varphi(y)^{(1-t)}) \leq t^s f(\varphi(x)) + (1-t)^s f(\varphi(y))$$

**Lemma 2.1.** (see [11]) For  $t \in [0, 1]$ , we have

$$(2.6) \quad \begin{aligned} (1-t)^n &\leq 2^{1-n} - t^n \quad \text{for } n \in [0, 1], \\ (1-t)^n &\geq 2^{1-n} - t^n \quad \text{for } n \in [0, \infty). \end{aligned}$$

The following inequality was used in the proof directly in [12].

**Lemma 2.2.** (see [13]) for  $t \in [0, 1]$  and  $x, y > 0$ , we have

$$(2.7) \quad tx + (1-t)y \geq y^{1-t} x^t.$$

**Lemma 2.3.** (see [10]) Let  $f : [a, b] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(a, b)$  with  $a < b$ . If  $f'' \in L[a, b]$ , then the following equality for fractional integrals holds:

$$(2.8) \quad \begin{aligned} &\frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \\ &= \frac{(b-a)^2}{2} \int_0^1 \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} f''(ta + (1-t)b) dt. \end{aligned}$$

**Lemma 2.4.** (see [9]) Let  $f : [a, b] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(a, b)$  with  $a < b$ . If  $f'' \in L[a, b]$ , then

$$(2.9) \quad \begin{aligned} &\frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] - f\left(\frac{a+b}{2}\right) \\ &= \frac{(b-a)^2}{2} \int_0^1 m(t) f''(ta + (1-t)b) dt. \end{aligned}$$

where

$$m(t) = \begin{cases} t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [0, \frac{1}{2}), \\ 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [\frac{1}{2}, 1]. \end{cases}$$

**Lemma 2.5.** (see [9]) Let  $f : [a, b] \rightarrow \mathbb{R}$  be a differentiable mapping on  $(a, b)$  with  $a < b$ . If  $f'' \in L[a, b]$ ,  $r > 0$ , then

$$(2.10) \quad \begin{aligned} &\frac{f(a)+f(b)}{r(r+1)} + \frac{2}{r+1} f\left(\frac{a+b}{2}\right) - \frac{\Gamma(\alpha+1)}{r(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \\ &= (b-a)^2 \int_0^1 k(t) f''(ta + (1-t)b) dt. \end{aligned}$$

Where

$$k(t) = \begin{cases} \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} - \frac{t}{r+1}, & t \in [0, \frac{1}{2}), \\ \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} - \frac{1-t}{r+1}, & t \in [\frac{1}{2}, 1]. \end{cases}$$

YuMei Liao and colleagues based on our study, they have provided the following theorems in [16],[17].

**Theorem 2.1.** *Let  $f : [0, b] \rightarrow R$  be a differentiable mapping. If  $|f'|$  is measurable and  $|f'|$  is decreasing and geometric-arithmetically  $s$ -convex on  $[0, b]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:*

$$(2.11) \quad \begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{(b-a)(2^{\alpha+s}|f'(b)| - |f'(a)| - |f'(b)|)}{2^{\alpha+s+1}(\alpha+s+1)} \\ & \quad + (b-a)|f'(a)| [0.5B(s+1, \alpha+1) - B_{0.5}(\alpha+1, s+1)] \\ & \quad + (b-a)|f'(b)| [B_{0.5}(s+1, \alpha+1) - 0.5B(a+1, \alpha+1)]. \end{aligned}$$

**Theorem 2.2.** *Let  $f : [0, b] \rightarrow R$  be a differentiable mapping. If  $|f''|$  is measurable and  $|f''|$  is decreasing and geometric-arithmetically  $s$ -convex on  $[0, b]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:*

$$(2.12) \quad \begin{aligned} & \left| \frac{f(a)+f(b)}{2} - \frac{\Gamma(\alpha+1)}{2(b-a)^\alpha} [J_{a^+}^\alpha f(b) + J_{b^-}^\alpha f(a)] \right| \\ & \leq \frac{(b-a)^2(|f''(a)|+|f''(b)|)}{2(\alpha+1)} \left( \frac{1}{s+1} - \frac{1}{\alpha+s+1} - B(s+1, \alpha+2) \right). \end{aligned}$$

### 3. MAIN RESULTS

**Lemma 3.1.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [\varphi(a), \varphi(b)] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(\varphi(a), \varphi(b))$ . If  $f'' \in L[\varphi(a), \varphi(b)]$ , then the following equality for fractional integral holds:*

$$(3.1) \quad \begin{aligned} & \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} [J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a))] \\ & = \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 \left( \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) f''(t\varphi(a) + (1-t)\varphi(b)) dt. \end{aligned}$$

*Proof.* By using Lemma 3 and Definition 5, we have

$$\begin{aligned} I &= \int_0^1 \left( \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) (f''(t\varphi(a) + (1-t)\varphi(b))) dt \\ &= \frac{1}{\varphi(b)-\varphi(a)} \left[ \int_0^1 ((1-t)^\alpha - t^\alpha) f'(t\varphi(a) + (1-t)\varphi(b)) dt \right] \\ &= \frac{f(\varphi(a))+f(\varphi(b))}{(\varphi(b)-\varphi(a))^2} - \frac{\alpha}{(\varphi(b)-\varphi(a))^2} \\ & \quad \times \left( \int_0^1 (1-t)^{\alpha-1} f(t\varphi(a) + (1-t)\varphi(b)) dt \right. \\ & \quad \left. + \int_0^1 t^{\alpha-1} f(t\varphi(a) + (1-t)\varphi(b)) dt \right) \\ &= \frac{f(\varphi(a))+f(\varphi(b))}{(\varphi(b)-\varphi(a))^2} - \frac{\alpha\Gamma(\alpha)}{(\varphi(b)-\varphi(a))^{\alpha+2}} \\ & \quad \times \left( \int_{\varphi(a)}^{\varphi(b)} (\varphi(x) - \varphi(a))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right. \\ & \quad \left. + \int_{\varphi(a)}^{\varphi(b)} (\varphi(b) - \varphi(x))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right) \\ &= \frac{f(\varphi(a))+f(\varphi(b))}{(\varphi(b)-\varphi(a))^2} - \frac{\Gamma(\alpha+1)}{(\varphi(b)-\varphi(a))^{\alpha+2}} [J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a))]. \end{aligned}$$



If  $I$  by multiplying  $\frac{(\varphi(b)-\varphi(a))^2}{2}$ , it obtain that

$$\begin{aligned} & \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 \left( \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) f'' t\varphi(a) + (1-t)\varphi(b) dt \\ & = \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right]. \end{aligned}$$

The proof is done. □

**Lemma 3.2.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [\varphi(a), \varphi(b)] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(\varphi(a), \varphi(b))$ . If  $f'' \in L[\varphi(a), \varphi(b)]$ , then the following equality for fractional integral holds:*

$$\begin{aligned} (3.2) \quad & \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \\ & = \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 m(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt, \end{aligned}$$

where

$$m(t) = \begin{cases} t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [0, \frac{1}{2}), \\ 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1}, & t \in [\frac{1}{2}, 1]. \end{cases}$$

*Proof.* By using Lemma 4 and Definition 5, we have:

$$\begin{aligned} I & = \int_0^1 m(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ & = \int_0^{\frac{1}{2}} \left( t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) (f''(t\varphi(a) + (1-t)\varphi(b))) dt \\ & \quad + \int_{\frac{1}{2}}^1 \left( 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) (f''(t\varphi(a) + (1-t)\varphi(b))) dt \\ & = I_1 + I_2. \end{aligned}$$

If use twice the partial integration method for  $I_1$ , we have

$$\begin{aligned} I_1 & = \int_0^{\frac{1}{2}} \left( t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ & = \frac{(\alpha-1+2^{1-\alpha})f'(\frac{\varphi(a)+\varphi(b)}{2})}{2(\alpha+1)(\varphi(a)-\varphi(b))} - \frac{f(\frac{\varphi(a)+\varphi(b)}{2})}{(\varphi(a)-\varphi(b))^2} + \frac{\Gamma(\alpha+1)}{(\varphi(a)-\varphi(b))^{\alpha+2}} \\ & \quad \times \left[ \int_{\frac{\varphi(a)+\varphi(b)}{2}}^{\varphi(b)} (\varphi(x) - \varphi(a))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right. \\ & \quad \left. + \int_{\frac{\varphi(a)+\varphi(b)}{2}}^{\varphi(a)} (\varphi(b) - \varphi(x))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right]. \end{aligned}$$

If use twice the partial integration method for  $I_2$ , we have

$$\begin{aligned} I_2 & = \int_{\frac{1}{2}}^1 \left( 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right) f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ & = - \left( \frac{(\alpha-1+2^{1-\alpha})f'(\frac{\varphi(a)+\varphi(b)}{2})}{2(\alpha+1)(\varphi(a)-\varphi(b))} \right) - \frac{f(\frac{\varphi(a)+\varphi(b)}{2})}{(\varphi(a)-\varphi(b))^2} \\ & \quad + \frac{\Gamma(\alpha+1)}{(\varphi(a)-\varphi(b))^{\alpha+2}} \left[ \int_{\varphi(a)}^{\frac{\varphi(a)+\varphi(b)}{2}} (\varphi(x) - \varphi(a))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right. \\ & \quad \left. + \int_{\varphi(a)}^{\frac{\varphi(a)+\varphi(b)}{2}} (\varphi(b) - \varphi(x))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right], \end{aligned}$$

by add uping  $I_1$  and  $I_2$ , and by multiplying  $\frac{(\varphi(b)-\varphi(a))^2}{2}$  with  $I$ , it obtain that:

$$\begin{aligned} & \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \\ & = \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 m(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt. \end{aligned}$$

The proof is done.

**Lemma 3.3.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [\varphi(a), \varphi(b)] \rightarrow \mathbb{R}$  be a twice differentiable mapping on  $(\varphi(a), \varphi(b))$ . If  $f'' \in L[\varphi(a), \varphi(b)]$ , then the following equality for fractional integral holds:*

$$(3.3) \quad \begin{aligned} & \frac{f(\varphi(a))+f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(\varphi(b)-\varphi(a))^\alpha} \\ & \quad \times \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \\ & = (\varphi(b) - \varphi(a))^2 \int_0^1 k(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt. \end{aligned}$$

Where

$$k(t) = \begin{cases} \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} - \frac{t}{r+1}, & t \in \left[0, \frac{1}{2}\right), \\ \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} - \frac{1-t}{r+1}, & t \in \left[\frac{1}{2}, 1\right). \end{cases}$$

□

*Proof.* By using Definition 5 and Lemma 5, we have

$$\begin{aligned} I &= \int_0^1 k(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ &= \frac{1}{r(r+1)(\alpha+1)} \\ & \quad \times \int_0^{\frac{1}{2}} \left[ (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - rt(\alpha+1) \right] \\ & \quad \times f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ & \quad + \frac{1}{r(r+1)(\alpha+1)} \\ & \quad \times \int_{\frac{1}{2}}^1 \left[ (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - r(1-t)(\alpha+1) \right] \\ & \quad \times f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ & = I_1 + I_2. \end{aligned}$$

If use twice the partial integration method for  $I_1$ , we have

$$\begin{aligned} I_1 &= \int_0^{\frac{1}{2}} \left[ (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - rt(\alpha+1) \right] \\ & \quad \times f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ &= \left[ (r+1)(1-2^{-\alpha}) - \frac{r(\alpha+1)}{2} \right] \frac{f'(\frac{\varphi(a)+\varphi(b)}{2})}{\varphi(a)-\varphi(b)} \\ & \quad + \frac{r(\alpha+1)f(\frac{\varphi(a)+\varphi(b)}{2})}{(\varphi(b)-\varphi(a))^2} + \frac{(\alpha+1)f(\varphi(b))}{(\varphi(b)-\varphi(a))^2} - \frac{(r+1)(\alpha+1)\alpha\Gamma(\alpha)}{(\varphi(b)-\varphi(a))^{\alpha+2}} \\ & \quad \times \left[ \int_{\frac{\varphi(a)+\varphi(b)}{2}}^{\varphi(b)} (\varphi(x) - \varphi(a))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right. \\ & \quad \left. + \int_{\frac{\varphi(a)+\varphi(b)}{2}}^{\varphi(b)} (\varphi(b) - \varphi(x))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right] \end{aligned}$$

If use twice the partial integration method for  $I_2$ , we have

$$\begin{aligned} I_2 &= \int_{\frac{1}{2}}^1 \left[ (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - r(1-t)(\alpha+1) \right] \\ & \quad \times f''(t\varphi(a) + (1-t)\varphi(b)) dt \\ &= \left[ \frac{r(\alpha+1)}{2} - (r+1)(1-2^{-\alpha}) \right] \frac{f'(\frac{\varphi(a)+\varphi(b)}{2})}{\varphi(a)-\varphi(b)} \\ & \quad + \frac{r(\alpha+1)f(\frac{\varphi(a)+\varphi(b)}{2})}{(\varphi(b)-\varphi(a))^2} + \frac{(\alpha+1)f(\varphi(a))}{(\varphi(b)-\varphi(a))^2} - \frac{(r+1)(\alpha+1)\alpha\Gamma(\alpha)}{(\varphi(b)-\varphi(a))^{\alpha+2}} \\ & \quad \times \left[ \int_{\varphi(a)}^{\frac{\varphi(a)+\varphi(b)}{2}} (\varphi(x) - \varphi(a))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right. \\ & \quad \left. + \int_{\varphi(a)}^{\frac{\varphi(a)+\varphi(b)}{2}} (\varphi(b) - \varphi(x))^{\alpha-1} f(\varphi(x)) d\varphi(x) \right]. \end{aligned}$$

By add uping  $I_1$  and  $I_2$ , and by multiplying with  $\frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)}$ , it obtain that:

$$\begin{aligned} & \frac{f(\varphi(a))+f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \\ & - \frac{\Gamma(\alpha+1)}{r(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] \\ & = (\varphi(b) - \varphi(a))^2 \int_0^1 k(t) f''(t\varphi(a) + (1-t)\varphi(b)) dt . \end{aligned}$$

The proof is done.

**Theorem 3.1.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|$  is measurable and  $|f''|$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:*

$$\begin{aligned} (3.4) \quad & \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] \\ & \leq \frac{(\varphi(b)-\varphi(a))^2 (|f''(\varphi(a))| + |f''(\varphi(b))|)}{2(\alpha+1)} \left( \frac{1}{s+1} - \frac{1}{\alpha+s+1} - B(s+1, \alpha+2) \right) . \end{aligned}$$

□

*Proof.* By using Definition 2, Lemma 2 and Lemma 6, we have

$$\begin{aligned} & \left| \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] \right| \\ & \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 \left| \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right| |f''(t\varphi(a) + (1-t)\varphi(b))| dt \\ & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \int_0^1 \left( 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right) |f''(\varphi^t(a) + \varphi^{(1-t)}(b))| dt \\ & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \int_0^1 \left( 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right) \\ & \quad \times [t^s |f''\varphi(a)| + (1-t)^s |f''\varphi(b)|] dt \\ & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''\varphi(a)| \\ & \quad \times \left[ -\int_0^1 t^{\alpha+s+1} dt + \int_0^1 t^s dt - \int_0^1 t^s (1-t)^{\alpha+1} dt \right] \\ & \quad + \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''\varphi(b)| \\ & \quad \times \left[ -\int_0^1 (1-t)^{\alpha+s+1} dt + \int_0^1 (1-t)^s dt - \int_0^1 t^{\alpha+1} (1-t)^s dt \right] \\ & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} (|f''\varphi(a)| + |f''\varphi(b)|) \\ & \quad \times \left[ -\frac{1}{\alpha+s+1} + \frac{1}{s+1} - B(s+1, \alpha+2) \right] . \end{aligned}$$

The poof is done.

**Theorem 3.2.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|^q$  is measurable and  $|f''|^q$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:*

$$\begin{aligned} (3.5) \quad & \left| \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] \right| \\ & \leq \frac{(\varphi(b)-\varphi(a))^2 \max\{1-2^{1-\alpha}, 2^{1-\alpha}-1\}}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right) . \end{aligned}$$

□

*Proof.* To achieve our aim ,we divide our proof into two cases.

Case 1:  $\alpha \in (0, 1)$ , by using Definition 2 , Hölder's inequality and Lemma 6 ,we have

$$\begin{aligned}
& \left| \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \right| \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 \left| \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right| |f''(t\varphi(a) + (1-t)\varphi(b))| dt \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \int_0^1 \left| 1-(1-t)^{\alpha+1}-t^{\alpha+1} \right|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 |f''(t\varphi(a) + (1-t)\varphi(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \int_0^1 \left| 1-(1-t)^{\alpha+1}-t^{\alpha+1} \right|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 |f''(\varphi^t(a)\varphi^{1-t}(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \int_0^1 \left| 1-(1-t)^{\alpha+1}-t^{\alpha+1} \right|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \left( \int_0^1 [(1-t)^\alpha + t^\alpha - 1]^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \left( \int_0^1 [2^{1-\alpha} - 1]^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2 (2^{1-\alpha} - 1)}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}}.
\end{aligned}$$

Where  $\frac{1}{p} + \frac{1}{q} = 1$ .

Case 2:  $\alpha \in [1, \infty)$ , by using Definition 2, Hölder's inequality and Lemma 6, we have

$$\begin{aligned}
& \left| \frac{f(\varphi(a))+f(\varphi(b))}{2} - \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \right| \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\
& \quad \times \left( \int_0^1 [1-(1-t)^\alpha - t^\alpha]^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2 (1-2^{1-\alpha})}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}}.
\end{aligned}$$

The proof is done.

**Theorem 3.3.** Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|$  is measurable and  $|f''|$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the

following inequality for fractional integrals holds:

$$\begin{aligned}
 & \left| \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right| \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2 |f''(\varphi(a))|}{2(\alpha+1)} \\
 (3.6) \quad & \times \left[ \frac{\alpha - \alpha 2^{-s-1} - 2^{-s-1}}{1+s} - \frac{\alpha+1}{2+s} + 2B(s+1, \alpha+2) + \frac{1}{\alpha+s+2} \right] \\
 & + \frac{(\varphi(b)-\varphi(a))^2 |f''(\varphi(b))|}{2(\alpha+1)} \\
 & \times \left[ \frac{\alpha 2^{-s-1} + 2^{-s-1} - 1}{1+s} + \frac{1}{\alpha+s+2} + 2B(s+1, \alpha+2) \right].
 \end{aligned}$$

□

*Proof.* By using Definition 5 and Lemma 7, we have

$$\begin{aligned}
 & \left| \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right| \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 |m(t)| |f''(t\varphi(a) + (1-t)\varphi(b))| dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 |m(t)| [f''(\varphi^t(a)\varphi^{1-t}(b))] dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 |m(t)| [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^{\frac{1}{2}} \left| t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right| [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \quad + \frac{(\varphi(b)-\varphi(a))^2}{2} \int_{\frac{1}{2}}^1 \left| 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right| [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''(\varphi(a))| \left[ \int_0^{\frac{1}{2}} [-t^s + (\alpha+1)t^{s+1} + t^s(1-t)^{\alpha+1} + t^{\alpha+s+1}] dt \right. \\
 & \quad \left. + \int_{\frac{1}{2}}^1 [\alpha t^s - (\alpha+1)t^{s+1} + t^s(1-t)^{\alpha+1} + t^{\alpha+s+1}] dt \right] + \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''(\varphi(b))| \\
 & \quad \times \left[ \int_0^{\frac{1}{2}} [-(1-t)^s + (\alpha+1)t(1-t)^{s+1} + (1-t)^{\alpha+s+1} + t^{\alpha+s}(1-t)^s] dt \right. \\
 & \quad \left. + \int_{\frac{1}{2}}^1 [\alpha(1-t)^s - (\alpha+1)t(1-t)^s + (1-t)^{\alpha+s+1} + t^{\alpha+1}(1-t)^s] dt \right] \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''(\varphi(a))| \left[ \frac{\alpha - \alpha 2^{-s-1} - 2^{-s-1}}{1+s} - \frac{\alpha+1}{s+2} + 2B(s+1, \alpha+2) + \frac{1}{\alpha+s+2} \right] \\
 & \quad + \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} |f''(\varphi(b))| \left[ \frac{\alpha 2^{-s-1} + 2^{-s-1} - 1}{1+s} + 2B(\alpha+2, s+1) + \frac{1}{\alpha+s+2} \right].
 \end{aligned}$$

**Theorem 3.4.** Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|$  is measurable and  $|f''|^q$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1)$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:

$$\begin{aligned}
 (3.7) \quad & \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \left( \frac{(\alpha+1)2^{-p-1} + (\alpha+0.5)^{p+1} - \alpha^{p+1}}{p+1} \right)^{\frac{1}{p}}.
 \end{aligned}$$

Where  $\frac{1}{p} + \frac{1}{q} = 1$ .

□

*Proof.* By using Lemma 2, Hölder's inequality and Lemma 7, we have

$$\begin{aligned}
& \left| \frac{\Gamma(\alpha+1)}{2(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] - f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) \right| \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \int_0^1 |m(t)| |f''(t\varphi(a) + (1-t)\varphi(b))| dt \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \left( \int_0^1 |m(t)|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 |f''(t\varphi(a) + (1-t)\varphi(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \left( \int_0^1 |m(t)|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 |f''(\varphi^t(a)\varphi^{(1-t)}(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2} \left( \int_0^1 |m(t)|^p dt \right)^{\frac{1}{p}} \\
& \quad \times \left( \int_0^1 [t^s |f''(\varphi(a))|^q + (1-t)^s |f''(\varphi(b))|^q] dt \right)^{\frac{1}{q}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\
& \quad \times \left( \int_0^{\frac{1}{2}} \left| t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right|^p dt + \int_{\frac{1}{2}}^1 \left| 1-t - \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{\alpha+1} \right|^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\
& \quad \times \left( (\alpha+1) \int_0^{\frac{1}{2}} t^p dt + \int_{\frac{1}{2}}^1 (\alpha-t+1)^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{2(\alpha+1)} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \left( \frac{(\alpha+1)2^{-p-1} + (\alpha+0.5)^{p+1} - \alpha^{p+1}}{p+1} \right)^{\frac{1}{p}}.
\end{aligned}$$

The proof is done.

**Theorem 3.5.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|$  is measurable and  $|f''|$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1]$ ,  $0 \leq a < b$ , then the*

following inequality for fractional integrals holds:

$$\begin{aligned}
 (3.8) \quad & \left| \frac{f(\varphi(a))+f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \right| \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \max \left\{ [r+1 - (r+1)2^{-\alpha}] \left( \frac{2^{-s-1}|f''(\varphi(a))| + (1-2^{-s-1})|f''(\varphi(b))|}{s+1} \right) \right. \\
 & \quad - r(\alpha+1) \left[ \frac{2^{-s-2}(s+1)|f''(\varphi(a))| + (1-t)^{s+1}(-ts-3t+1)f''(\varphi(a))}{(s+1)(s+2)} \right], \\
 & \quad \left. r(\alpha+1) \left[ \frac{2^{-s-2}(s+1)|f''(\varphi(a))| + (1-t)^{s+1}(-ts-3t+1)f''(\varphi(a))}{(s+1)(s+2)} \right] \right\} \\
 & \quad + \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \max \{ (r+1 - (r+1)2^{-\alpha} - r(\alpha+1)) \\
 & \quad \times \left[ \frac{(1-2^{-s-1})|f''(\varphi(a))| - 2^{-s-1}|f''(\varphi(b))|}{s+1} \right] \\
 & \quad + r(\alpha+1) \left[ \frac{(1-2^{-s-1})(s+1)|f''(\varphi(a))| + 2^{-s-2}(s+3)|f''(\varphi(b))|}{(s+1)(s+2)} \right], \\
 & \quad r(\alpha+1) \left[ \frac{(1-2^{-s-1})|f''(\varphi(a))| - 2^{-s-1}|f''(\varphi(b))|}{s+1} \right. \\
 & \quad \left. - \frac{(1-2^{-s-2})(s+1)|f''(\varphi(a))| + 2^{-s-2}(s+3)|f''(\varphi(b))|}{(s+1)(s+2)} \right] \}.
 \end{aligned}$$

□

*Proof.* By using Definition 3, Lemma 3 and Lemma 8, we have

$$\begin{aligned}
 & \left| \frac{f(\varphi(a))+f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \right| \\
 & \leq (\varphi(b) - \varphi(a))^2 \int_0^1 |k(t)| [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \int_0^{\frac{1}{2}} \left| (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - tr(\alpha+1) \right| \\
 & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \quad + \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \int_{\frac{1}{2}}^1 \left| (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - r(\alpha+1)(1-t) \right| \\
 & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\
 & \leq \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \max \left\{ [r+1 - (r+1)2^{-\alpha}] \left( \frac{2^{-s-1}|f''(\varphi(a))| + (1-2^{-s-1})|f''(\varphi(b))|}{s+1} \right) \right. \\
 & \quad - r(\alpha+1) \left[ \frac{2^{-s-2}(s+1)|f''(\varphi(a))| + (1-t)^{s+1}(-ts-3t+1)f''(\varphi(a))}{(s+1)(s+2)} \right], \\
 & \quad \left. r(\alpha+1) \left[ \frac{2^{-s-2}(s+1)|f''(\varphi(a))| + (1-t)^{s+1}(-ts-3t+1)f''(\varphi(a))}{(s+1)(s+2)} \right] \right\} \\
 & \quad + \frac{(\varphi(b)-\varphi(a))^2}{r(r+1)(\alpha+1)} \max (r+1 - (r+1)2^{-\alpha} - r(\alpha+1)) \left[ \frac{(1-2^{-s-1})|f''(\varphi(a))| - 2^{-s-1}|f''(\varphi(b))|}{s+1} \right] \\
 & \quad + r(\alpha+1) \left[ \frac{(1-2^{-s-1})(s+1)|f''(\varphi(a))| + 2^{-s-2}(s+3)|f''(\varphi(b))|}{(s+1)(s+2)} \right], \\
 & \quad r(\alpha+1) \left[ \frac{(1-2^{-s-1})|f''(\varphi(a))| - 2^{-s-1}|f''(\varphi(b))|}{s+1} \right. \\
 & \quad \left. - \frac{(1-2^{-s-2})(s+1)|f''(\varphi(a))| + 2^{-s-2}(s+3)|f''(\varphi(b))|}{(s+1)(s+2)} \right]
 \end{aligned}$$

Where we have used the following inequality:

$$\begin{aligned} & \int_0^{\frac{1}{2}} \left| (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - tr(\alpha+1) \right| \\ & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\ & \leq [(r+1)(1-2^{-\alpha})] \left[ \frac{2^{-s-1} |f''(\varphi(a))| + (1-2^{-s-1}) |f''(\varphi(b))|}{s+1} \right] \\ & \quad - r(\alpha+1) \left[ \frac{2^{-s-2} |f''(\varphi(a))|}{s+2} + \frac{1-(s+3)2^{-s-2} |f''(\varphi(b))|}{(s+2)(s+1)} \right] \end{aligned}$$

and

$$\begin{aligned} & \int_0^{\frac{1}{2}} \left| -r-1 + (r+1) \left[ (1-t)^{\alpha+1} + t^{\alpha+1} \right] + tr(\alpha+1) \right| \\ & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\ & \leq r(\alpha+1) \left[ \frac{2^{-s-2} |f''(\varphi(a))|}{s+2} + \frac{1-(s+3)2^{-s-2} |f''(\varphi(b))|}{(s+2)(s+1)} \right], \end{aligned}$$

and

$$\begin{aligned} & \int_{\frac{1}{2}}^1 \left| r+1 + tr(\alpha+1) - (r+1) \left[ (1-t)^{\alpha+1} + t^{\alpha+1} \right] - r(\alpha+1) \right| \\ & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\ & \leq [r+1 - (r+1)2^{-\alpha} - r(\alpha+1)] \left[ \frac{(1-2^{-s-1}) |f''(\varphi(a))| - 2^{-s-1} |f''(\varphi(b))|}{s+1} \right] \\ & \quad + r(\alpha+1) \left[ \frac{2^{-s-2} |f''(\varphi(a))|}{s+2} + \frac{(s+3)2^{-s-2} |f''(\varphi(b))|}{(s+2)(s+1)} \right], \end{aligned}$$

and

$$\begin{aligned} & \int_{\frac{1}{2}}^1 \left| -r-1 - tr(\alpha+1) + (r+1) \left[ (1-t)^{\alpha+1} + t^{\alpha+1} \right] + r(\alpha+1) \right| \\ & \quad \times [t^s |f''(\varphi(a))| + (1-t)^s |f''(\varphi(b))|] dt \\ & \leq r(\alpha+1) \\ & \quad \times \left[ \frac{(1-2^{-s-1}) |f''(\varphi(a))| - 2^{-s-1} |f''(\varphi(b))|}{s+1} - \frac{(1-2^{-s-2}) |f''(\varphi(a))|}{s+2} - \frac{(s+3)2^{-s-2} |f''(\varphi(b))|}{(s+2)(s+1)} \right]. \end{aligned}$$

The proof is done.

**Theorem 3.6.** *Let  $I$  be an interval  $a, b \in I$  with  $0 \leq a < b$  and  $\varphi : I \rightarrow \mathbb{R}$  a continuous increasing function. Let  $f : [0, \varphi(b)] \rightarrow \mathbb{R}$  be a differentiable mapping and  $1 < q < \infty$ . If  $|f''|$  is measurable and  $|f''|^q$  is decreasing and geometric-arithmetically  $\varphi - s$ -convex on  $[0, \varphi(b)]$  for some fixed  $\alpha \in (0, \infty)$ ,  $s \in (0, 1)$ ,  $0 \leq a < b$ , then the following inequality for fractional integrals holds:*

$$\begin{aligned} (3.9) \quad & \left| \frac{f(\varphi(a)) + f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a) + \varphi(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(\varphi(b) - \varphi(a))^\alpha} \left[ J_{\varphi(a)^+}^\alpha f(\varphi(b)) + J_{\varphi(b)^-}^\alpha f(\varphi(a)) \right] \right| \\ & \leq \frac{(\varphi(b) - \varphi(a))^2}{[r(r+1)(\alpha+1)]^{1+p-1}} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\ & \quad \times \left( \max \left\{ [(r+1)(1-2^{-\alpha})]^{p+1} - \left[ \frac{2+r(1-\alpha)-(1+r)2^{-\alpha}}{2} \right]^{p+1}, [r(\alpha+1)]^{p+1} 2^{-p-1} \right\} \right. \\ & \quad \left. + \max \left\{ [r+1 - (r+1)2^{-\alpha}]^{p+1} - \left[ \frac{2+r(1-\alpha)-(1+r)2^{-\alpha}}{2} \right]^{p+1}, \left[ \frac{r(\alpha+1)}{2} \right]^{p+1} \right\} \right)^{\frac{1}{p}}. \end{aligned}$$

Where  $\frac{1}{p} + \frac{1}{q} = 1$ .

□



*Proof.* By using Hölder's inequality and Lemma 7, we have

$$\begin{aligned}
& \left| \frac{f(\varphi(a))+f(\varphi(b))}{r(r+1)} + \frac{2}{r+1} f\left(\frac{\varphi(a)+\varphi(b)}{2}\right) - \frac{\Gamma(\alpha+1)}{r(\varphi(b)-\varphi(a))^\alpha} \left[ J_{\varphi(a)+}^\alpha f(\varphi(b)) + J_{\varphi(b)-}^\alpha f(\varphi(a)) \right] \right| \\
& \leq (\varphi(b) - \varphi(a))^2 \int_0^1 |k(t)| |f''(t\varphi(a) + (1-t)\varphi(b))| dt \\
& \leq (\varphi(b) - \varphi(a))^2 \left( \int_0^1 |k(t)|^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f''(t\varphi(a) + (1-t)\varphi(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq (\varphi(b) - \varphi(a))^2 \left( \int_0^1 |k(t)|^p dt \right)^{\frac{1}{p}} \left( \int_0^1 |f''(\varphi^t(a)\varphi^{(1-t)}(b))|^q dt \right)^{\frac{1}{q}} \\
& \leq (\varphi(b) - \varphi(a))^2 \left( \int_0^1 |k(t)|^p dt \right)^{\frac{1}{p}} \left( \int_0^1 [t^s |f''(\varphi(a))|^q + (1-t)^s |f''(\varphi(b))|^q] dt \right)^{\frac{1}{q}} \\
& \leq (\varphi(b) - \varphi(a))^2 \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \left( \int_0^1 |k(t)|^p dt \right)^{\frac{1}{p}} \\
& \leq (\varphi(b) - \varphi(a))^2 \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\
& \quad \times \left( \int_0^{\frac{1}{2}} \left| \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{r(\alpha+1)} - \frac{t}{r+1} \right|^p dt + \int_{\frac{1}{2}}^1 \left| \frac{1-(1-t)^{\alpha+1}-t^{\alpha+1}}{r(\alpha+1)} - \frac{1-t}{r+1} \right|^p dt \right)^{\frac{1}{p}} \\
& \leq \frac{(\varphi(b)-\varphi(a))^2}{[r(r+1)(\alpha+1)]^{1+p-1}} \left( \frac{|f''(\varphi(a))|^q + |f''(\varphi(b))|^q}{s+1} \right)^{\frac{1}{q}} \\
& \quad \left( \times \max \left\{ [(r+1)[1-2^{-\alpha}]]^{p+1} - \left[ \frac{2+r(1-\alpha)-(1+r)2^{-\alpha-1}}{2} \right]^{p+1}, [r(\alpha+1)]^{p+1} 2^{-p-1} \right\} \right. \\
& \quad \left. + \max \left\{ [(r+1)[1-2^{-\alpha}]]^{p+1} - \left[ \frac{2+r(1-\alpha)+(1+r)2^{-\alpha-1}}{2} \right]^{p+1}, \left[ \frac{r(\alpha+1)}{2} \right]^{p+1} \right\} \right)^{\frac{1}{p}}.
\end{aligned}$$

Where we have used the following inequalities:

$$\begin{aligned}
& \int_0^{\frac{1}{2}} \left| (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - tr(\alpha+1) \right|^p dt \\
& \leq \frac{[(r+1)[1-2^{-\alpha}]]^{p+1} - \left[ \frac{2+r(1-\alpha)-(1+r)2^{-\alpha-1}}{2} \right]^{p+1}}{r(\alpha+1)(p+1)},
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\frac{1}{2}}^1 \left| -r-1 + (r+1) \left[ (1-t)^{\alpha+1} + t^{\alpha+1} \right] + tr(\alpha+1) \right|^p dt \\
& \leq \frac{[r(\alpha+1)]^{p+1} 2^{-p-1}}{r(\alpha+1)(p+1)},
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\frac{1}{2}}^1 \left| (r+1) \left[ 1 - (1-t)^{\alpha+1} - t^{\alpha+1} \right] - r(1-t)(\alpha+1) \right|^p dt \\
& \leq \frac{[(r+1)[1-2^{-\alpha}]]^{p+1} - \left[ \frac{2+r(1-\alpha)+(1+r)2^{-\alpha-1}}{2} \right]^{p+1}}{r(\alpha+1)(p+1)},
\end{aligned}$$

and

$$\begin{aligned}
& \int_{\frac{1}{2}}^1 \left| -r-1 - tr(\alpha+1) + (r+1) \left[ (1-t)^{\alpha+1} + t^{\alpha+1} \right] + r(\alpha+1) \right|^p dt \\
& \leq \frac{\left[ \frac{r(\alpha+1)}{2} \right]^{p+1}}{r(\alpha+1)(p+1)}.
\end{aligned}$$

The proof is done.

□

## REFERENCES

- [1] Baleanu, D, Machado, JAT, Luo, ACJ: Fractional Dynamics and Control. Springer, New York (2012)
- [2] Diethelm, K: The Analysis of Fractional Differential Equations. Lecture Notes in Mathematics.(2010)
- [3] Kilbas, AA, Srivastava, HM , Trujillo, JJ: Theory and Applications of Fractional Differential Equations.Elsevier, Amsterdam (2006)
- [4] Lakshmikantham, V, Leela, S, Devi, JV: Theory of Fractional Dynamic Systems. Cambridge Scientific Publishers, Cambridge (2009)
- [5] Miller, KS, Ross, B: An Introduction to the Fractional Calculus and Differential Equations. Wiley, New York (1993)
- [6] Michalski, MW: Derivates of noninteger Order and Their Applications. Diss. Math. CC-CXXVIII. Inst. Math., Polish Acad.Sci.(1993)
- [7] Podlubny, I: Fractional Differential Equations.Academic Press, San Diego (1999)
- [8] Tarasov, VE. Fractional Dynamics: Application of Fractional Calculus to Dynamics of Particles, Fields and Media.Springer, Berlin (2011)
- [9] Zhang, Y, Wang, J: On some new Hermite-Hadamard inequalities involving Riemann-Liouville fractional integrals. J.Inequal. Appl. 2013, 220 (2013)
- [10] Wang, J, Li,X, Feckan, M, Zhou, Y: Hermite-Hadamard-type inequalities for Riemann-Liouville fractional integrals via two kinds of convexity. Appl. Anal. (2012).doi: 10.1080/00036811.2012.727986
- [11] Deng, J, Wang, J: Fractional Hermite-Hadamard inequalities for  $(\alpha, m)$ -logarithmically convex functions.J. Inqual.Appl. 2013, 364 (2013). doi: 10.1186/1029-242X-2013-364
- [12] Shuang, Y, Yin, HP, QI, F: Hermite-Hadamard type integral inequalities for geometric-arithmetically  $s$ -convex functions. Analysis 33, 197-208 (2013)
- [13] Liao, Y, Deng, J, Wang, j: Riemann-Liouville fractional Hermite-Hadamard inequalities. Part I: for once differentiable Geometric-Arithmetically  $s$ -convex functions. J.Inequal. Appl. 2013, 443 (2013)
- [14] DiDonato, AR, Jarnagin, MP: The efficient calculation of the incomplete beta-function ratio for half-integer values of the parameters. Math. Comput. 21, 652-662 (1967)
- [15] E. A. Youness, E-convex sets, E-convex functions and E-convex programming, J. of Optim.Theory and Appl., 102, No.2(1999), 439-450
- [16] Liao, Y, Deng, J, Wang, j: Riemann-Liouville fractional Hermite-Hadamard inequalities. Part I: for once differentiable Geometric-Arithmetically  $s$ -convex functions. J.Inequal. Appl. 2013, 443 (2013)
- [17] Liao, Y, Deng, J, Wang, j: Riemann-Liouville fractional Hermite-Hadamard inequalities. Part II: for twice differentiable Geometric-Arithmetically  $s$ -convex functions. J.Inequal. Appl. 2013, 517 (2013)

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, UNIVERSITY OF KAHRAMANMARAŞ  
 SÜTÇÜ İMAM, 46100, KAHRAMANMARAŞ, TURKEY  
*E-mail address:* [sumeyye\\_ermeydan@hotmail.com](mailto:sumeyye_ermeydan@hotmail.com)  
*E-mail address:* [hyildir@ksu.edu.tr](mailto:hyildir@ksu.edu.tr)

**ON A BOUNDARY VALUE PROBLEM WITH RETARDED  
ARGUMENT IN THE DIFFERENTIAL EQUATION**

F. AYCA CETINKAYA, KHANLAR R. MAMEDOV, AND GIZEM CERCI

ABSTRACT. In this work a boundary value problem on the half axis is studied. The special solution of this boundary value problem is defined. The simplicity of the eigenvalues is shown and it is proven that all positive values of the parameter  $\lambda$  are the eigenvalues of this boundary value problem.

Received: 19–August–2016

Accepted: 29–August–2016

1. INTRODUCTION

In this work we study the boundary value problem

$$(1) \quad y''(x) + \lambda y(x) + M(x)y(x - \Delta(x)) = 0 \quad (0 \leq x < \infty),$$

$$(2) \quad y'(0) - hy(0) = 0,$$

$$(3) \quad y(x - \Delta(x)) \equiv \phi(x - \Delta(x)), \text{ if } x - \Delta(x) < 0,$$

$$(4) \quad \sup_{[0, \infty)} |y(x)| < \infty,$$

where  $M(x)$  and  $\Delta(x) \geq 0$  are defined and continuous on the half axis  $[0, \infty)$ ,  $\lambda$  is a real parameter ( $-\infty < \lambda < \infty$ ),  $h$  is an arbitrary real number and  $\phi(x)$  is a continuous initial function on the initial set

$$E_0 = \{x - \Delta(x) : x - \Delta(x) < 0, x > 0\} \cup \{0\}$$

with  $\phi(0) = 1$ .

The literature for the boundary value problems for differential equations of the second order with retarded arguments begins with [1, 2, 3, 4, 5, 6, 7, 8, 9].

Differential equations with retarded argument, describe processes with aftereffect; they find many applications, particularly in the theory of automatic control, in the theory of self-oscillatory systems, in the study of problems connected with combustion in rocket engines, in a number of problems in economics, biophysics, and many other fields. Equations with retarded argument appear, for example each time when in some physical or technological problem, the force operating at the mass point depends on the velocity and the position of this point, not only at the given instant, but also at some given previous instant.

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference

The presence of retardations in the system studied often proves to be a result of a phenomenon which essentially influences the course of the process. For example, in automatic control systems the retardation is the time interval which the system requires to react to an input impulse. Various physical applications of such problems can be found in [8].

The rest of this paper is organized as follows. First, the equivalent integral representation for the solution of the boundary value problem (1)-(4) is constructed. Then, the simplicity of the eigenvalues is shown. It is proven that all positive values of the parameter  $\lambda$  are the eigenvalues of the boundary value problem (1)-(4).

## 2. THE SPECIAL SOLUTION

Let  $w(x, \lambda)$  be a solution of (1) which satisfies the conditions

$$(5) \quad w(0, \lambda) = 1, \quad w'(0, \lambda) = h,$$

$$(6) \quad w(x - \Delta(x), \lambda) \equiv \phi(x - \Delta(x)), \quad \text{if } x - \Delta(x) < 0.$$

From Theorem I.2.1 (see [8]) it follows that under conditions (5), (6) there exists a unique solution of (1) on the half axis  $[0, \infty)$ .

**Lemma 2.1.** *Equation (1) together with the initial conditions (5), (6) are equivalent, for each value of  $\lambda > 0$  to the integral equation:*

$$(7) \quad w(x, \lambda) = \cos sx + \frac{h \sin sx}{s} - \frac{1}{s} \int_0^x M(t) \sin s(x-t) w(t - \Delta(t), \lambda) dt, \quad (s^2 = \lambda).$$

*Proof.* If we seek the solution of equation

$$w''(x, \lambda) + \lambda w(x, \lambda) = -q(x)w(x - \Delta(x), \lambda)$$

as

$$w(x, \lambda) = c_1 \cos sx + c_2 \sin sx,$$

by applying the method of variation of parameters we have

$$w(x, \lambda) = \tilde{c}_1 \cos sx + \tilde{c}_2 \sin sx - \frac{1}{s} \int_0^x q(t) \sin s(x-t) w(t - \Delta(t), \lambda) dt.$$

Taking condition (5) into consideration we find

$$\tilde{c}_1 = 1 \quad \text{and} \quad \tilde{c}_2 = \frac{h}{s}.$$

□

Before giving the following theorem, it will be useful to keep in mind that; here, the multiplicity of an eigenvalue of a boundary value problem is defined to be the number of linearly independent eigenfunctions corresponding to this eigenvalue.

**Theorem 2.1.** *The boundary value problem (1)-(4) can have only simple eigenvalues.*

*Proof.* Let  $\tilde{\lambda}$  be an eigenvalue of the boundary value problem (1)-(4) and  $\tilde{\varphi}(x, \tilde{\lambda})$  a corresponding eigenfunction. By (2) and (5),

$$W \left\{ \tilde{\varphi} \left( 0, \tilde{\lambda} \right), w \left( 0, \tilde{\lambda} \right) \right\} = \begin{vmatrix} \tilde{\varphi} \left( 0, \tilde{\lambda} \right) & 1 \\ \tilde{\varphi}' \left( 0, \tilde{\lambda} \right) & h \end{vmatrix} = 0,$$

and according to Theorem II.2.2. in [8] the functions  $\tilde{\varphi}(x, \tilde{\lambda})$  and  $w(x, \tilde{\lambda})$  are linearly dependent on  $[0, \infty)$ . Hence it follows that  $w(x, \tilde{\lambda})$  is an eigenfunction for the boundary value problem (1)-(4) and all eigenfunctions of this boundary value problem which correspond to the eigenvalue  $\tilde{\lambda}$  are pairwise linearly dependent.  $\square$

### 3. EXISTENCE THEOREM

**Theorem 3.1.** *Let*

$$(8) \quad \sup_{t \in E_0} |\phi(x)| = \phi_0 < \infty$$

and in equation (1) let

$$(9) \quad \int_0^\infty |M(t)| dt = M_\infty < \infty.$$

Then all positive values of the parameter  $\lambda$  are eigenvalues of the boundary value problem (1)-(4).

*Proof.* By (7), if  $\lambda > 0$ , then

$$(10) \quad w(x, \lambda) = R_\lambda \sin(st - \psi_\lambda) - \frac{1}{s} \int_0^x M(t) \sin s(x-t) w(t - \Delta(t), \lambda) dt,$$

where

$$R_\lambda = \sqrt{1 + \frac{h^2}{\lambda}}, \quad \cos \psi_\lambda = \frac{1}{R_\lambda}, \quad \sin \psi_\lambda = \frac{h}{sR_\lambda}, \quad (0 \leq \psi_\lambda < 2\pi).$$

Let  $x_0 \in (0, \infty)$ , and  $N_\lambda(x_0) = \max_{[0, x_0]} |w(x, \lambda)|$ . Evidently,  $N_\lambda(x_0) \geq N_\lambda(x)$  ( $x_0 \geq x$ )

and, from (10), (5), (6), (8) and (9) one of the following inequalities holds:

$$(11) \quad N_\lambda(x_0) \leq R_\lambda + \frac{1}{s} \int_0^{x_0} |M(t)| N_\lambda(t) dt$$

or

$$N_\lambda(x_0) \leq R_\lambda + \frac{\phi_0}{s} \int_0^{x_0} |M(t)| dt \leq R_\lambda + \frac{\phi_0 M_\infty}{s}.$$

By Lemma II.3.5 in [8] it follows from (11) that

$$N_\lambda(x_0) \leq R_\lambda \exp \frac{1}{s} \int_0^{x_0} |M(t)| dt \leq R_\lambda \exp \frac{M_\infty}{s},$$

and for  $s > 0$

$$N_\lambda(x_0) \leq \max \left\{ R_\lambda \exp \frac{M_\infty}{s}; R_\lambda + \frac{\phi_0 M_\infty}{s} \right\} < \infty.$$

The bound obtained is valid for any  $\lambda > 0$  and is independent of  $x_0$ , proving the theorem.  $\square$

### REFERENCES

- [1] Bellman, R. and Cook, K. L., *Differential-Difference Equations*, New York Academic Press: London, 1963.
- [2] Demidenko, D. M. and Likhoshvai V. A., *On differential equations with retarded argument*, Siberian Mathematical Journal. **46** (3) (2005), 417-430.
- [3] Elsgolc, L. E., *Qualitative Methods of Mathematical Analysis*, GITTL, Moscow, 1955; English transl., Transl. Math. Monographs, vol. 12, Amer. Math. soc., Providence, R. I., 1964.
- [4] Elsgolc, L. E., *Introduction to the Theory of Differential Equations with Deviating Arguments*, Nauka, Moscow, 1964; English transl., Holden-Day, San Fransisco, Calif., 1966.

- [5] Krasovskii, N. N., *Certain Problems in the Theory of the Stability of Motion*, Fizmatgiz, Moscow, 1959; English transl., *Stability of Motion. Application of the Ljapunov's Second Method to Differential Systems and Equations with Delay*, Stanford Univ. Press, Stanford, Calif., 1963.
- [6] Myshkis, A. D., *Linear Differential Equations with Retarded Argument*, GITTL, Moscow, 1951; German transl., VEB Deutscher Verlag, Berlin, 1955.
- [7] Norkin, S. B., *On a boundary problem of Sturm-Liouville type for second-order differential equation with retarded argument*, *Izvestiya Vysshikh Uchebnykh Zavedenii Matematika*. **6 (7)** (1958), 203-214 (Russian).
- [8] Norkin, S. B., *Differential Equations of the Second Order with Retarded Argument*, *Translations of Mathematical Monographs*, Vol. 31, AMS: Providence, 1972.
- [9] Pinney, E., *Ordinary Difference-Differential Equations*, Univ. of California Press, Berkeley, 1958; Russian transl., IL, Moscow, 1961.

MERSIN UNIVERSITY, DEPARTMENT OF MATHEMATICS, MERSIN, TURKEY

*E-mail address:* aycacetinkaya@mersin.edu.tr

*E-mail address:* hanlar@mersin.edu.tr

*E-mail address:* cercigizem@gmail.com

## Ω– ALGEBRAS ON CO-HEYTING VALUED SETS

MEHMET CITIL AND SINEM TARSUSLU(YILMAZ)

ABSTRACT. As known, co-Heyting algebras are dual to Heyting algebras. co-Heyting algebra has many studying areas as topos theory, co-intuitionistic logic, linguistics, quantum theory, etc.

In this paper, we studied on co-Heyting valued sets. The co-Heyting valued Ω–algebra, co-Heyting Valued Algebra Homomorphism are defined and some properties of these sets are examined.

Received: 05–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

co- Heyting algebra that is a lattice which dual is Heyting algebra. co-Heyting algebra has applications in many different areas.

**Definition 1.1.** [1]A Boolean algebra is an algebra  $(H, \vee, \wedge, -, 0_H, 1_H)$  where  $(H, \vee, \wedge, 0_H, 1_H)$  is a distributive lattice and for all  $a \in H$ ,

$$a \wedge \bar{a} = 0 \text{ and } a \vee \bar{a} = 1$$

**Definition 1.2.** [1]A Heyting algebra is an algebra  $(H, \vee, \wedge, \rightarrow, 0_H, 1_H)$  such that  $(H, \vee, \wedge, 0_H, 1_H)$  is an lattice and for all  $a, b, c \in H$ ,

$$a \leq b \rightarrow c \Leftrightarrow a \wedge b \leq c$$

$(H, \vee, \wedge, 0_H, 1_H)$  is a Heyting algebra with  $\forall a, b \in H$ ,

$$a \rightarrow b = \bigvee \{c : a \wedge c \leq b, c \in H\}.$$

**Proposition 1.1.** [4]An algebra  $(H, \vee, \wedge, \rightarrow, 0_H, 1_H)$  is a Heyting algebra if and only if  $(H, \vee, \wedge, 0_H, 1_H)$  is an lattice and the following identities hold for all  $a, b, c \in H$ ,

- (1)  $a \rightarrow a = 1$
- (2)  $a \wedge (a \rightarrow b) = a \wedge b$
- (3)  $b \wedge (a \rightarrow b) = b$
- (4)  $a \rightarrow (b \wedge c) = (a \rightarrow b) \wedge (a \rightarrow c)$

**Definition 1.3.** [1]A co-Heyting algebra is an algebra  $(H^*, \vee, \wedge, \leftrightarrow, 0_{H^*}, 1_{H^*})$  such that  $(H^*, \vee, \wedge, 0_{H^*}, 1_{H^*})$  is an lattice and for all  $a, b, c \in H^*$ ,

$$a \leftrightarrow b \leq c \Leftrightarrow a \leq b \vee c$$

---

<sup>13<sup>rd</sup></sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
 2010 *Mathematics Subject Classification.* Primary 03C05.

*Key words and phrases.* Omega Algebra, co-Heyting Valued Algebra, co-Heyting Valued Algebra Homomorphism.

$(H^*, \vee, \wedge, 0_{H^*}, 1_{H^*})$  is a co-Heyting algebra with  $\forall a, b \in H^*$ ,

$$a \hookrightarrow b = \bigwedge \{c : a \vee c \geq b, c \in H^*\}.$$

A co-Heyting algebra with the ordering reversed will yield a Heyting algebra. The implication operation in this algebra will be  $a \rightarrow b = b \hookrightarrow a$ .

It is clear that  $H$  and  $H^*$  are same sets with different order relations.  $1_{H^*}$  and  $0_{H^*}$  are greatest and least elements of  $H^*$ , respectively.

**Proposition 1.2.** *An algebra  $(H^*, \vee, \wedge, \hookrightarrow, 0_{H^*}, 1_{H^*})$  is a co-Heyting algebra if and only if  $(H^*, \vee, \wedge, 0_{H^*}, 1_{H^*})$  is a lattice and the following identities hold for all  $a, b, c \in H^*$ ,*

- (1)  $a \hookrightarrow a = 0$
- (2)  $a \vee (b \hookrightarrow a) = a \vee b$
- (3)  $b \vee (b \hookrightarrow a) = b$
- (4)  $(b \vee c) \hookrightarrow a = (b \hookrightarrow a) \vee (c \hookrightarrow a)$

*Proof.* (1)  $\forall a \in H^*$ ,

$$a \hookrightarrow a = \bigwedge \{c : a \vee c \geq a, c \in H^*\} = 0$$

(2) From definition it is obtained that,

$$a \vee (b \hookrightarrow a) \geq b \Rightarrow a \vee (b \hookrightarrow a) \geq a \vee b$$

and

$$\begin{aligned} (a \vee b) \vee a &\geq b \Rightarrow a \vee b \geq b \hookrightarrow a \text{ and } a \leq a \vee b \\ &\Rightarrow a \vee b \geq a \vee (b \hookrightarrow a) \end{aligned}$$

(3)  $\forall a, b \in H^*$ ,

$$b \leq a \vee b \Rightarrow b \vee (b \hookrightarrow a) = b$$

(4)  $\forall a, b, c \in H^*$ ,

$$\begin{aligned} (b \hookrightarrow a) \vee (c \hookrightarrow a) \vee a &= (a \vee (b \hookrightarrow a)) \vee (a \vee (c \hookrightarrow a)) \geq b \vee c \\ &\Rightarrow (b \hookrightarrow a) \vee (c \hookrightarrow a) \geq (b \vee c) \hookrightarrow a \end{aligned}$$

On the other hand,  $(b \vee c) \hookrightarrow a \geq (b \hookrightarrow a) \vee (c \hookrightarrow a)$ . □

## 2. CO-HEYTING VALUED SETS

In this section, the concepts of co-Heyting valued set, co-Heyting valued function are defined and some properties of these structures are examined.

**Definition 2.1.** Let  $H^*$  be a complete co-Heyting algebra and  $X$  be a universal.  $H^*$ -valued set is determined with  $[=]$  function

$$[=] : X \times X \rightarrow H^*, [=](a, b) = [a = b]$$

which satisfy the following conditions.

- (1)  $[a = b] \geq [b = a]$
- (2)  $[a = b] \vee [b = c] \geq [a = c]$

Let  $X$  be a universal.  $u \in X$ ,  $E(u)$  means the degree of existence the element  $u$ . For  $H^*$ -valued sets we will use,

$$E(u) = [u \in X].$$

So,  $[u \in X] = [u = u]$ .



**Definition 2.2.** Let  $A$  be a  $H^*$ -valued set. The subset of  $A$  is a  $s : A \rightarrow H^*$  function with following conditions.

- (1)  $[x \in s] \vee [x = y] \geq [y \in s]$
- (2)  $[x \in s] \geq [x \in A]$

**Definition 2.3.** Let  $(X, =)$  and  $(Y, =)$  are  $H^*$ -valued sets. If  $f : X \times Y \rightarrow H^*$  function satisfy the following conditions then called  $H^*$ -valued function and it is shown  $f : X \rightarrow Y$ .

- F1  $f(x, y) \geq [x = x] \vee [y = y]$
- F2  $[x = x'] \vee f(x, y) \vee [y = y'] \geq f(x', y')$
- F3  $f(x, y) \vee f(x, y') \geq [y = y']$
- F4  $[x = x] \geq \bigwedge \{f(x, y) : y \in Y\}$

**Notation 1:**  $f(x, y) := [f(x) = y]$

**Definition 2.4.** Let  $(X, =)$  be an  $H^*$ -valued set.  $I : X \times X \rightarrow H^*$ ,  $I(x, x') = [x = x']$  function is called unit function.

**Definition 2.5.** Let  $(X, =)$ ,  $(Y, =)$  and  $(Z, =)$  are  $H^*$ -valued sets and  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$  are  $H^*$ -valued functions. For  $x \in X, z \in Z$ ,

$$(g \circ f)(x, z) = \bigwedge \{f(x, y) \vee g(y, z) : y \in Y\}.$$

**Proposition 2.1.** Let  $(X, =)$ ,  $(Y, =)$  and  $(Z, =)$  are  $H^*$ -valued sets and  $f : X \rightarrow Y$ ,  $g : Y \rightarrow Z$  are  $H^*$ -valued functions. The function  $(g \circ f) : X \rightarrow Z$  is a  $H^*$ -valued function.

*Proof.* (i) Let  $x \in X, z \in Z$ ,

$$\begin{aligned} (g \circ f)(x, z) &= \bigwedge \{f(x, y) \vee g(y, z) : y \in Y\} \\ &\geq \bigwedge \{[x = x] \vee [y = y] \vee [z = z] : y \in Y\} \\ &= [x = x] \vee [z = z] \vee \bigwedge \{[y = y] : y \in Y\} \\ &\geq [x = x] \vee [z = z] \end{aligned}$$

(ii) Let  $x, x' \in X$  and  $z, z' \in Z$ ,

$$\begin{aligned} [x = x'] \vee (g \circ f)(x, z) \vee [z = z'] &= [x = x'] \vee \bigwedge \{f(x, y) \vee g(y, z) : y \in Y\} \vee [z = z'] \\ &= \bigwedge \{[x = x'] \vee f(x, y) \vee g(y, z) \vee [z = z'] : y \in Y\} \\ &= \bigwedge \left\{ \begin{array}{l} [x = x'] \vee f(x, y) \vee [y = y] \\ \vee g(y, z) \vee [z = z'] : y \in Y \end{array} \right\} \\ &\geq \bigwedge \{f(x', y') \vee g(y', z') : y \in Y\} = f(x', y') \vee g(y', z') \\ &\geq \bigwedge \{f(x', y') \vee g(y', z') : y' \in Y\} = (g \circ f)(x', z') \end{aligned}$$

(iii) Let  $x \in X$  and  $z, z' \in Z$ ,

$$\begin{aligned} (g \circ f)(x, z) \vee (g \circ f)(x, z') &= \bigwedge \{f(x, y) \vee g(y, z) : y \in Y\} \vee \bigwedge \{f(x, t) \vee g(t, z') : t \in Y\} \\ &= \bigwedge \{f(x, y) \vee f(x, y) \vee g(y, z) \vee g(y, z') : y \in Y\} \\ &\geq \bigwedge \{[y = y] \vee [z = z'] : y \in Y\} \\ &\geq [z = z'] \end{aligned}$$

(iv) Let  $x \in X$ ,

$$\begin{aligned} \bigwedge \{(g \circ f)(x, z) : z \in Z\} &= \bigwedge \left\{ \bigwedge \{f(x, y) \vee g(y, z) : y \in Y\} : z \in Z \right\} \\ &= \bigwedge \left\{ \bigwedge \{f(x, y) : y \in Y\} : z \in Z \right\} \vee \\ &\quad \bigwedge \left\{ \bigwedge \{g(y, z) : y \in Y\} : z \in Z \right\} \\ &\leq [x = x] \vee \bigwedge \{[y = y] : y \in Y\} = [x = x] \end{aligned}$$

□

**Definition 2.6.** Let  $(X, =)$  and  $(Y, =)$  are  $H^*$ -valued sets and  $f : X \rightarrow Y$  is  $H^*$ -valued function.

(1)  $f$  is a monomorphism.  $\Leftrightarrow \forall x, x' \in X, y \in Y$ ,

$$f(x, y) \vee f(x', y) \geq [x = x']$$

(2)  $f$  is an epimorphism.  $\Leftrightarrow \forall y \in Y$ ,

$$[y = y'] \geq \bigwedge \{f(x, y) : x \in X\}$$

**Definition 2.7.** Let  $(X, =)$  be a  $H^*$ -valued set.  $R : X \times X \rightarrow H^*$  is called  $H^*$ -valued equivalence relation.  $\Leftrightarrow$

- R1  $R(x, y) \vee [x = x] = R(x, y), R(x, y) \vee [y = y] = R(x, y)$
- R2  $R(x, y) \vee [x = x'] \geq R(x', y), R(x, y) \vee [y = y'] \geq R(x, y')$
- R3  $[x = x] \geq R(x, x) :$
- R4  $R(x, y) \geq R(y, x)$
- R5  $R(x, y) \vee R(y, z) \geq R(x, z)$

**Example 2.1.** Let  $(X, =), (Y, =)$  are  $H^*$ -valued sets and  $f : X \rightarrow Y$  is  $H^*$ -valued function.  $\forall x_1, x_2 \in X$ ,

$$C_f(x_1, x_2) = [f(x_1) = f(x_2)]$$

function is a  $H^*$ -valued equivalence relation on  $X$ .

**Definition 2.8.** Let  $R$  be a  $H^*$ -valued equivalence relation on  $X$ .  $d : X \rightarrow H^*$  is called equivalence class of  $R \Leftrightarrow$

- d1  $d(x) \vee R(x, x') \geq d(x')$
- d2  $d(x) \vee d(y) \geq R(x, y)$

$d(x)$  is the equivalence class of  $x \in X$ .

**Proposition 2.2.** Let  $R$  be a  $H^*$ -valued equivalence relation on  $X$  and  $d_1, d_2$  are equivalence class of  $R$ .

$$\begin{aligned} \bigwedge \{d_1(x) : x \in X\} &= \bigwedge \{d_2(x) : x \in X\} \text{ and} \\ d_1(x) &\geq d_2(x) \Rightarrow d_1 = d_2 \end{aligned}$$

*Proof.*  $x_0 \in X$ ,

$$\begin{aligned} d_2(x_0) &= \bigwedge \{d_2(x) \vee d_1(x) : x \in X\} \\ &\geq \bigwedge \{R(x_0, x) \vee d_1(x) : x \in X\} \\ &\geq d_1(x_0) \end{aligned}$$

□

**Proposition 2.3.** Let  $(X, =), (Y, =)$  are  $H^*$ -valued sets and  $f : X \rightarrow Y$  is  $H^*$ -valued function.  $f$  is surjective  $\Leftrightarrow \forall y \in Y,$

$$\bigwedge \{[f(x) = y] : x \in X\} = [y = y]$$

### 3. $H^*$ -VALUED Ω-ALGEBRAS

Now, let  $\Omega$  is defined as follows,

$$\Omega = \{\omega : X^n \times X \rightarrow H^* : \omega \text{ satisfy F1-F4 conditions}\}$$

It means that, if  $\omega \in \Omega,$   $\omega$  is  $H^*$ -valued function. The concept of  $H^*$ -valued  $\Omega$ -algebra can be defined as following;

**Definition 3.1.**  $A = \langle X, \Omega \rangle$  is  $H^*$ -valued  $\Omega$ -algebra.  $\Leftrightarrow$  For  $\omega \in \Omega$  and  $((x_1, x_2, \dots, x_n), c) \in X^n \times X,$

$$\bigwedge \left\{ \left\{ \bigvee [x_i \in A] \vee \omega((x_1, x_2, \dots, x_n), d) \right\} : d \in X \right\} \geq \omega((x_1, x_2, \dots, x_n), c)$$

**Example 3.1.** Let  $A = \langle X, \Omega \rangle$  be a  $H^*$ -valued  $\Omega$ -algebra.

$$\{\Theta\} : A \rightarrow H^*, [x \in \{\Theta\}] = 1_{H^*}$$

is a subset of  $A.$

$$E = \langle \{\Theta\}, \Omega \rangle$$

is a  $H^*$ -valued  $\Omega$ -algebra.  $E$  is called trivial  $H^*$ -valued  $\Omega$ -algebra.

**Definition 3.2.** Let  $A = \langle X, \Omega \rangle$  be a  $H^*$ -valued  $\Omega$ -algebra. If  $K \subseteq X, B : K \rightarrow H^*$  is  $H^*$ -valued set,  $(B, =) \subseteq (A, =)$  and for all  $\omega \in \Omega, \omega \downarrow_B$  satisfy the (1) then  $B$  is  $H^*$ -valued  $\Omega$ -subalgebra of  $A.$

**Example 3.2.** Let  $A = \langle X, \Omega \rangle$  be a  $H^*$ -valued  $\Omega$ -algebra.  $E = \langle \{\Theta\}, \Omega \rangle$  is  $H^*$ -valued  $\Omega$ -subalgebra.

**Definition 3.3.** Let  $A = \langle X, \Omega \rangle$  and  $B = \langle Y, \Omega \rangle$  are similar  $H^*$ -valued  $\Omega$ -algebras and  $f : A \rightarrow B$  be  $H^*$ -valued function.  $f$  is a  $H^*$ -valued  $\Omega$ -algebra homomorphism  $\Leftrightarrow$

- H1  $[x = x] = [f(x) = f(x)]$
- H2  $[x = x'] \leq [f(x) = f(x')]$
- H3  $f(\omega(x_1, x_2, \dots, x_n), y) = \bigvee \{f(x_i, y_i) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0\}$

**Example 3.3.** Let  $A = \langle X, \Omega \rangle$  be a  $H^*$ -valued  $\Omega$ -algebra and  $f : E \rightarrow A, g : E \rightarrow A$  are  $H^*$ -valued functions.  $\forall x, I(\{\Theta\}, x) = 1_{H^*}$   $H^*$ -valued  $\Omega$ -algebra homomorphism exist. This homomorphism is unique.

**Proposition 3.1.** Let  $A, B, C$  are similar  $H^*$ -valued  $\Omega$ -algebras. If  $f : A \rightarrow B, g : B \rightarrow C$  are  $H^*$ -valued  $\Omega$ -algebra homomorphisms then  $(g \circ f) : A \rightarrow C$  is a  $H^*$ -valued  $\Omega$ -algebra homomorphism.

*Proof.* Let  $x_1, x_2, \dots, x_n \in A, z \in C$ ,

$$\begin{aligned}
& (g \circ f)(\omega(x_1, x_2, \dots, x_n), z) = \bigwedge \{f(\omega(x_1, x_2, \dots, x_n), y) \vee g(y, z) : y \in B\} \\
& = \bigwedge \left\{ \bigvee \left\{ f(x_i, y_i) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0 \right\} \right. \\
& \quad \left. \bigvee g(y, z) : y \in B \right\} \\
& = \bigwedge \left\{ \bigvee \left\{ f(x_i, y_i) : i = 1, \dots, n \right\} \vee \right. \\
& \quad \left. \{g(\omega(y_1, y_2, \dots, y_n), z) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0\} \right\} \\
& = \bigwedge \left\{ \bigvee \left\{ \bigvee \left\{ f(x_i, y_i) : i = 1, \dots, n \right\} \vee \right. \right. \\
& \quad \left. \left. \left\{ \bigvee g(y_i, z_i) : z = \omega(z_1, z_2, \dots, z_n), g(y_i, z_i) > 0 : \right\} \right\} \right. \\
& \quad \left. y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0 \right\} \\
& = \bigwedge \left\{ \bigvee \left\{ \left. \begin{array}{l} f(x_i, y_i) \vee g(y_i, z_i) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0, \\ z = \omega(z_1, z_2, \dots, z_n), g(y_i, z_i) > 0 : y_i \in B \end{array} \right\} \right\} \right. \\
& = \bigvee \left\{ \bigwedge \left\{ \left. \begin{array}{l} f(x_i, y_i) \vee g(y_i, z_i) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0, \\ z = \omega(z_1, z_2, \dots, z_n), g(y_i, z_i) > 0 : y_i \in B \end{array} \right\} \right\} \right. \\
& = \bigvee \left\{ \left\{ \left. \begin{array}{l} (g \circ f)(x_i, z_i) : y = \omega(y_1, y_2, \dots, y_n), f(x_i, y_i) > 0, \\ z = \omega(z_1, z_2, \dots, z_n), g(y_i, z_i) > 0 \end{array} \right\} : y_i \in B \right\} \right. \\
& = \bigvee \{(g \circ f)(x_i, z_i) : z = \omega(z_1, z_2, \dots, z_n), (g \circ f)(x_i, z_i) > 0, i = 1, \dots, n\}
\end{aligned}$$

□

#### 4. CONCLUSION

In this paper, we introduced the  $H^*$ -valued  $\Omega$ -algebra and  $H^*$ -valued  $\Omega$ -algebra homomorphism. To study these concepts, firstly we defined  $H^*$ -valued set and  $H^*$ -valued function. In continuation of this study, injection and projection mappings on  $H^*$ -valued  $\Omega$ -algebra can be defined, isomorphism theorems can be proved.

#### REFERENCES

- [1] Birkhoff G., Lattice Theory, American Mathematical Society, United States of America, p.418,1940.
- [2] Faith C., Algebra: Rings, Modules and Categories I, Springer-Verlag, Berlin
- [3] F.V. Lawvere, Intrinsic co-Heyting boundaries and the Leibniz rule in certain toposes, in A.Carboni, et.al., Category theory, Proceedings, Como. 1990.
- [4] Çuvalcıǧlı G., Heyting Deęerli Omega Cebirlerde Serbestlik, Çukurova Üniversitesi Fen Bilimleri Enstitüsü, Doktora Tezi, Adana, 2002, 68 s.
- [5] Çuvalcıođlu G., On Fuzzy  $\Omega$ -word Algebras, Adv. Studies in Contemp. Math. Volume 8(2), 103-110, 2004.

DEPARTMENT OF MATHEMATICS, FACULTY OF SCIENCE AND ARTS, SÜTCÜ İMAM UNIVERSITY,  
KAHRAMANMARAŞ, TURKEY  
*E-mail address:* citil@ksu.edu.tr

MERSİN UNIVERSITY, DEPARTMENT OF MATHEMATICS, MERSİN, TURKEY  
*E-mail address:* sinemyilmaz@gmail.com

## SOME CLASSIFICATIONS OF AN X FDK-SPACE

ŞEYDA SEZGEK AND İLHAN DAĞADUR

ABSTRACT. In this paper, we introduce definitions of double wedge and double conull FDK-spaces. Also, we give some important corollary for any double sequence space about these definitions.

Received: 27–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

In [4], the first study on double sequences was examined by Bromwich. And than it was investigated by many authors such as Hardy [6], Moricz [7], Tripathy [16], Başarır and Sonalcan [2]. The notion of regular convergence for double sequences was defined by Hardy [6]. After that both the theory of topological double sequence spaces and the theory of summability of double sequences were studied by Zeltser [17]. The statistical and Cauchy convergence for double sequences were examined by Mursaleen and Edely [8] and Tripathy [15] in recent years. Many recent improvements containing the summability by four dimensional matrices might be found in [10].

$\Omega$  denotes the space of all complex valued double sequences which is a vector space with coordinatewise addition and scalar multiplication. Any vector subspace of  $\Omega$  is called as a double sequence space. The space  $\mathcal{M}_u$  of all bounded double sequences is defined by

$$\mathcal{M}_u := \left\{ x \in \Omega : \|x\|_\infty := \sup_{k,l} |x_{kl}| < \infty \right\},$$

which is a Banach space with the norm  $\|\cdot\|_\infty$ . In addition we consider the double sequence spaces

$$\begin{aligned} \Phi &:= \text{span}\{e^{kl} : k, l \in \mathbb{N}\} = \{x \in \Omega : \exists N_0 \in \mathbb{N}, \forall (k, l) \in \mathbb{N}^2/[1, N_0]^2 : x_{kl} = 0\} \\ \Phi_1 &:= \Phi \cup \{e\} \\ \mathcal{L}_u &:= \left\{ x \in \Omega : \sum_{k,l} |x_{kl}| < \infty \right\} \end{aligned}$$

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference 2010 *Mathematics Subject Classification.* 40A05, 40C05, 40B05, 40D25.

*Key words and phrases.* Double sequence, FDK-space, wedge FDK-space, conull FDK-space.

$$\begin{aligned} \mathcal{BV} &:= \left\{ x \in \Omega : \|x\|_{\mathcal{BV}} := \sum_{k,l} |x_{kl} - x_{k+1,l} - x_{k,l+1} + x_{k+1,l+1}| < \infty \right\} \\ \mathcal{C}_0 &:= \left\{ x \in \mathcal{C} : \forall l \in \mathbb{N} : \lim_k x_{kl} = 0 \right\} \\ \mathcal{C}_c &:= \left\{ x \in \Omega : \forall l \in \mathbb{N} : \lim_{k,l} x_{kl} \text{ exist} \right\}. \end{aligned}$$

Throughout this paper  $e$  denotes the double sequence of ones;  $(\delta^{ij})$ ,  $i, j = 1, 2, \dots$ , with the one in the  $(i, j)$  position.

A subspace  $E$  of the vector space  $\Omega$  is called DK-space, if all the seminorms  $r_{kl} : E \rightarrow \mathbb{R}, x \mapsto |x_{kl}|$  ( $k, l \in \mathbb{N}$ ) are continuous. An FDK space is a DK-space with a complete, metrizable, locally convex topology. A normable FDK-space is called BDK-space.

## 2. MAIN RESULTS

In this section conull (strongly conull) and double wedge (weak double wedge) FDK-spaces are defined and several characterizations are given.

**Definition 2.1.** If  $(E, \tau)$  is a FDK-space containing  $\Phi$ , and  $\delta^{ij} \rightarrow 0$  in  $\tau$ , then  $(E, \tau)$  is called a double wedge space.

**Definition 2.2.** If  $(E, \tau)$  is a FDK-space containing  $\Phi$ , and  $\delta^{ij} \rightarrow 0$  (weakly) in  $\tau$ , then  $(E, \tau)$  is called a weak double wedge space.

**Definition 2.3.** Let  $(E, \tau)$  is an FDK-space containing  $\Phi_1$ . If  $\forall f \in E'$ ,

$$f(e) = \lim_{m,n \rightarrow \infty} \sum_{k,l=1}^{m,n} f(\delta^{kl})$$


then  $(E, \tau)$  is called a conull FDK-space.

**Definition 2.4.** Let  $(E, \tau)$  is an FDK-space containing  $\Phi_1$ . If  $\forall f \in E'$ ,

$$e = \lim_{m,n \rightarrow \infty} \sum_{k,l=1}^{m,n} \delta^{kl}$$

then  $(E, \tau)$  is called a strong conull FDK-space.

Clearly, each strong conull (wedge) FDK-space is a conull (weak wedge) FDK-space and the following diagram is hold:



Indeed, let  $E$  be a strong conull FDK-space. Then we have  $e^{(kl)} \rightarrow e$ . Hence  $P(e^{(kl)} - e) \rightarrow 0$  ( $k, l \rightarrow \infty$ ) for the seminorm  $P$  in  $\tau$ . In this case, the following

equation is hold

$$\begin{aligned}
\delta^{kl} &= e^{(k,l)} - e^{(k-1,l-1)} - \sum_{i=k,j=1}^{k,l} \delta^{ij} - \sum_{i=1,j=l}^{k,l} \delta^{ij} \\
&= e^{(k,l)} - e^{(k-1,l-1)} - \sum_{i=k,j=1}^{k,l} \delta^{ij} - \sum_{i=1,j=l}^{k,l} \delta^{ij} + e^{(k-1,l-1)} - e^{(k-1,l-1)} \\
&= e^{(k,l)} - e^{(k,l-1)} - e^{(k-1,l)} + e^{(k-1,l-1)} + e - e + e - e .
\end{aligned}$$

So, we have

$$\begin{aligned}
P(\delta^{kl}) &= P\left(e^{(k,l)} - e^{(k,l-1)} - e^{(k-1,l)} + e^{(k-1,l-1)} + e - e + e - e\right) \\
&\leq P\left(e^{(k,l)} - e\right) + P\left(e^{(k,l-1)} - e\right) + P\left(e^{(k-1,l)} - e\right) + P\left(e^{(k-1,l-1)} - e\right).
\end{aligned}$$

It is clearly that since  $P(\delta^{kl}) \rightarrow 0$ ,  $k, l \rightarrow \infty$ ,  $E$  is double wedge space.

We recall that the  $\alpha$ -dual of a subset  $E$  of  $\Omega$  is defined to be ([14])

$$E^\alpha := \left\{ x = (x_{kl}) : \sum_{k,l=1}^{\infty, \infty} |x_{kl} y_{kl}| < \infty, \forall y = (y_{kl}) \in E \right\}.$$

**Lemma 2.1.** *If  $z^{(mn)} \in \mathcal{C}_0$ ,  $m, n = 1, 2, \dots$ , then there exists  $z \in \mathcal{C}_0$  such that*

$$\lim_{i,j \rightarrow \infty} \frac{z_{ij}^{(mn)}}{z_{ij}} = 0 \quad (m, n = 1, 2, \dots).$$

Furthermore, for any such  $z$ , we have  $z^\alpha \subseteq \bigcap_{m,n=1}^{\infty, \infty} \{z^{(mn)}\}^\alpha$ .

*Proof.* Let  $z^{(mn)} \in \mathcal{C}_0$ . We can choose two sequences  $(i_k)$ ,  $(j_l)$  of positive integers such that

$$1 = i_0 < i_1 < i_2 < \dots \quad 1 = j_0 < j_1 < j_2 < \dots$$

and

$$\max_{\substack{1 \leq m \leq k \\ 1 \leq n \leq l}} |z_{ij}^{(mn)}| < \frac{1}{4^{kl}} \quad \left( \begin{array}{l} i \geq i_k, j \geq j_l \\ k, l = 1, 2, \dots \end{array} \right).$$

Define  $z \in \Omega$  as follows:

$$z_{ij} = \frac{1}{2^{kl}} \quad \left( \begin{array}{l} i_k \leq i < i_{k+1}, j_l \leq j < j_{l+1} \\ k, l = 0, 1, 2, \dots \end{array} \right).$$

Clearly,  $z \in \mathcal{C}_0$  and, fixing  $m, n$

$$\left| \frac{z_{ij}^{(mn)}}{z_{ij}} \right| < \frac{1}{2^{kl}} \quad \left( \begin{array}{l} i \geq i_k, j \geq j_l \\ k \geq m, l \geq n \end{array} \right).$$

Thus,  $\lim_{i,j \rightarrow \infty} \frac{z_{ij}^{(mn)}}{z_{ij}} = 0 \quad \forall m, n$ .

Now let  $x \in z^\alpha$ , then

$$\sum_{i,j=1}^{\infty, \infty} |x_{ij} z_{ij}| < \infty.$$

Moreover, for  $m, n = 1, 2, \dots$  we have

$$\begin{aligned} \sum_{i,j=1}^{\infty, \infty} |x_{ij} z_{ij}^{(mn)}| &< \sum_{i,j=1}^{\infty, \infty} |x_{ij} z_{ij}| \frac{1}{2^{kl}} \\ &< \frac{1}{2^{kl}} \sum_{i,j=1}^{\infty, \infty} |x_{ij} z_{ij}| < \infty. \end{aligned}$$

Hence,  $x \in \bigcap_{m,n=1}^{\infty, \infty} \{z^{(mn)}\}^\alpha$ . The proof is completed.  $\square$

$s = (s_m), t = (t_n)$  always denote strictly increasing of nonnegative integers with  $s_1 = 0, t_1 = 0$ . We will be interested in spaces of the form:

$$m|(s, t)| = \left\{ x \in \Omega : \sup_{m,n} \sum_{\substack{k=s_m+1 \\ l=t_n+1}}^{s_{m+1}, t_{n+1}} |x_{kl}| < \infty \right\}$$

which becomes a BDK-space under the norm:

$$x \rightarrow \sup_{m,n} \sum_{\substack{k=s_m+1 \\ l=t_n+1}}^{s_{m+1}, t_{n+1}} |x_{kl}|.$$

**Theorem 2.1.** *Let  $(E, \tau)$  be an FDK-space. These are equivalent:*

- i)  $E$  is a double wedge space,
- ii)  $E$  contains  $z^\alpha$  for some  $z \in \mathcal{C}_0$ ,
- iii)  $E$  contains  $m|(s, t)|$  for some  $s, t$  and the inclusion mapping is compact,
- iv)  $E$  contains  $\mathcal{L}_u$  and the inclusion mapping is compact.

*Proof.* ( $i \Rightarrow ii$ ) Let  $\{p_{mn}\}$  be a defining family of seminorms for the topology  $\tau$  and let

$$z_{ij}^{(mn)} = p_{mn}(\delta^{ij}) \quad (m, n, i, j = 1, 2, \dots).$$

Then  $z^{(mn)} \in \mathcal{C}_0$ ,  $m, n = 1, 2, \dots$ , since  $E$  is a double wedge space. Suppose  $y \in \bigcap_{m,n=1}^{\infty, \infty} \{z^{(mn)}\}^\alpha$ , then for each  $m, n$

$$\sum_{i,j} |y_{ij} z_{ij}^{(mn)}| < \infty.$$

Therefore,

$$\sum_{i,j} |y_{ij} p_{mn}(\delta^{ij})| = \sum_{i,j} p_{mn}(y_{ij} \delta^{ij}) < \infty$$

is obtained. Since the space  $E$  is complete  $\sum_{i,j} y_{ij} \delta^{ij}$  converges in  $(E, \tau)$  to, say  $x$ , or  $y^{(mn)} \rightarrow x$ . Thus  $y_{ij}^{(mn)} \rightarrow x_{ij}$  for each  $i, j$ ; also we always have  $y_{ij}^{(mn)} \rightarrow y_{ij}$  for each  $i, j$ . Consequently  $y = x$ ; that is

$$\bigcap_{m,n=1}^{\infty, \infty} \{z^{(mn)}\}^\alpha \subseteq E.$$

Choosing  $z$  as in Lemma 2.1, (ii) follows.



(ii  $\Rightarrow$  iii) Let us choose strictly increasing sequences  $(s_m), (t_n)$  of positive integers such that  $s_1 = 0, t_1 = 0$  and

$$|z_{ij}| \leq \frac{1}{2^{mn}}$$

whenever  $i \geq s_m, j \geq t_n, m, n \geq 2$ .

For  $x \in m|(s, t)|$  and any positive integers  $k, l, u, v$  such that  $l \geq k$  and  $v \geq u$  we have

$$\sum_{\substack{i=s_k+1 \\ j=t_p+1}}^{s_{l+1}, t_{v+1}} |x_{ij}z_{ij}| = \sum_{m=k, n=u}^{l, v} \sum_{\substack{i=s_m+1 \\ j=t_n+1}}^{s_{m+1}, t_{n+1}} |x_{ij}z_{ij}| \leq \|x\| \sum_{m=k, n=u}^{l, v} \frac{1}{2^{mn}}.$$

Hence  $x \in z^\alpha$ . That is;  $m|(s, t)| \subseteq z^\alpha \subseteq E$ . Also, the inclusion theorem  $i : (m|(s, t)|, \|\cdot\|) \rightarrow (E, \tau)$  is compact.

(iii  $\Rightarrow$  iv) Since  $\mathcal{L}_u \subset m|(s, t)|$  always true, the inclusion theorem  $i : \mathcal{L}_u \rightarrow m|(s, t)|$  is continuous. By hypothesis  $\mathcal{L}_u \subset m|(s, t)| \subset E$  and because of  $i : (m|(s, t)|, \|\cdot\|) \rightarrow (E, \tau)$  is compact  $i^* : \mathcal{L}_u \rightarrow E$  is compact.  $\square$

**Corollary 2.1.** *The intersection of all double wedge FDK-spaces is  $\mathcal{L}_u$ .*

*Proof.* Let  $E$  be a double wedge FDK-space. Then we have

$$\bigcap E = \bigcap \{z^\alpha : z \in \mathcal{C}_0\} = \mathcal{C}_0^\alpha.$$

Now we need show that  $\mathcal{C}_0^\alpha = \mathcal{L}_u$ . Since  $\mathcal{C}_0 \subset \mathcal{M}_u, \mathcal{M}_u^\alpha \subset \mathcal{C}_0^\alpha$ . That is,  $\mathcal{L}_u = \mathcal{M}_u^\alpha \subset \mathcal{C}_0^\alpha$  is obtained. Suppose that  $x = (x_{kl}) \in \mathcal{C}_0^\alpha$  but  $x = (x_{kl}) \notin \mathcal{L}_u$ . Then for  $y = (y_{kl}) = \frac{1}{k!l!} \in \mathcal{C}_0$  we have

$$\sum |x_{kl}y_{kl}| = \sum |x_{kl} \frac{1}{k!l!}| = e^2 \sum |x_{kl}| = \infty.$$

This means that  $x = (x_{kl}) \notin \mathcal{C}_0^\alpha$  which contradicts the hypothesis. So,  $x = (x_{kl})$  must be in  $\mathcal{L}_u$ . That is,  $\mathcal{L}_u \subset \mathcal{C}_0^\alpha$ . This completes the proof.  $\square$

The one to one mapping  $S^{(2)}$  of  $\Omega$  to itself defined by

$$S^{(2)}x = \begin{pmatrix} x_{11} & x_{11} + x_{12} & \dots & \dots & \dots \\ x_{11} + x_{21} & x_{11} + x_{12} + x_{21} + x_{22} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & \sum_{k,l=1}^{m,n} x_{kl} & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

$$(S^{(2)})^{-1}x = \begin{pmatrix} x_{11} & x_{12} - x_{11} & \dots & \dots & \dots \\ x_{21} - x_{11} & x_{22} - x_{12} - x_{21} + x_{11} & \dots & \dots & \dots \\ \dots & \dots & \dots & \dots & \dots \\ \dots & \dots & \dots & x_{mn} - x_{m,n-1} - x_{m-1,n} + x_{m-1,n-1} & \dots \\ \dots & \dots & \dots & \dots & \dots \end{pmatrix}$$

**Theorem 2.2.** *i)  $(E, \tau)$  is strongly conull FDK-space if and only if the space  $(S^{(2)})^{-1}(E)$  is a double wedge FDK-space.*

*ii)  $(E, \tau)$  is conull FDK-space if and only if the space  $(S^{(2)})^{-1}(E)$  is a weak double wedge FDK-space.*

*Proof.* i) Let  $(E, \tau)$  is strongly conull FDK-space and  $\{P_{mn}\}$  is a set of seminorms on the topology  $\tau$ . Then a topology with the set of seminorms  $\{q_{mn}\}$  make  $(S^{(2)})^{-1}(E)$  is a FDK-space such that

$$q_{mn}(x) := P_{mn}(S^{(2)}(x)).$$

Since  $(E, \tau)$  is strongly conull FDK-space, for all  $m, n \in \mathbb{N}$   $P_{mn}(e - e^{(mn)}) \rightarrow 0$ . Otherwise,

$$S^{(2)}x = e - e^{(mn)} = \sum_{\substack{k=m+1 \\ l=n+1}}^{\infty, \infty} \delta^{kl} + \sum_{\substack{k=1 \\ l=n+1}}^{m, \infty} \delta^{kl} + \sum_{\substack{k=m+1 \\ l=1}}^{\infty, n} \delta^{kl}$$

is hold. Thus we have

$$\begin{aligned} x &= (S^{(2)})^{-1} \left( \sum_{\substack{k=m+1 \\ l=n+1}}^{\infty, \infty} \delta^{kl} + \sum_{\substack{k=1 \\ l=n+1}}^{m, \infty} \delta^{kl} + \sum_{\substack{k=m+1 \\ l=1}}^{\infty, n} \delta^{kl} \right) \\ &= (S^{(2)})^{-1} \begin{pmatrix} 0 & \dots & 0 & 1 & \dots & 1 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 0 & \dots & 0 & 1 & \dots & 1 \\ 1 & \dots & 1 & 1 & \dots & 1 \\ \dots & \dots & \dots & \dots & \dots & \dots \\ 1 & \dots & \dots & \dots & \dots & 1 \end{pmatrix} \\ &= \delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}. \end{aligned}$$

From the definition of  $(q_{mn})$  the following equation is obtained:

$$q_{mn}(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) = P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right).$$

Since  $P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right) \rightarrow 0$  ( $m, n \rightarrow \infty$ ), we can say  $q_{mn}(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) \rightarrow 0$  ( $m, n \rightarrow \infty$ ).

In this case, we have  $(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) \rightarrow 0$  ( $m, n \rightarrow \infty$ ) according to the topology of the space  $(S^{(2)})^{-1}(E)$ . Since  $\delta^{m+1,1} \rightarrow 0$  and  $\delta^{1,n+1} \rightarrow 0$ ,  $\delta^{m+1,n+1} \rightarrow 0$  is hold. This gives that  $(S^{(2)})^{-1}(E)$  is a double wedge FDK-space.

Now we suppose that  $(S^{(2)})^{-1}(E)$  is a double wedge FDK-space. Then we have for all  $m, n \in \mathbb{N}$   $q_{mn}(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) \rightarrow 0$  ( $m, n \rightarrow \infty$ ). Since

$$q_{mn}(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) = P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right),$$

$P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right) \rightarrow 0$  ( $m, n \rightarrow \infty$ ) is obtained. So  $E$  strongly conull FDK-space.

ii) Let  $(E, \tau)$  is conull FDK-space and let us define the topology of  $(S^{(2)})^{-1}(E)$  as the proof of (i). Then  $q_{mn}(x) := P_{mn}(S^{(2)}(x))$  and since  $(S^{(2)})^{-1} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right) = \delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}$ , we have

$$q_{mn}(\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1}) = P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right).$$

Because of  $E$  is conull FDK space,  $P_{mn} \left( e - \sum_{k,l=1}^{m,n} \delta^{kl} \right) \rightarrow 0$  (weak) ( $m, n \rightarrow \infty$ ). Hence  $\delta^{m+1,1} + \delta^{1,n+1} - \delta^{m+1,n+1} \rightarrow 0$  (weak) ( $m, n \rightarrow \infty$ ). Consequently  $\delta^{m+1,n+1} \rightarrow 0$  (weak) ( $m, n \rightarrow \infty$ ) is obtained. That is,  $(S^{(2)})^{-1}(E)$  is a weak double wedge FDK-space. The other hand of the proof is as the proof of (i).  $\square$

## REFERENCES

- [1] Altay B., Baar F., *Some new spaces of double sequences* J. Math. Anal. Appl. 309 (2005) 70-90.
- [2] Başarır M., Sonalcan O.: *On Some Double Sequence Spaces*, J. Indian Acad. Math., 21 (1999), 193-200.
- [3] Bennett G. , *A New Class of Sequence Spaces with Applications in Summability Theory*, J. Reine Agnew. Math., 266 (1974), 49-75.
- [4] Bromwich T. J. , *An Introduction to The Theory of Infinite Series* , Macmillan and Co. Ltd., New York (1965).
- [5] Goffman C., Petersen G. M., *Submethods of Regular Matrix Summability Methods*, Canad. J. Math. 8, 40-46 (1956).
- [6] Hardy G. H., *On The Convergence of Certain Multiple Series*, Proc.London Math. Soc., s 2-1(1)(1904), 124-128.
- [7] Moricz F., *Extension of The Spaces  $c$  and  $c_0$  from Single to Double Sequence*, Acta Math. Hungarica, 57(1991), 129-136.
- [8] Mursaleen, M. and Edely ,Osama H.H., *Statistical Convergence of Double sequences*, J.Math. Anal. Appl. 288(2003)223-231.
- [9] Mursaleen M., Mohiuddine S. A., *Convergence Methods for Double Sequences and Applications*, Springer (2014).
- [10] Patterson R. F., *Analogues of Some Fundamental Theorems of Summability Theory*, Int. J. Math. Math. Sci., 23, (2000) 1-9.
- [11] Pringsheim A., *Elementare Theorie der Unendliche Doppelreihen*, Sitzungs Berichte der Math. Akad. der Wissenschaften zu MÜch. Ber. 7, 101-153 (1898).
- [12] Robison G. M. , *Divergent Double sequences and series*, Transactions of the American Mathematical Society, 28, no .1, (1926), 50-73.
- [13] Snyder A. K., *Conull and coregular FK-spaces* , Math. Z., 90 (1965), 376-381.
- [14] Tripathy B. C., Sarma B., *Generalized Köthe-Toeplitz Dual of Some Double Sequence Spaces*, Fasciculi Mathematici 40(2008), 119-125.
- [15] Tripaty B. C., *Statistically Convergent Double Sequences*, Tamkang J. Math., 34, (2003), 230-237.
- [16] Tripaty B. C., *Generalized Difference Paranormed Statistically Convergent Sequences Defined by Orlicz Function in a Locally Convex Spaces*, Soochow J. Math., 30 (2004), 431-446.
- [17] Zeltser M., *Investigation of Double Sequence Spaces by Soft and Hard Analytical Methods*, Diss. Math. Univ. Tartu. 25, Tartu University Press, Univ. of Tartu, Faculty of Mathematics and Computer Science, Tartu (2001).

MERSIN UNIVERSITY, DEPARTMENT OF MATHEMATICS MERSIN / TURKEY

*E-mail address:* seydaezgek@gmail.com

*E-mail address:* ilhandagadur@yahoo.com

## A NOTE ON POROSITY CLUSTER POINTS

MAYA ALTINOK AND MEHMET KÜÇÜKASLAN

ABSTRACT. Porosity cluster points of real valued sequences was defined and studied in [3]. In this paper we give the relation between porosity cluster points and distance function.

Received: 17–July–2016

Accepted: 29–August–2016

### 1. INTRODUCTION

Porosity is appeared in the papers of Denjoy [5], [6], Khintchine [11] and, Dolzenko [7]. It has many applications in theory of free boundaries [10], generalized subharmonic functions [8], complex dynamics [12], quasisymmetric maps [14], infinitesimal geometry [4] and other areas of mathematics.

Let  $A \subset \mathbb{R}^+ = [0, \infty)$ , then the right upper porosity of  $A$  at the point 0 is defined as

$$p^+(A) := \limsup_{h \rightarrow 0^+} \frac{\lambda(A, h)}{h}$$

where  $\lambda(A, h)$  denotes the length of the largest open subinterval of  $(0, h)$  that contains no point of  $A$  (for more information look [13]). The notion of right lower porosity of  $A$  at the point 0 is defined similarly.

In [1], the notation of porosity which was defined at zero for the subsets of real numbers, has been redefined at infinity for the subsets of natural numbers.

Let  $\mu : \mathbb{N} \rightarrow \mathbb{R}^+$  be a strictly decreasing function such that  $\lim_{n \rightarrow \infty} \mu(n) = 0$ , (it is called scaling function) and let  $E$  be a subset of  $\mathbb{N}$ .

Upper porosity and lower porosity of the set  $E$  at infinity were defined respectively in [1] as follows:

$$(1.1) \quad \bar{p}_\mu(E) := \limsup_{n \rightarrow \infty} \frac{\lambda_\mu(E, n)}{\mu(n)}, \quad p_\mu(E) := \liminf_{n \rightarrow \infty} \frac{\lambda_\mu(E, n)}{\mu(n)},$$

where

$$\lambda_\mu(E, n) := \sup\{|\mu(n^{(1)}) - \mu(n^{(2)})| : n \leq n^{(1)} < n^{(2)}, (n^{(1)}, n^{(2)}) \cap E = \emptyset\}.$$

Using the definition of upper porosity, all subsets of natural numbers can be classify as follows:  $E \subseteq \mathbb{N}$  is

- (i) porous at infinity if  $\bar{p}_\mu(E) > 0$ ;
- (ii) strongly porous at infinity if  $\bar{p}_\mu(E) = 1$ ;

---

<sup>1</sup>3<sup>rd</sup> International Intuitionistic Fuzzy Sets and Contemporary Mathematics Conference  
 2010 *Mathematics Subject Classification.* 40A05, 14F45.

*Key words and phrases.* local upper porosity, porosity of subsets of natural numbers, porosity convergence, porosity limit points, porosity cluster points .

(iii) nonporous at infinity if  $\bar{p}_\mu(E) = 0$ . Throughout this paper, we will consider only the upper porosity of subsets of  $\mathbb{N}$ .

Let us recall the definition of  $\bar{p}_\mu$ -convergence of real valued sequences for any scaling function:

**Definition 1.1.** [2] A sequence  $x = (x_n)_{n \in \mathbb{N}}$  is said to  $\bar{p}_\mu$ -convergent to  $l$  if for each  $\varepsilon > 0$ ,

$$\bar{p}_\mu(A_\varepsilon) > 0,$$

where  $A_\varepsilon := \{n : |x_n - l| \geq \varepsilon\}$ . It is denoted by  $x \rightarrow l(\bar{p}_\mu)$  or  $(\bar{p}_\mu - \lim_{n \rightarrow \infty} x_n = l)$ .

Let  $x' = (x_{n_k})$  be a subsequence of  $x = (x_n)$  for monotone increasing sequence  $(n_k)_{k \in \mathbb{N}}$  and  $K := \{n_k : k \in \mathbb{N}\}$ , then we abbreviate  $x' = (x_{n_k})$  by  $(x)_K$ .

**Definition 1.2.** [3]. Let  $x = (x_n)$  be a sequence and  $(x)_K$  be a subsequence of  $x = (x_n)$ . If

- (i)  $\bar{p}_\mu(K) > 0$ , then  $(x)_K$  is called  $\bar{p}_\mu$ -thin subsequence of  $x = (x_n)$ ,
- (ii)  $\bar{p}_\mu(K) = 1$ , then  $(x)_K$  is called a strongly  $\bar{p}_\mu$ -thin subsequence of  $x = (x_n)$ ,
- (iii)  $\bar{p}_\mu(K) = 0$ , then  $(x)_K$  is a  $\bar{p}_\mu$ -nonthin (or  $\bar{p}_\mu$ -dense) subsequence of  $x = (x_n)$ .

**Definition 1.3.** [3]. A number  $\alpha$  is said to be  $\bar{p}_\mu$ -limit point of the sequence  $x = (x_n)$  if it has a  $\bar{p}_\mu$ -nonthin subsequence that converges to  $\alpha$ .

The set of all  $\bar{p}_\mu$ -limit points of  $x = (x_n)$  is denoted by  $L_{\bar{p}_\mu}(x)$ .

**Definition 1.4.** [3]. A number  $\beta$  is said to be a  $\bar{p}_\mu$ -cluster point of  $x = (x_n)$  if for every  $\varepsilon > 0$ , the set

$$\{n : |x_n - \beta| < \varepsilon\}$$

is nonporous. i.e.,

$$\bar{p}_\mu(\{n : |x_n - \beta| < \varepsilon\}) = 0.$$

For a given sequence  $x = (x_n)$ ; the symbol  $\Gamma_{\bar{p}_\mu}(x)$  denotes the set of all  $\bar{p}_\mu$ -cluster points.

## 2. MAIN RESULTS

Some results about the set of  $L_{\bar{p}_\mu}(x)$ -cluster, and  $L_{\bar{p}_\mu}(x)$ -limit points of given real valued sequences has been investigated in [3]. In this paper, as a continuation of [3] the same subject will be studied.

**Theorem 2.1.** *Assume that  $x = (x_n)$  is monotone increasing (or decreasing) sequence of real numbers. If  $\sup x_n < \infty$  (or  $\inf x_n < \infty$ ), then  $\sup x_n \in \Gamma_{\bar{p}_\mu}(x)$  (or  $\inf x_n \in \Gamma_{\bar{p}_\mu}(x)$ ).*

*Proof.* The proof will be given only for monotone increasing sequences. The other case can be proved by following similar steps. From the definition of supremum for any  $\varepsilon > 0$ , there exists an  $n_0 = n_0(\varepsilon) \in \mathbb{N}$  such that the inequality

$$\sup x_n - \varepsilon < x_{n_0} \leq \sup x_n$$

holds. Since the sequence is monotone increasing, then we have

$$(2.1) \quad \sup x_n - \varepsilon < x_{n_0} < x_n \leq \sup x_n < \sup x_n + \varepsilon$$

for all  $n > n_0$ .

From (2.1), for any  $\varepsilon > 0$  there exists an  $n_0 = n_0(\varepsilon) \in \mathbb{N}$  such that following inequality

$$(2.2) \quad |x_n - \sup x_n| < \varepsilon$$

holds for all  $n > n_0(\varepsilon)$ .

From (2.2), following inclusion

$$\mathbb{N} \setminus \{1, 2, 3, \dots, n_0\} \subset \{n : |x_n - \sup x_n| < \varepsilon\}$$

and the inequality

$$\bar{p}_\mu(\{n : |x_n - \sup x_n| < \varepsilon\}) \leq \bar{p}_\mu(\mathbb{N} \setminus \{1, 2, \dots, n_0\})$$

hold.

Since  $\bar{p}_\mu(\mathbb{N} \setminus \{1, 2, 3, \dots, n_0\}) = 0$ , then from Lemma 1.1 in [3] we have

$$\bar{p}_\mu(\{n : |x_n - \sup x_n| < \varepsilon\}) = 0.$$

This gives the desired proof.  $\square$

**Corollary 2.1.** *If  $x = (x_n)$  is a bounded sequence, then  $\sup x_n$  and  $\inf x_n$  are belong to  $\Gamma_{\bar{p}_\mu}(x)$ .*

Let  $A, B \subset \mathbb{R}$  and recall the distance between  $A$  and  $B$  is defined as

$$d(A, B) := \inf\{|a - b| : a \in A, b \in B\}.$$

**Theorem 2.2.** *Let  $x = (x_n)$  be a real valued sequence. If  $\Gamma_{\bar{p}_\mu}(x) \neq \emptyset$ , then  $d(\Gamma_{\bar{p}_\mu}(x), x) = 0$ .*

*Proof.* Assume  $\Gamma_{\bar{p}_\mu}(x) \neq \emptyset$ . Let us consider an arbitrary element  $y^* \in \Gamma_{\bar{p}_\mu}(x)$ . Then, for an arbitrary  $\varepsilon > 0$  we have

$$\bar{p}_\mu(\{n : |x_n - y^*| < \varepsilon\}) = 0.$$

So, the set  $\{x_n : |x_n - y^*| < \varepsilon\}$  has at least countable number elements of  $x = (x_n)$ . Let us denote this set by  $D$  where  $D := \{n_k : |x_{n_k} - y^*| < \varepsilon\} \subset \mathbb{N}$ . Therefore, we have

$$\begin{aligned} 0 \leq \text{dist}(\Gamma_{\bar{p}_\mu}(x), x) &= \inf\{|y - x_n| : y \in \Gamma_{\bar{p}_\mu}(x), n \in \mathbb{N}\} \\ &\leq \inf\{|y^* - x_{n_k}| : n_k \in D\} < \varepsilon. \end{aligned}$$

So, for every  $\varepsilon > 0$ , we have  $0 \leq d(\Gamma_{\bar{p}_\mu}(x), x) < \varepsilon$ .  $\square$

**Theorem 2.3.** *Let  $x = (x_n)$  be a real valued sequence and  $\gamma \in \mathbb{R}$  be an arbitrary fixed point. If  $d(\gamma, x) \neq 0$ , then  $\gamma \notin \Gamma_{\bar{p}_\mu}(x)$ .*

*Proof.* From the hypothesis we have

$$d(\gamma, x) := \inf\{|x_k - \gamma| : k \in \mathbb{N}\} = m > 0.$$

From the assumption the inequality

$$|x_k - \gamma| \geq m$$

holds for all  $k \in \mathbb{N}$ . It means that the open interval  $(\gamma - m, \gamma + m)$  has no elements of the sequence  $x = (x_n)$ . So, we have

$$\bar{p}_\mu(\{k : |x_k - \gamma| < m\}) = 1.$$

For  $0 < \varepsilon < m$  we have

$$\bar{p}_\mu(\{k : |x_k - \gamma| < \varepsilon\}) = 1.$$

So,  $\gamma \notin \Gamma_{\bar{p}_\mu}(x)$ . □

*Remark 2.1.* If  $d(\gamma, x) = 0$ , it is not necessary for  $\gamma \in \Gamma_{\bar{p}_\mu}(x)$ .

Let us consider  $(x_n) = (\frac{1}{n})_{n \in \mathbb{N}}$ . For  $\gamma = \frac{1}{2}$ , it is clear that  $d(\frac{1}{2}, \frac{1}{n}) = 0$  holds, but  $\frac{1}{2} \notin \Gamma_{\bar{p}_\mu}(x) = \{0\}$ .

#### REFERENCES

- [1] M. Altınok, O. Dovgoshey, M. Küçükcaslan, *Local one-sided porosity and pretangent spaces*, Analysis, **36(3)** (2016), 147-171.
- [2] M. Altınok, M. Küçükcaslan, *On porosity-convergence of real valued sequences*, (Accepted in Alexandru Ioan Cuza Dn Ia Matematica in 2016).
- [3] M. Altınok, *Porosity limit and cluster points of real valued sequences*, (Submitted 2016).
- [4] V. Bilet, O. Dovgoshey, *Boundedness of pretangent spaces to general metric spaces*, Ann. Acad. Sci. Fenn. Math., **39** (2009), 73-82.
- [5] A. Denjoy, *Sur une propriété des séries trigonométriques*, Verlag v.d. G. V. der Wie-en Natuur. Afd., **29** (1920), 220-232.
- [6] A. Denjoy, *Leçons sur le calcul des coefficients d'une série trigonométrique, Part II, Métrique et topologie d'ensembles parfaits et de fonctions*, Gauthier-Villars, Paris, 1941.
- [7] E. P. Dolženko, *Boundary properties of arbitrary functions*, (Russian), Izv. Akad. Nauk SSSR Ser. Math. **31** (1967), 3-14.
- [8] O. Dovgoshey and J. Riihentausta, *Mean value type inequalities for quasilinearly subharmonic functions*, Glasgow Math. J., **55(2)** (2013), 349-368.
- [9] J. A. Fridy *Statistical limit points*, Proceeding of the American Mathematical Society, **118(4)** (1993), 1187-1192.
- [10] L. Karp, T. Kilpenläinen, A. Petrosyan and H. Shahgholian, *On the porosity of free boundaries in degenerate variational inequalities*, J. Differential Equations, **164** (2000), 110-117.
- [11] A. Khintchine, *An investigation of the structure of measurable functions*, (Russian), Mat. Sbornik, **31** (1924), 265-285.
- [12] F. Przytycki and S. Rohde, *Porosity of Collet-Eckmann Julia sets*, Fund. Math., **155** (1998), 189-199.
- [13] B. S. Thomson, *Real Functions*, Lecture Notes in Mathematics, **1170**, Springer-Verlag, Berlin, Heidelberg, New York, Tokyo, 1985.
- [14] J. Väisälä, *Porous seta and quasisymmetric maps*, Trans. Amer. Math. Soc., **299** (1987), 525-533.

MERSIN UNIVERSITY, FACULTY OF SCIENCES AND ARTS DEPARTMENT OF MATHEMATICS, 33343 MERSIN, TURKEY

*E-mail address:* [mayaaltinok@mersin.edu.tr](mailto:mayaaltinok@mersin.edu.tr)

*E-mail address:* [mkkaslan@gmail.com.tr](mailto:mkkaslan@gmail.com.tr)

196

IFSCOM2016

**SECTION II  
ABSTRACTS**



## Fracral Reconstruction of Dilaton Field

**Mustafa SALTİ, Oktay AYDOĞDU**

Department of Physics, Faculty of Arts and Science, Mersin University,  
Mersin-33343, Turkey

msalti@mersin.edu.tr, oktaydogdu@mersin.edu.tr

### Abstract

Numerous papers have been presented[1,2,3,4,5] to implement the dynamics of scalar field describing nature of the dark energy by establishing a connection between the pilgrim/new agegraphic/Ricci/ghost/holographic energy density and a scalar field definition. These works showed that the analytical form of potential in terms of the scalar field cannot be obtained due to the complexity of the involved equations. On the other hand, writing a meaningful quantum gravity theory is one of the tough puzzles in modern theoretical physics[6,7]. In the quantum gravity theories, the universe is described as a dimensional flow and one can discuss whether and how these attractive features are connected with the ultraviolet-divergence problem[8]. That's why, such important points motivated us to reconstruct the potential and dynamics of the dilaton scalar field model[9] according to the evolutionary behavior of the extended holographic energy description[10] in fractal geometry.

### Referances

- [1] Sahni, V., *Dark Matter and Dark Energy*, Lect. Notes Phys. Vol.653, 141-180, 2004.
- [2] Chiba, T., Okabe, T. and Yamaguchi, M., *Kinetical Driven Quintessence*, Phys. Rev. D, Vol.62, No.2, 023511-02351, 2000.
- [3] Feng, B., Wang, X.L. and Zhang, X.M., *Dark energy constraints from the cosmic age and supernova*, Phys. Lett. B, Vol.607, Nos.1-2, 35-41, 2005.
- [4] Caldwell, R.R., *A phantom menace? Cosmological consequences of a dark energy component with super-negative equation of state*, Phys. Lett. B, Vol.545, Nos.1-2, 23-29, 2002.
- [5] Copeland, E.J., Sami, M. and Tsujikawa, S., *Dynamics of dark energy*, Int. J. Mod. Phys. D, Vol.15, No.11, 1753-1935, 2006.
- [6] Calcagni, G., *Fractal Universe and Quantum Gravity*, Phys. Rev. Lett. Vol.104, No.25, 251301, 2010.
- [7] Calcagni, G., *Quantum field theory, gravity and cosmology in a fractal universe*, JHEP, Vol.2010(3), No.120, 1-38, 2010.
- [8] Salti, M. and Aydogdu, A., *Holographic Tachyon in Fractal Geometry*, Math. Comput. Appl. Vol. 21, No.21, 1-12, 2016.

[9] Piazza, F. and Tsujikawa, S., *Dilatonic ghost condensate as dark energy*, JCAP, Vol.2004(7), No.004, 1-28, 2004.

[10] Miao, L., Xiao-Dong, L., Shuang, W. and Yi, W., *Dark Energy*, Communications in Theoretical Physics, Vol.56, No.3, 525-604, 2011.

## The Cauchy Problem for Complex Intuitionistic Fuzzy Differential Equations

A. EL ALLAOU, S. MELLIANI

LMACS, Laboratoire de Mathématiques Appliquées & Calcul Scientifique  
Sultan Moulay Slimane University, PO Box 523, 23000 Beni Mellal  
Morocco  
said.melliani@gmail.com

### Abstract

In this paper, we discuss the existence of a solution to the Cauchy problem for complex intuitionistic fuzzy differential equations. We first propose definitions of complex intuitionistic fuzzy sets and discuss entailed results which parallel those of complex fuzzy sets.

**Keywords :** complex intuitionistic fuzzy sets, complex intuitionistic fuzzy differential equations.

### References

- [1] K. Atanassov, (1986), *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, 20, 87-96.
- [2] K. Atanassov, (1999), *Intuitionistic Fuzzy Sets : Theory and Applications*, Springer Physica-Verlag, Heidelberg.
- [3] K. Atanassov, T., Vassilev, P. M., and Tsvetkov, R. T. (2013), *Intuitionistic Fuzzy Sets*, Mea-sures and integrals Bulgarian Academic Monographs (12), Professor Marin Drinov Academic Publishing House, Sofia.
- [4] R. Ettoussi, S. Melliani, M. Elomari and L. S. Chadli, *Solution of intuitionistic fuzzy differential equations by successive approximations method*, 19th Int. Workshop on IFSs, Burgas, 4-6 June 2015 Notes on Intuitionistic Fuzzy Sets ISSN 1310-4926 Vol. 21, 2015, No. 2, 51-62.
- [5] M. Elomari, S. Melliani, R. Ettoussi and L. S. Chadli, *Intuitionistic fuzzy semigroup*, 19th Int. Workshop on IFSs, Burgas, 4-6 June 2015 Notes on Intuitionistic Fuzzy Sets ISSN 1310-4926 Vol. 21, 2015, No. 2, 43-50.

- [6] S. Melliani, M. Elomari and R. Ettoussi, L. S. Chadli, *Intuitionistic fuzzy metric space*, Notes on Intuitionistic Fuzzy Sets ISSN 1310-4926 Vol. 21, 2015, No. 1, 43-53.
- [7] J. Nieto, *The Cauchy problem for continuous fuzzy differential equations*, Fuzzy Sets Syst. 102 (1999) 259-262.
- [8] D. Ramot, R. Milo, M. Friedman, A. Kandel, *Complex fuzzy sets*, IEEE Trans. Fuzzy Syst. 10 (2002) 171-186.
- [9] D.E. Tamir, L. Jin, A. Kandel, *A new interpretation of complex membership grade*, Int. J. Intell. Syst. 26 (2011) 285-312.

## Numerical Solution of Intuitionistic Fuzzy Differential Equations by Runge-Kutta Method of Order Four

**B. Ben AMMA, L. S. CHADLI**

LMACS, Laboratory of Applied Mathematics and Scientific Calculus  
Sultan Moulay Slimane University, PO Box 523, 23000 Beni Mellal  
Morocco  
bouchrabenamma@gmail.com

### Abstract

This paper presents solution for first order fuzzy differential equation by Runge-Kutta method of order four. This method is discussed in detail and this is followed by a complete error analysis. The accuracy and efficiency of the proposed method is illustrated by solving an intuitionistic fuzzy initial value problem.

**Keywords :** intuitionistic fuzzy Cauchy problem, Runge-Kutta method of order four.

### Referances

- [1] K. Atanassov, Intuitionistic fuzzy sets, Fuzzy Sets and Systems **20**(1986), pp. 87-96.
- [2] K. Atanassov, Intuitionistic Fuzzy Sets, Springer Physica-Verlag, Berlin, 1999.
- [3] J.J Buckley , and Feuring T., Fuzzy differential equations, Fuzzy Sets and Systems, **110**(2000), pp. 43-54.
- [4] B. Ben Amma, S. Melliani and L.S. Chadli, Numerical Solution Of Intuitionistic Fuzzy Differential Equations By Euler and Taylor Method (submitted on Notes on Intuitionistic Fuzzy Sets)**21**(2)(2016).

- [5] C. Duraisamy , and Usha B., Numerical Solution of Differential Equation, by Runge-Kutta Method of Order Four, European Journal of Scientific Research, **67**(2012), pp. 324-337.
- [6] M. Friedman , and Kandel A., Numerical solution of fuzzy differential equations, Fuzzy Sets and Systems, **105**(1999), pp. 133-138.
- [7] S. Melliani, M. Elomari, L.S. Chadli and R. Ettoussi,, Intuitionistic Fuzzy Metric Space, Notes on Intuitionistic Fuzzy sets, **21**(1) (2015), pp. 43-53.
- [8] S. Melliani, M. Elomari, M. Atraoui and L. S. Chadli, Intuitionistic fuzzy differential equation with nonlocal condition, Notes on Intuitionistic Fuzzy Sets, **21**(4)(2015), pp. 58-68.
- [9] R. Ettoussi, S. Melliani and L.S. Chadli, Solution of Intuitionistic Fuzzy Differential Equations(submitted on Notes on Intuitionistic Fuzzy Sets)**21**(2)(2016).
- [10] Zadeh, L.A., Fuzzy sets, Inf. Control, **8**(1965), pp. 338-353.

## Fractional Differential Equations with Intuitionistic Fuzzy Data

**R. ETTOUSSI, L. S. CHADLI**  
razika.imi@gmail.com

### Abstract

The purpose of this paper is to study the existence and uniqueness of solution for fractional differential equation with intuitionistic fuzzy data where the intuitionistic fuzzy fractional derivatives and integral are considered in the Riemann-Liouville sense. Finally we give an example.

**Keywords :** intuitionistic fuzzy number, Fractional differential equations.

### Referances

- [1] K. Atanassov , Intuitionistic fuzzy sets, Fuzzy Sets and Systems 20(1986), pp. 87-96.
- [2] K. Atanassov , Intuitionistic Fuzzy Sets, Springer Physica-Verlag, Berlin, (1999).
- [3] I. M. Gelfand and G. E. Shilvov, Generalized Functions, 1(1958), Moscow.
- [4] A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, Theory and Applications of Fractional Differential Equations, Elsevier Science B.V, Amsterdam, 2006.
- [5] G. E. Shilove, Generalized functions and partial differential equations, in Mathematics and its Applications, Science publishers, Inc., 1968.

- [6] A. J. Turski, B. Atamaniuk and E. Turska, On the appearance of the fractional derivative in the behavior of real materials, *J. Appl. Mechanics*, 51 (1984), 294-298.
- [7] S. Melliani, M. Elomari, L. S. Chadli and R. Ettoussi, Intuitionistic Fuzzy metric spaces, *Notes on intuitionistic Fuzzy sets*, 21(1)(2015), pp. 43-53.
- [8] S. Melliani, M. Elomari, L. S. Chadli and R. Ettoussi, Intuitionistic fuzzy Fractional differential equations, *Notes on Intuitionistic Fuzzy Sets*, 21(4) (2015), pp. 76-89.
- [9] D. Junsheng, A. Jianye and Xu Mingyu; Solution of system of fractional differential equations by Adomian Decomposition Method, *Appl. Math. J. Chinese Univ. Ser. B* 2007, 22: 7-12.
- [10] V. Lakshmikantham and J. Vasundhara Devi; Theory of fractional differential equations in a Banach Space. *European Journal of Pure and Applied Mathematics*, 1(1)(2008), pp. 38-45.

## Solving Second Order Intuitionistic Fuzzy Initial Value Problems with Heaviside Function

Ömer AKIN, Selami BAYEG

TOBB Economy and Technology University, Department of Mathematics,  
Ankara

omerakin@etu.edu.tr, sbayeg@etu.edu.tr

### Abstract

In this work, we examined the solution of the following second order intuitionistic fuzzy initial value problem through intuitionistic Zadeh's Extension Principle [17]:

$$y''(x) + \bar{a}_1^i y'(x) + \bar{a}_2^i y(x) = \sum_{j=1}^r \bar{b}_j^i g_j(x); \quad (0.0.1)$$

$$y(0) = \bar{\gamma}_0^i; \quad (0.0.2)$$

$$y'(0) = \bar{\gamma}_1^i. \quad (0.0.3)$$

Here  $\bar{a}_1^i$ ,  $\bar{a}_2^i$ ,  $\bar{\gamma}_0^i$ ,  $\bar{\gamma}_1^i$  and  $\bar{b}_j^i$  ( $j=1,2,\dots,r$ ) are intuitionistic fuzzy numbers and  $g_i(x)$  ( $i=1,2,\dots,r$ ) are continuous functions on the interval  $[0, \infty)$ . We reformulated the approach in [2] and [3] for finding an analytical form of alpha and beta cuts for the solution of intuitionistic fuzzy initial value problem for the second order differential equation with the help of Heaviside step function. Firstly we reformulated the general solution of the crisp differential equation corresponding to Eq. 0.1 and applied intuitionistic Zadeh's Extension Principle to intuitionistically fuzzify the solution. Then, we obtained the analytical form

of  $(\alpha, \beta)$ -cuts of the solution of the fuzzy initial value problem by using interval operations and Heaviside step function. Finally, we have illustrated some examples by using this algorithm.

**Keywords :** Intuitionistic Fuzzy Initial Value Problem, Intuitionistic Zadeh's Extension Principle, Heaviside Function.

### Referances

- [1] L. Atanassova, *On Intuitionistic Fuzzy Versions of L. Zadeh's Extension Principle*, NIFS, vol. 13. pp. 33–36, 2007.
- [2] O.Akin, T.Khaniyev, S.Bayeg, B. Turken *Solving a Second Order Fuzzy Initial Value Problem using, Turk. J. Math. Comput. Sci.,4(2016) 16–25,2015.*
- [3] Buckley, J. J.,Feuring, T.,*Fuzzy initial value problem for N-th order linear differential equations*, Fuzzy Sets and Systems, 121, 247–255, 2001.
- [4] ...

## Equivalence Among Three 2-Norms on the Space of $p$ -Summable Sequences

**Sukran KONCA, Mochammad IDRIS**

Department of Mathematics, Bitlis Eren University, 13000, Bitlis, Turkey  
 Department of Mathematics, Institute of Technology Bandung 40132,  
 Bandung, Indonesia  
 skonca@beu.edu.tr, idemath@gmail.com

### Abstract

There are two known 2-norms defined on the space of  $p$ -summable sequences of real numbers. The first 2-norm is a special case of Gähler's formula [Mathematische Nachrichten, 1964], while the second is due to Gunawan [Bulletin of the Australian Mathematical Society, 2001]. The aim of this paper is to define a new 2-norm on  $\ell^p$  and prove the equivalence among these three 2-norms.

**Keywords :** 2-normed spaces; the space of  $p$ -summable sequences; completeness; norm equivalence.

### Referances

- [1] Gähler, S., *2-metrische räume und ihre topologische struktur*, Mathematische Nachrichten, Vol. **26**, 115-148, 1963.

- [2] Gozali, S.M., Gunawan H. and Neswan, O., *On  $n$ -norms and bounded  $n$ -linear functionals in a Hilbert space*, Annals of Functional Analysis, Vol. **1**, 72-79, 2010.
- [3] Gunawan, H., *The space of  $p$ -summable sequences and its natural  $n$ -norms*, Bulletin of the Australian Mathematical Society, Vol. **64**, 137-147, 2001.
- [4] Idris, M., Ekariani, S. and Gunawan, H., *On the space of  $p$ -summable sequences*, Matematicki Vesnik, Vol. **65** No. 1, 58-63, 2013.
- [5] Konca, S., Idris, M., Gunawan, H. and Basarir, M.  *$p$ -Summable sequence spaces with 2-inner products*, Contemporary Analysis and Applied Mathematics, Vol. **3** No. 2, 213-226, 2015.
- [6] Konca, S., Idris, M. and Gunawan, H. *A new 2-inner product on the space of  $p$ -summable sequences*, Journal of Egyptian Mathematical Society, Vol. **24** No. 2, 244-249, 2016.
- [7] Konca, S., Gunawan, H. and Başarır, M., *Some remarks on  $\ell^p$  as an  $n$ -normed space*, Mathematical Sciences and Applications E-Notes, Vol. **2** No. 2, 45-50, 2014.
- [8] Misiak, A.,  *$n$ -inner product spaces*, Mathematische Nachrichten, Vol. **140**, 299-319, 1989.
- [9] Mutaqin, A. and Gunawan, H., *Equivalence of  $n$ -norms on the space of  $p$ -summable sequences*, Journal of the Indonesian Mathematical Society, Vol. **16**, 2010.
- [10] Wibawa-Kusumah, R.A. and Gunawan, H. *Two equivalent  $n$ -norms on the space of  $p$ -summable sequences*, Periodica Mathematica Hungarica, Vol. **67** No. 1, 63-69, 2013.

## Some Properties of Soft Mappings on Soft Metric Spaces

**Murat Ibrahim YAZAR, Cigdem GUNDUZ (ARAS)**

Department of Primary Education, Karamanoğlu Mehmetbey University  
Yunus Emre Campus, 70100, Karaman, Turkey  
Kocaeli University, Department of Mathematics, Kocaeli, 41380-Turkey  
myazar@kmu.edu.tr, carasgunduz@gmail.com

### Abstract

In this study we define the soft topology generated by the soft metric and show that every soft metric space is a soft normal space. We also investigate some properties of soft continuous mappings on soft metric spaces and finally we give a few examples of soft contraction mapping on soft metric spaces.

**Keywords:** soft metric space, soft normal space, soft continuous mapping, soft contraction mapping.

### Referances

- [1] Yazar, M.I., Gunduz(Aras),C. and Bayramov, S., *Fixed point theorems of soft contractive mappings*, Filomat, Vol. **30**, No.2, 269-279,2016.
- [2] Bayramov, S. and Gunduz(Aras), C., *Soft locally compact and soft para-compact spaces*, Journal of Mathematics and System Science, ,Vol. **3**, 122-130, 2013.
- [3] Das, S. and Samanta, S. K., *Soft metric*, Ann. Fuzzy Math. Inform., ,Vol. **6**, No. 1, 77-94, 2013.
- [4] Gunduz (Aras), C. ,Sonmez, A. and Çakallı, H., *On Soft Mappings*, arXiv:1305.4545v1 [math.GM] 16 May 2013.
- [5] Long-Guang, H. and Xian,Z., *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl.,Vol. **332**, 1468–1476, 2007.
- [6] Majumdar, P. and Samanta, S. K. , *On soft mappings*, Comput. Math. Appl. ,Vol. **60**, 2666-2672, 2010.
- [7] Molodtsov D., *Soft set theory-first results*, Comput. Math. Appl., Vol. **37**,19-31, 1999.
- [8] Shabir, M. and Naz, M., *On soft topological spaces*, Comput. Math. Appl., Vol. **61**, 1786-1799, 2011.
- [9] Rhoades, B.E., *A comparison of various definition of contractive mappings*, Trans. Amer. Math. Soc., Vol. **226**, 257-290, 1977.

## Soft Totally Bounded Spaces in Soft Metric Spaces

**Cigdem GUNDUZ (ARAS), Murat Ibrahim YAZAR**  
 Kocaeli University , Department of Mathematics, Kocaeli, 41380-Turkey  
 Department of Primary Education, Karamanoğlu Mehmetbey University  
 Yunus Emre Campus, 70100, Karaman, Turkey  
 carasgunduz@gmail.com, myazar@kmu.edu.tr

### Abstract

In this study we define the soft topology generated by the soft metric and show that every soft metric space is a soft normal space. We also investigate some properties of soft continuous mappings on soft metric spaces and finally we give a few examples of soft contraction mapping on soft metric spaces.

**Keywords:** Soft set, soft metric space, soft sequential compact, soft totally bounded sets.

### Referances



- [1] Das, S. and Samanta, S. K., *Soft metric*, Ann. Fuzzy Math. Inform. ,Vol. **6**, No. 1, 77-94, 2013.
- [2] Molodtsov D., *Soft set theory-first results*, Comput. Math. Appl., Vol. **37**,19-31, 1999.
- [3] Shabir, M. and Naz, M., *On soft topological spaces*, Comput. Math. Appl., Vol. **61**, 1786-1799, 2011.
- [4] Bayramov, S. and Gunduz(Aras), C., *Soft locally compact and soft para-compact spaces*, Journal of Mathematics and System Science, ,Vol. **3**, 122-130, 2013.

## Some Generalized Fixed Point Type Theorems on an $S$ -Metric Space

**Nihal TAŞ, Nihal YILMAZ ÖZGÜR**

Balıkesir University, Department of Mathematics, 10145 Balıkesir, Turkey  
nihaltas@balikesir.edu.tr, nihal@balikesir.edu.tr

### Abstract

In this talk, we give new contractive mappings on an  $S$ -metric space. We investigate some generalizations of the Banach's contraction principle and new fixed point type theorems using the notion of periodic index on an  $S$ -metric space.

**Keywords:**  $S$ -metric, Banach's contraction principle, periodic index.

### Referances

- [1] Chang, S.S. and Zhong, Q.C., *On Rhoades' open questions*, Proc. Amer. Math. Soc., Vol. **109**, No. 1, 269-274, 1990.
- [2] Gupta, A., *Cyclic contraction on  $S$ -metric space*, International Journal of Analysis and Applications, Vol. **3**, No. 2, 119-130, 2013.
- [3] Hieu,N.T., Ly, N.T. and Dung, N.V., *A generalization of Ciric quasi-contractions for maps on  $S$ -metric spaces*, Thai Journal of Mathematics, Vol. **13**, No. 2, 369-380, 2015.
- [4] Özgür, N.Y. and Taş, N., *Some generalizations of fixed-point theorems on  $S$ -metric spaces*, Essays in Mathematics and Its Applications in Honor of Vladimir Arnold, New York, Springer, 2016.
- [5] Özgür, N.Y. and Taş, N., *Some fixed point theorems on  $S$ -metric spaces*, submitted for publication.
- [6] Özgür, N.Y. and Taş, N., *Some new contractive mappings on  $S$ -metric spaces and their relationships with the mapping (S25)*, submitted for publication.

- [7] Rhoades, B.E., *A comparison of various definitions of contractive mappings*, Trans. Amer. Math. Soc., Vol. **226**, 257-290, 1977.
- [8] Sedghi, S., Shobe, N. and Aliouche, A., *A generalization of fixed point theorems in  $S$ -metric spaces*, Mat. Vesnik, Vol. **64**, No. 3, 258-266, 2012.
- [9] Sedghi, S. and Dung, N.V., *Fixed point theorems on  $S$ -metric spaces*, Mat. Vesnik, Vol. **66**, No. 1, 113-124, 2014.

## A New Generalization of Soft Metric Spaces

Nihal TAŞ, Nihal YILMAZ ÖZGÜR

Balıkesir University, Department of Mathematics, 10145 Balıkesir, Turkey  
 nihaltas@balikesir.edu.tr, nihal@balikesir.edu.tr

### Abstract

In this talk, we describe the notion of a soft  $S$ -metric as a generalization of a soft metric. We investigate some basic and topological properties of this new metric. Also we give some existence and uniqueness conditions of fixed-point theorems on a complete soft  $S$ -metric space. We verify our results with some examples.

**Keywords:** Soft  $S$ -metric space, fixed point, topological properties

### References

- [1] Das, S. and Samanta, S. K., *Soft real sets, soft real numbers and their properties*, J. Fuzzy Math., Vol. **20**, No. 3, 551-576, 2012.
- [2] Das, S. and Samanta, S. K., *Soft metric*, Ann. Fuzzy Math. Inform., Vol. **6**, No. 1, 77-94, 2013.
- [3] Maji, P. K., Biswas, P. and Roy, A. R., *Soft set theory*, Comput. Math. Appl., Vol. **45**, 555-562, 2003.
- [4] Molodtsov, D., *Soft set theory - first results*, Comput. Math. Appl., Vol. **37**, 19-31, 1999.
- [5] Özgür, N.Y. and Taş, N., *A note on "application of fuzzy soft sets to investment decision making problem"*, Journal of New Theory, **1**, No. 7, 1-10, 2015.
- [6] Özgür, N.Y. and Taş, N., *Some generalizations of fixed-point theorems on  $S$ -metric spaces*, Essays in Mathematics and Its Applications in Honor of Vladimir Arnold, New York, Springer, 2016.
- [7] Özgür, N.Y. and Taş, N., *Some generalizations of the Banach's contraction principle on complete soft  $S$ -metric spaces*, submitted for publication.

- [8] Sedghi, S., Shobe, N. and Aliouche, A., *A generalization of fixed point theorems in S-metric spaces*, Mat. Vesnik, Vol. **64**, No. 3, 258-266, 2012.
- [9] Sedghi, S. and Dung, N.V., *Fixed point theorems on S-metric spaces*, Mat. Vesnik, Vol. **66**, No. 1, 113-124, 2014.

## Some Separation Axioms in Fuzzy Soft Topological Spaces

**Vildan ÇETKİN, Halis AYGÜN**

Kocaeli University, Department of Mathematics, Umuttepe Campus,  
41380, Kocaeli-TURKEY  
vcetkin@gmail.com, halis@kocaeli.edu.tr

### Abstract

Molodtsov (1999) proposed a completely new concept called soft set theory to model uncertainty, which associates a set with a set of parameters. Pei and Miao (2005) showed that soft sets are a class of special information systems. Later, Maji et al. (2001) introduced the concept of a fuzzy soft set which combines a fuzzy set and a soft set. From then on, many authors have contributed to (fuzzy) soft set theory in the different fields such as algebra, topology and etc. Soft topology is a relatively new and promising domain which can lead to the development of new mathematical models and innovative approaches that will significantly contribute to the solution of complex problems in natural sciences. Separation is an essential part of topology, on which a lot of work has been done. The aim of this work is to generalize some low-level separation axioms in fuzzifying topology and fuzzy topology to the fuzzifying soft topology and fuzzy soft topology by considering parametrization. So, we obtain some fundamental properties and characterizations of proposed separations.

**Keywords:** Fuzzy soft set, fuzzy soft topology, separation axiom

### Referances

- [1] Ahmad, B. and Kharal, A., *On fuzzy soft sets*, Advances in Fuzzy Systems, Article ID 586507, 2009.
- [2] Aygünoğlu, A., Çetkin, V., and Aygün, H., *An introduction to fuzzy soft topological spaces*, Hacettepe Journal of Mathematics and Statistics, **48**, No. 2, 197-208, 2014.
- [3] Çetkin, V., and Aygün, H., *On convergence of fuzzy soft filters*, 3rd International Eurasian Conference on Mathematical Sciences and Applications, Vienna, 25-28 August 2014.

- [4] Çetkin, V., and Aygün, H., *Lowen functors in fuzzy soft topological spaces*, the 4th International Fuzzy Systems Symposium, Istanbul, 5-6 November 2016.
- [5] Hong-Yan, L., Fu-Gui, S., *Some separation axioms in I-fuzzy topological spaces*, Fuzzy Sets and Systems, 159, 573-587, 2008.
- [6] Kharal,, A., Ahmad, B., *Mappings on fuzzy soft classes*, Advances in Fuzzy Systems, Article ID 407890, 2009.
- [7] Maji, P. K., Biswas, R., Roy, A. R., *Fuzzy soft sets*, Journal of Fuzzy Mathematics, **9**, No. 3, 589-602, 2001
- [8] Molodtsov, D., *Soft set theory: First results*, Computers and Mathematics with Applications, **37**, No. 4/5, 19-31, 1999.
- [9] Pei, D., Miao, D., *From soft sets to informations systems*, Granular Computing, 2005 IEEE International Conference on (2), 2005, pp. 617-621.
- [10] Yueli, Y., Jinming, F., *On separation axioms in I-fuzzy topological spaces*, Fuzzy Sets and Systems, 157, 780-793,2006.

## Fuzzy Equilibrium Analysis of a Transportation Network Problem

**Gizem TEMELCAN\***, Hale Gonce KOCKEN,  
İnci ALBAYRAK

\*PhD Student at Mathematical Engineering Program, Yildiz Technical University, Istanbul, TURKEY

Mathematical Engineering Department, Yildiz Technical University, Istanbul, TURKEY

temelcan.gizem@gmail.com, halegk@gmail.com, ibayrak@yildiz.edu.tr

### Abstract

In this paper, we focused on the solution process of a fuzzy transportation network equilibrium problem. This problem aims to minimize the total travel time of vehicles on traffic flows between specified origin and destination points. The link travel time for a vehicle is taken as a linear function of link flow (the number of vehicles on that link). Thus, the objective function can be formulated in terms of link flows and link travel times in a quadratic form while satisfying the flow conservation constraints. The parameters of this problem are path lengths, number of lanes, average velocity of a vehicle, vehicle-length, clearance, spacing, link capacity and free flow travel time. Considering a road network, path lengths and number of lanes are taken as crisp numbers. The average velocity of a vehicle and the vehicle-length are imprecise in nature, so these are taken as triangular fuzzy numbers. Since the remaining parameters, that are clearance, spacing, link capacity and free flow travel time, are determined by the average velocity of a vehicle and vehicle-length, all of them will be triangular

fuzzy numbers. Finally, the original fuzzy transportation network problem is converted to a fuzzy quadratic programming problem, and it is solved with an existing approach from the literature. A numerical experiment is illustrated.

**Keywords:** Fuzzy transportation network equilibrium problem, fuzzy quadratic programming, triangular fuzzy numbers.

### Referances

- [1] Leblanc, L.J, Morlok, E.K and Pierskalla, W.P., *An Efficient Approach to Solving the Road Network Equilibrium Traffic Assignment Problem*, Transportation Research, Vol. **9**, 309-318, 1975.
- [2] Leblanc, L.J., *The Use of Large Scale Mathematical Programming Models in Transportation Systems*, Transportation Research, Vol. **10**, 419-421, 1976.
- [3] Branston, D., *Link Capacity Functions: A Review*, Transportation Research, Vol. **10**, 233-236, 1976.
- [4] Daganzo, C.F., *On the Traffic Assignment Problem with Flow Dependent Costs-I*, Transportation Research, Vol. **11**, 433-437, 1977.
- [5] Sheffi, Y., *Urban Transportation Networks: Equilibrium Analysis with Mathematical Programming Methods*, Prentice-Hall, USA, 1985.

## A Special Type of Sasakian Finsler Structures on Vector Bundles

Nesrin CALIŞKAN, Ayşe Funda YALINIZ

Faculty of Education, Department of Mathematics Education, Uşak  
University, 64200, Uşak, Turkey

Faculty of Arts and Sciences, Department of Mathematics, Dumlupınar  
University, 43100, Kütahya, Turkey

nesrin.caliskan@usak.edu.tr, caliskan.nesrin35@gmail.com,  
fundayaliniz@gmail.com

### Abstract

Sasakian Finsler structures can be obtained on horizontal and vertical distributions of vector bundles. In this paper, Sasakian Finsler structures satisfying  $R(X^H, Y^H)C^* = 0$  on horizontal distribution of vector bundles are examined where  $R$  is Riemann curvature tensor,  $C^*$  is quasi-conformal curvature tensor and  $X^H, Y^H$  are elements of family of vector fields on horizontal distribution. In this regard some structure theorems are examined.

**Keywords:** Quasi-conformal curvature tensor, Sasakian Finsler structure.

### References

- [1] Antonelli P.L., *Handbook of Finsler Geometry*, Kluver Academic Publishers, ISBN-10:1402015550, 1437 p., 2003.
- [2] Besse A.L., *Einstein manifolds*, Series: Classics in Mathematics, ISBN 978-3-540-74120-6, doi 10.1007/978-3-540-74311-8, 516 p. 2002.
- [3] De U.C., Jun J.B. and Gaz A.K., *Sasakian manifolds with quasi-conformal curvature tensor*, Bull. Korean Math. Soc., Vol. **45**, No. 2, 313-319, 2008.
- [4] Tripathi M.M., Gupta P. and Kim J.S., *Index of quasi-conformally symmetric semi-Riemannian manifolds*, Int. J. Math. Math. Sci., Vol. **2012**, Article ID 461383, 14 p., 2012.
- [5] Yano K. and Kon M., *Structures on manifolds*, Series in Pure Mathematics, Vol. **3**, World Scientific-Singapore, 508 p., 1984.
- [6] Yalnız A.F. and Çalışkan N., *Sasakian Finsler manifolds*, Turk. J. Math., Vol. **37**, No. 2, 319-339, 2013.

## On Some Properties and Applications of the Quasi-Resolvent Operators of the Infinitesimal Operator of a Strongly Continuous Linear Representation of the Unit Circle Group in a Complex Banach Space

**Abdullah ÇAVUŞ**

Emeritus Professor, Worked for Karadeniz Technical University, Faculty  
of Science, Department of Mathematics, Trabzon, Turkey  
abcavus@gmail.com

### Abstract

Let  $A$  be the infinitesimal operator of a strongly continuous linear representation of the unit circle group on a complex Banach space  $H$ . In this talk, the quasi-resolvent operator of  $A$  which is denoted by  $R_\lambda$  is defined by the spectrum of  $A$ . Some properties and inter relations of operators  $R_\lambda$  are introduced, and by using them, some theorems on existence of periodic solutions to the non-linear equations  $\phi(A)x = f(x)$  are stated and proven, where  $\phi(A)$  is a polynomial of  $A$  and  $f$  is a continuous mapping of  $H$  into itself.

## Existence and Nonexistence for Nonlinear Problems with Singular Potential

**B. ABDELLAOUI\***, **K. BIROUD\***, **J. DAVILA\*\***, **F. MAHMOUDI\*\***

\*Département de Mathématiques, Université Abou Bakr Belkaïd  
Tlemcen 13000, Algeria.

\*\*Departamento de Ingenieria Matematica, CMM Universidad de Chile,  
Casilla 170-3 Correo 3, Santiago, Chile  
boumediene.abdellaoui@uam.es, kh\_biroud@yahoo.fr,  
jdavila@dim.uchile.cl, fmahmoudi@dim.uchile.cl

### Abstract

Let  $\Omega \subset \mathbb{R}^N$  be a bounded regular domain of  $\mathbb{R}^N$  we consider the following class of elliptic problem

$$\begin{cases} -\Delta u = \frac{u^q}{d^2} & \text{in } \Omega, \\ u > 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where  $0 < q \leq 2^* - 1$ . We investigate the question of existence and nonexistence of positive solutions depending on the range of the exponent  $q$ .

## Intuitionistic Fuzzy Soft Generalized Superconnected

**I. BAKHADACH, S. MELLIANI, L. S. CHADLI**  
LMACS, Laboratory of Applied Mathematics & Computing sciences  
Sultan Moulay Slimane University, PO Box 523, 23000 Beni Mellal  
Morocco  
said.melliani@gmail.com

### Abstract

Shabir and Naz [7] introduced the notion of soft topological spaces which are defined over an initial universe with a fixed set of parameters. They also studied some of basic concepts of soft topological spaces. In the present study, we introduce some new concepts in intuitionistic fuzzy soft topological spaces such

as intuitionistic fuzzy soft generalized superconnected. We also give characterizations and properties of this notion.

**Keywords:** intuitionistic fuzzy soft set, Intuitionistic fuzzy soft topology, intuitionistic fuzzy soft mapping.

### References

- [1] K. Atanassov, On Intuitionistic Fuzzy Sets Theory, Springer, (2012) Berlin
- [2] A. Csázar, Generalized open sets in generalized topologies , Acta math. Hungar., 106 (2005) 53–66.
- [3] A. Kharal and B. Ahmad, Mappings on soft classes, to appear in New Math. Nat. Comput.
- [4] K. Kannan, Soft Generalized Closed Sets In Soft Topological Spaces, J. Theoret. Appl. Inf. Tech., 37 (2012) 17–21.
- [5] P. K. Maji, R. Biswas and A. R. Roy, Intuitionistic fuzzy soft sets, J. Fuzzy Mathematics, 9 (3) (2001) 677–693.
- [6] D. Molodtsov, Soft set theory-first results, Computers and Mathematics with Applications, 37 (1999) 19–31.
- [7] M. Shabir and M. Naz , On soft topological spaces , Comput. Math. Appl., 61 (2011) 1786-1799.
- [8] I. Zorlutuna, M. Akdag, W. K. Min and S. Atmaca, Remarks on soft topological spaces, Ann. Fuzzy Math. Inform., 3 (2) (2012) 171–185.
- [9] I. Zorlutuna, M. Akdag, W. K. Min and S. Atmaca, Remarks on soft topological spaces, Ann. Fuzzy Math. Inform., 3 (2) (2012) 171–185.

## Vietoris Topology in the Context of Soft Set

İzzettin DEMİR

Department of Mathematics, Duzce University, 81620, Duzce-Turkey  
 izzettindemir@duzce.edu.tr

### Abstract

In the present paper, we study the notion of a Vietoris topology by using soft sets. We obtain some properties related to the first countability of soft Vietoris topology. Then, we focus on second countability of it.

**Keywords:** Soft set, Soft Vietoris topology, Soft first countability, Soft second countability.

### References



- [1] Akdağ, M. and Erol, F., *On hyperspaces of soft sets*, Journal of New Theory, Vol. **7**, 86-97, 2015.
- [2] Andrijevic, D., Jelic, M. and Mrsevic, M., *Some properties of hyperspaces of Cech closure spaces*, Filomat, Vol. **24**, 53-61, 2010.
- [3] Aygünöglu, A. and Aygün, H., *Some notes on soft topological spaces*, Neural Comput. Appl., Vol. **21**, 113–119, 2012.
- [4] Demir, İ. and Özbakır, O.B., *Soft Hausdorff spaces and their some properties*, Annals of Fuzzy Mathematics and Informatics, Vol. **8**, No. 5, 769-783, 2014.
- [5] Di Maio, G. and Hola, L., *On hit-and-miss hyperspace topologies*, Rend. Accad. Sci. Fis. Mat. Napoli, Vol. **62**, 103-124, 1995.
- [6] Hola, L. and Levi, S., *Decomposition properties of hyperspace topologies*, Set-valued Analysis, Vol. **5**, 309-321, 1997.
- [7] Hussain, S. and Ahmad, B., *Soft separation axioms in soft topological spaces*, Hacettepe Journal of Mathematics and Statistics, Vol. **44**, 559-568, 2015.
- [8] Nazmul, S. and Samanta, S.K., *Neighbourhood properties of soft topological spaces*, Ann. Fuzzy Math. Inform., Vol. **6**, 1–15, 2013.
- [9] Shabir, M. and Naz, M., *On soft topological spaces*, Comput. Math. Appl., Vol. **61**, 1786–1799, 2011.
- [10] Shakir, Q. R., *On Vietoris soft topology I*, J. Sci. Res., Vol. **8**, 13-19, 2016.
- [11] Smithson, R. E., *First countable hyperspaces*, Proc. Amer. Math. Soc., Vol. **36**, 325-328, 1976.
- [12] Zsilinszky, L., *Baire spaces and hyperspace topologies*, Proc. Amer. Math. Soc., Vol. **124**, 2575-2584, 1996.

## On Totally Umbilical and Minimal Cauchy Riemannian Lightlike Submanifolds of an Indefinite Kaehler Manifold

Burçin DOĞAN, Ali YAKAR, Erol YAŞAR  
Mersin University

bdogan@mersin.edu.tr, yakar1986@gmail.com, yerol@mersin.edu.tr

### Abstract

In this talk, we survey Cauchy Riemannian lightlike submanifolds of an indefinite Kaehler manifold. Firstly, we mention definition of Cauchy Riemannian (CR, SCR and GCR) lightlike submanifolds of an indefinite Kaehler manifold. Then, we investigate minimal and totally umbilical Cauchy Riemannian lightlike submanifolds and give some examples for these classes.

### Referances

- [1] Bejancu A., *Geometry of CR-Submanifolds*, Kluwer Academic, (1986).
- [2] Chen B.Y., *CR-Submanifolds of a Kähler Manifold, I-II*, J. Differential Geometry, 16 (1981), 305-322, 493-509.
- [3] Duggal K.L. and Bejancu A., *Lightlike submanifolds of semi-Riemannian Manifolds and Applications*, Kluwer Academic, (1996).
- [4] Kupeli D.N., *Singular Semi-Riemannian Geometry*, Kluwer Acad. Publishers, Dordrecht, (1996).
- [5] Duggal K.L. and Sahin B., *Screen Cauchy Riemann Lightlike Submanifolds*, Acta Math. Hungar., 106(1-2), (January 2005).
- [6] Duggal K.L. and Sahin B., *Generalized Cauchy Riemann Lightlike Submanifolds*, Acta Math. Hungar., 112(1-2), (2006).
- [7] Duggal K.L. and Sahin B., *Differential Geometry of Lightlike Submanifolds*, Birkhauser, (2010).
- [8] Sachdeva R., Kumar R. and Bhatia S.S., *Totally umbilical hemi-slant lightlike submanifolds*, New York J. Math., 21 (2015), 191-203.
- [9] Sahin B., *CR-lightlike Altmanifoldlar  $\mathbb{A}^{\pm n}$  Geometrisi* *Äæzerine*, Doktora Tezi, İnönü University Fen Bil. Ens., (2000).
- [10] Yano K. and Kon M., *CR-submanifolds of Kählerian and Sasakian Manifolds*, Birkhauser, (1983).

## Algebraic Properties of Dual Quasi-Quaternions

**Murat BEKAR, Yusuf YAYLI**

Necmettin Erbakan University, Department of Mathematics-Computer  
Sciences, 42090, Konya, Turkey  
Ankara University, Department of Mathematics, 06100, Ankara, Turkey  
mbekar@konya.edu.tr, yayli@science.ankara.edu.tr

### Abstract

In this work, we consider the algebra of dual quasi-quaternion and give some algebraic properties of this algebra.

### References

- [1] Ercan Z., Yuce S., *On properties of the Dual Quaternions*, European J. of Pure and Appl. Math., Vol. 4, No. 2, 142-146, 2011.
- [2] Majernik V., *Quaternion Formulation of the Galilean Space-Time Transformation*, Acta Phy. Slovaca, Vol. 56, No. 1, 9-14, 2006.
- [3] Rosenfeld B.A., *Geometry of Lie Groups*, Kluwer Academic Publishers, Dordrecht, 1997.

[4] Jafari M., *On the Properties of Quasi-quaternion Algebra*, Commun. Fac. Sci. Univ. Ank. Series A1 Vol. **63**, No. 1, 1-10, 2014.

## Fixed Intuitionistic Fuzzy Point Theorem in Hausdorff Intuitionistic Fuzzy Metric Spaces

Ferhan SOLA ERDURAN

Department of Mathematics Faculty of Science Gazi University 06500  
Teknikokullar, Ankara, Turkey  
ferhansola@yahoo.com

### Abstract

Atanassov (see [2,3]) introduced and studied the concept of intuitionistic fuzzy sets (i-fuzzy set, for short) and later there has been much progress in the study of i-fuzzy sets by many authors (see [1,4,5,10]). Using the idea of i-fuzzy set, Park [10] defined the notion of intuitionistic fuzzy metric space with the help of continuous t-norms and continuous t-conorms as a generalization of fuzzy metric space due to George and Veeramani [7] and proved some known results of metric spaces for intuitionistic fuzzy metric space. In 2001 Estruch and Vidal [6] introduced the concept of intuitionistic fuzzy mapping (i-fuzzy mapping, for short) and gave an intuitionistic version of Heilpern's mentioned theorem (see [9]). After that, Gregori et al [8] defined Hausdorff intuitionistic fuzzy metric on a family of non-empty compact subsets of a given intuitionistic fuzzy metric space.

In this study we modify concept of Hausdorff intuitionistic fuzzy metric using i-fuzzy sets and obtain fixed i-fuzzy point results for i-fuzzy mappings.

**Keywords:** i-fuzzy mapping, i-fuzzy point.

### Referances

- [1] Alaca, C., Turkoglu, D. and Yildiz, C., *Fixed points in intuitionistic fuzzy metric spaces*, Chaos, Solitons & Fractals, **29(5)**, 1073-1078, 2006.
- [2] Atanassov, K., *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, **20**, 87-96, 1986.
- [3] Atanassov, K., *Review and new results on intuitionistic fuzzy sets*, Preprint IM-MFAIS-1-88, 1988.
- [4] Çoker, D., *An introduction to Intuitionistic fuzzy topological spaces*, Fuzzy Sets and Systems, **88**, 81-89, 1997.
- [5] Efe, H. and Yildiz, C., *On The Hausdorff Intuitionistic Fuzzy Metric Spaces*, International Journal of Pure and Applied Mathematics, **31(2)**, 143-165, 2006.

- [6] Estruch, V.D. and Veeramani, P., *A note on fixed fuzzy points for fuzzy mappings*, Rend. Istit. Mat. Univ. Trieste, **32**, 39-45, 2001.
- [7] George, A. and Vidal, A., *On some results in fuzzy metric spaces*, Fuzzy Sets and Systems, **64**, 395-399, 1994.
- [8] Gregori, V., Romaguera, S. and Veeramani, P., *A note on intuitionistic fuzzy metric spaces*, Chaos, Solitons & Fractals, **28**, 902-905, 2006.
- [9] Heilpern, S., *Fuzzy mappings and fixed point theorem*, J. Math. Anal. Appl., **83(2)**, 566-569, 1981.
- [10] Park, J.H., *Intuitionistic fuzzy metric spaces*, Chaos, Solitons & Fractals, **22**, 1039-1046, 2004.

## Orthonormal Systems in Spaces of Number Theoretical Functions

**Erdener Kaya**  
Mersin University  
kayaerdener@mersin.edu.tr

### Abstract

In this paper we regard some (for number theory important) examples of set algebras  $\mathcal{A}$  on  $\mathbb{N}$ . In each example we obtain the measure space  $\Omega := (\beta\mathbb{N}, \sigma(\overline{\mathcal{A}}), \bar{\delta})$  by the model of Indlekofer which is based on the Stone-Cech compactification of  $\mathbb{N}$ .

Let  $\mathcal{E}(\mathcal{A})$  be the set of simple functions on  $\mathcal{A}$  and let  $\mathcal{L}^{*\alpha}(\mathcal{A})$  be  $\|\cdot\|_\alpha$  the closure of  $\mathcal{E}(\mathcal{A})$  with

$$\|f\|_\alpha := \left\{ \limsup_{x \rightarrow \infty} \frac{1}{x} \sum_{n \leq x} |f(n)|^\alpha \right\}^{\frac{1}{\alpha}}, 1 \leq \alpha < \infty.$$

Now our aim was to give a description of a complete orthonormal system for  $L^{*2}(\mathcal{A})$  in each regarded case where  $L^{*2}(\mathcal{A})$  is denoted the quotient space  $\mathcal{L}^{*2}(\mathcal{A})$  modulo null-functions.

**Keywords:** Stone-Cech compactification, function spaces, complete orthonormal systems.

### References

- [1] Indlekofer, K.-H., *A new method in probabilistic number theory*, Probability Theory and Applications, Math. Appl., **80**, 299-308, 1992.

- [2] Indlekofer, K.-H., *New approach to probabilistic number theory-compactifications and integration*, *Advanced Studies in Pure Mathematics*, **49**, 133-170, 2005.
- [3] Indlekofer, K.-H., *Some remarks on almost-even and almost-periodic functions*, *Arch. Math.*, **37**, 353-358, 1981.
- [4] Indlekofer, K.-H., Y.-W. Lee and R. Wagner, *Mean behaviour of uniformly summable  $q$ -multiplicative functions*, *Ann. Univ. Sci. Budapest.*, **25**, 171-294, 2005.
- [5] Schwarz, W. and Spilker J., *Arithmetical Functions: An Introduction to Elementary and Analytic Properties of Arithmetic Functions and to Some of Their Almost-Periodic Properties*, Cambridge University Press, Cambridge, 388p. 1994.

## A Step Size Strategy for Numerical Integration of the Hurwitz Stable Differential Equation Systems

**Gülnur ÇELİK KIZILKAN, Kemal AYDIN**

Necmettin Erbakan University, Faculty of Science, Department of  
Mathematics and Computer Science, Konya, Turkey

Selçuk University, Faculty of Science, Department of Mathematics,  
Konya, Turkey

gckizilkan@konya.edu.tr, kaydin@selcuk.edu.tr

### Abstract

For the efficiency of the numerical integration of the Cauchy problems, it is not practical to use constant step size. There are some studies in the literature about the variable step size for numerical integration (for example see; [3,4,5]). One of these studies has given in [1,2]. On the region  $D = (t, X) : |t - t_0| \leq T, |x_j - x_{j0}| \leq b_j$ , in [1,2] the step size strategy for the Cauchy Problem

$$X' = AX, X(t_0) = X_0$$

has proposed such that the local error  $\|LE_i\| \leq \delta_L$ , where  $\delta_L$  is the error level that is determined by user. Here  $X(t) = (x_j(t))$ ,  $X_0 = (x_{j0})$ ;  $x_{j0} = x_j(t_0)$ ,  $A \in R^{N \times N}$ ,  $X(t)$ ,  $X_0$  and  $b = (b_j) \in R^N$ .

In this study, we aimed to develop the step size strategy in [1,2] for the system

$$X'^{N \times N} - \text{Hurwitz stable matrix.}$$

A step size strategy and an algorithm for the Hurwitz systems which calculate the step sizes based on the given strategy and numerical solutions are given. The numerical solutions obtained with the new strategy and algorithm are compared with the results in [1,2]. The given strategy and algorithm are applied to some industrial problems.

**Keywords:** Variable step size, Hurwitz stable differential equation systems, numerical integration, step size strategy

### References

- [1] Çelik Kızıllkan, G., *Step size strategies on the numerical integration of the systems of differential equations*, Ph.D. Thesis, Selcuk University Graduate Natural and Applied Sciences, Konya (in Turkish), 2009.
- [2] Çelik Kızıllkan, G., Aydın, K., *Step size strategies based on error analysis for the linear systems*, SDU Journal of Science, Vol. **6**, No. 2, 149-159, 2011.
- [3] Holsapple, R., Iyer, R., Doman, D., *Variable Step-Size Selection Methods for Implicit Integration Schemes for ODEs*, IJNAM, Vol. **4**, No. 2, 212-242, 2007.
- [4] Ramos, H., Vigo-Aguiar, J., *Variable Stepsize Störmer-Cowell Methods*, Math Comput Model, Vol. **42**, 837-846, 2005.
- [5] Shampine, L.F., Allen, R. C., Pruess, S., *Fundamentals of Numerical Computing*, John Wiley Sons, INC, New York, 1996.

## Schur Stabilitiy in Floating Point Arithmetic: Systems with Constant Coefficients

Ahmet Duman\*, Ali Osman ÇIBIKDİKEN,  
Gülnur ÇELİK KIZILKAN\*, Kemal AYDIN

\*Necmettin Erbakan University, Faculty of Science, Department of  
Mathematics and Computer Science, Konya, Turkey  
Necmettin Erbakan University, Faculty of Engineering and Architecture,  
Department of Computer Engineering, Konya, Turkey  
Selçuk University, Faculty of Science, Department of Mathematics,  
Konya, Turkey  
aduman@konya.edu.tr, aocdiken@konya.edu.tr, gckizilkan@konya.edu.tr,  
kaydin@selcuk.edu.tr

### Abstract

The representation of the numbers  $\mathbb{F} = \mathbb{F}(\gamma, p_-, p_+, k) = \{0\} \cup \{z | z = \pm \gamma^{p(z)} m_\gamma(z)\}$ , which are called floating point numbers [1,9,13,14]. Computers use floating point numbers for computing. These numbers are also called computer numbers or machine numbers [3,10,15,17]. If the results of arithmetic operations are elements of set  $\mathbb{F}$ , they are directly stored in the memory. Otherwise they are stored with error [4,7,11,12,16,18].

The linear difference equation in order of  $N$  as  

$$y(n + N) = a_N y(n + N - 1) + \dots + a_1 y(n)$$

can be transform to one order system as

$$x(n+1) = Ax(n), \quad n\text{-integer number,}$$

where matrix  $A$  ( $N \times N$  dimension) is the companion matrix. It is well known that the solution of the Cauchy problem

$$x(n+1) = Ax(n), \quad x(0) = x_0$$

is  $x(n) = A^n x_0$  (see, [1,8]). In [2], an algorithm has given which have computed power of the companion matrix.

The matrix  $fl(A^n)$  is the computed companion matrix  $A^n$  in floating point arithmetic. The effects of floating point arithmetic in the computation of the companion matrix  $A^n$  were investigated [5,6]. Error bounds were obtained for  $\|A^n - fl(A^n)\|$ , where  $A \in M_N(D)$ . Additionally, Schur stability were investigated according to floating point arithmetic. The obtained results were supported with numerical examples.

**Keywords:** floating point arithmetics, difference equation, error analysis, companion matrix.

## Referances

- [1] Akın, O. and Bulgak H., *Linear difference equations and stability theory*, Selçuk University Research Centre of Applied Mathematics, No. 2, Konya (in Turkish), 1998.
- [2] Aydın, K., *On a power of the companion matrix*, SDU Journal of Science (E-Journal), Vol. 3, No. 2. 230-235, 2008.
- [3] Behrooz, P., *Computer Architecture: From Microprocessors to Supercomputers*, Oxford University Press, 2005.
- [4] Çibikdiken, A.O., *Error estimation of elementary matrix operations*, Master Thesis, Selçuk University Graduate Natural and Applied Sciences, Konya (in Turkish), 2002.
- [5] Çibikdiken, A.O., *The effect of floating point arithmetic on fundamental matrix computation of periodic linear difference equation system*, Ph.D. Thesis, Selçuk University Graduate Natural and Applied Sciences, Konya (in Turkish), 2008.
- [6] Çibikdiken, A.O., Aydın, K., *Computation of monodromy matrix on floating point arithmetic with Godunov Model*, Konuralp Journal of Mathematics, Volume 4, No 1, 2016.
- [7] Dahlquist, G., Björck, A., *Numerical Mathematics and Scientific Computation*, SIAM, Vol. 1, 2008.
- [8] Elaydi, S.N., *An Introduction to Difference Equations*, Second Edition, Springer-Verlag, New York, 1999.
- [9] Godunov, S.K., Antonov, A.G., Kiriluk, O.P., Kostin, V.I., *Guaranteed Accuracy in Mathematical Computations*, Prentice-Hall, Englewood Cliffs, NJ, 1993.
- [10] Goldberg, D., *What every computer scientist should know about floating-point arithmetic*, J. ACM Comput. Surv., Vol. 23, No. 1, 5-48, 1991.
- [11] G.H. Golub, C.F. Van Loan, *Matrix Computations*, (third ed.)The Johns Hopkins University Press, Baltimore (1996).

- [12] Higham, N.J., *Accuracy and Stability of Numerical Algorithms*, SIAM, 1996.
- [13] Kulisch, U.W., *Mathematical foundation of computer arithmetic*, IEEE Trans. Comput., C-26 (7), 610-621, 1977.
- [14] Kulisch, U.W., Miranker, W.L., *Computer Arithmetic in Theory and Practice*, Academic Press Inc., 1981.
- [15] Overton, M.L., *Numerical Computing with IEEE Floating Point Arithmetic*, SIAM, 2001.
- [16] Shampine, L.F., Allen, R.C., Pruess, S., *Fundamentals of Numerical Computing*, John Wiley - Sons Inc., 1997.
- [17] Sterbenz, P.H., *Floating-Point Computation*, Prentice-Hall Inc., Englewood Cliffs, NJ, 1974.
- [18] Wilkinson, J.H., *Rounding Errors in Algebraic Processes*, Prentice-Hall Inc., 1963.

## The Numerical Solution of Some SIR Epidemic Models with Variable Step Size Strategy

**Gülnur ÇELİK KIZILKAN, Ahmet DUMAN,  
Kemal AYDIN\***

Necmettin Erbakan University, Faculty of Science, Department of  
Mathematics and Computer Science, Konya, Turkey

\*Selçuk University, Faculty of Science, Department of Mathematics,  
Konya, Turkey

gckizilkan@konya.edu.tr, aduman@konya.edu.tr, kaydin@selcuk.edu.tr

### Abstract

Selection of step size is one of the most important concepts in numerical integration of differential equation systems. Even to use constant step size, it must be investigated how should be selected the step size in the first step of numerical integration. Because, if the selected step size is large in numerical integration, computed solution can diverge from the exact solution. And if the chosen step size is small; calculation time, number of arithmetic operations, the calculation errors start to increase. So, it will be sensible to use small step sizes in the region where the solution changes rapidly and to use bigger step size in the region where the solution changes slowly. So, it is not practical to use constant step size in numerical integration. In literature, step size strategies have been given for the numerical integration. One of these strategies is given for the Cauchy problem

$$X'(t) = AX(t) + \varphi(t, X), X(t_0) = X_0 \quad (0.0.4)$$

where  $A = (a_{ij}) \in R^{N \times N}$ ,  $X \in R^N$  and  $\varphi \in C^1([t_0 - T, t_0 + T] \times R^N)$  in [3,4].



Many dynamical system models are represented by non-linear differential equation systems as in (0.0.4). The epidemic models is one of these systems also attracted attention in recent years (for example see, [1,2,5,6,7]). The classical epidemic model is SIR model.

In this study, we have aimed to investigate the effectiveness of the variable step size strategy for some SIR epidemic models. We have applied the variable step size strategy to the SIR model and its modifications.

**Keywords:** Step size strategy, variable step size, epidemic model, SIR model, system of non-linear differential equations.

### Referances

- [1] Chauhan, S., Misra, O.P., Dhar, J., *Stability analysis of SIR model with vaccination*, Am. J. Comput. Appl. Math., Vol. 4, No. 1, 17-23, 2014.
- [2] Chinviriyasit, S., Chinviriyasit, W., *Numerical modelling of on SIR epidemic model with diffusion*, Appl. Math. Comput., Vol. 216, 395-409, 2010.
- [3] Çelik Kızıllan, G., *Step size strategies on the numerical integration of the systems of differential equations*, Ph.D. Thesis, Selcuk University Graduate Natural and Applied Sciences, Konya (in Turkish), 2009.
- [4] Çelik Kızıllan, G., Aydın, K., *Step size strategies for the numerical integration of systems of differential equations*, J. Comput. Appl. Math., Vol. 236, No. 15, 3805-3816, 2012.
- [5] Harko, T., Lobo, S.N.F., Mak, M.K., *Exact analytical solutions of the susceptible- infected- recovered (SIR) epidemic model and of the SIR model with equal death and birt dates*, Appl. Math. Comput., Vol. 236, 84-94, 2014.
- [6] Keeling, J.M., Rohani, P., *Modeling infectious diseases in humans and animals*, Princeton University Press, New Jersey, 2008.
- [7] Murray, J. D., *Mathematical biology: I. An introduction*, Third Edition, Springer, New York, 2002.

## When is an Archimedean $f$ -Algebra Finite Dimensional?

Faruk POLAT

Çankırı Karatekin University, Department of Mathematics, Çankırı,  
Turkey  
faruk.polat@gmail.com

Abstract

In this note, we give necessary and sufficient conditions for an Archimedean  $f$ -algebra to be of finite dimensional. As an application, we give a positive answer to a question raised by Bresar in [1].

### Referances

[1] Brešar, M., *Finite dimensional zero product determined algebras are generated by idempotents*, Expo Math, Vol. **34**, No.1, 130-143, 2016.

## By Calculating for Some Linear Positive Operators to Compare of the Errors in the Approximations

**Muzeyyen OZHAVZALI**

Kırıkkale University, Art and Science Faculty, Department of Mathematics, Kırıkkale, Turkey  
thavzalimuzeyyen@hotmail.com

### Abstract

In mathematics, we investigate a Korovkin-type approximation theorem for sequences of positive linear operators on the space of all continuous real valued functions defined on  $[a, b]$  in "Approximation theory". In this paper, we get some approximation properties for sequences of positive linear operators constructed by means of the Bernstein operator and give a Korovkin-type approximations properties for them. We research convergence and approximation properties for type generalized Stancu operators and Bernstein operator to give some examples. We also made a comparison between the approximations obtained by them with calculating the errors in the approximations for different continuous functions. Recently, some authors draw graphics of some modified operators and calculating the errors in approximations[1,2]. Figures of these kind of operator are very difficult because these operators have many of properties such that integrals, summations etc. Consequently figures and numerical results verify the theoretical results in the view of different aspects.

**Keywords:** Approximation, Positive linear operators, Korovkin-type theorem, Comparison, Errors, Figures.

### Referances

- [1] Buyukyazici, İ., *Approximation by Stancu-Chlodowsky polynomials*, Computers and Mathematics with Applications 59 (2009) 274-282 Berens H., Lorentz G.G., *Inverse theorems for Bernstein polynomials*, Indiana Univ. Math. J., 21(8)(1972), 693-708.
- [2] Buyukyazici, İ., *A Korovkin Type Approximation Theorem and Its Application*, Fasciculi Mathematici, Nr 45, 2010,
- [3] Cheney E. W., Sharma A., *Bersntein power series*, Canad. J. Math. 16, 241-253 (1960).
- [4] D.D.Stancu, *Approximation of functions by a new class of linear polinomial operators*, REv. Roumaine Math Pures Appl. 13(1968) 1113-1194.
- [5] Gadjiev, A.D., *On Korovkin Type Theorems*, Math. Notes, 20,(5-6), 996-998, 1976.
- [6] Korovkin P. P., *Linear operators and approximation theory*, Hindustan Publish Co., Delphi (1960).

## Intuitionistic Fuzzy Soft Neighborhoods

**Banu PAZAR VAROL, Abdülkadir AYGÜNOĞLU,  
Halis AYGÜN**  
Department of Mathematics, Kocaeli University, Umuttepe Campus,  
41380, Kocaeli - Turkey  
banupazar@kocaeli.edu.tr, abdulcadir.aygunoglu@kocaeli.edu.tr,  
halis@kocaeli.edu.tr

### Abstract

In this paper, we introduce the concept of intuitionistic fuzzy soft point as a generalization of intuitionistic fuzzy point and study some basic properties. We consider the neighborhood structures of an intuitionistic fuzzy point and generate an intuitionistic fuzzy soft topology by using the systems of neighborhood.

**Keywords:** Intuitionistic fuzzy soft set, intuitionistic fuzzy soft point, intuitionistic fuzzy soft topology, neighborhood.

### Referances

- [1] K. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, 64(2)87-96, 1986.
- [2] Z. Li and R. Cui, *On the topological structure of intuitionistic fuzzy soft set*, Annals of Fuzzy Mathematics and Informatics, 5(1), 229-239, 2013.
- [3] P.K. Maji, A.R. Roy and R. Biswas, *On intuitionistic fuzzy soft sets*, Journal of Fuzzy Mathematics 12(3), 669-683, 2004.

- [4] D. Molodtsov, *Soft set theory-First results*, Computers and Mathematics with Application, 37(4/5), 19-31, 1999.
- [5] B.Pazar Varol, A. Aygünoğlu and H. Aygün, *Neighborhood structures of fuzzy soft topological spaces*, Journal of Intelligent and Fuzzy Systems, 27(4), 2127-2135, 2014.
- [6] J. L. Seok and P.L. Eun, *The category of intuitionistic fuzzy topological spaces*, Bull. Korean Math. Soc. 37(1), 63-76, 2000.
- [7] K.Serkan and M. Akdağ, *On intuitionistic fuzzy soft continuous mappings*, Journal of New Results in Science, 4, 55-70, (2014).
- [8] K. Serkan and I. M. Akif, *Connectedness on intuitionistic fuzzy soft topological spaces*, Journal of New Theory, 1, 02-16, 2015.
- [9] Y.Yin, H. Li and Y.B. Jun, *On algebraic structures of intuitionistic fuzzy soft sets*, Computers and Mathematics with Applications 61, 2896-2911, 2012.
- [10] L.A. Zadeh, *Fuzzy sets*, Information and Control, 8, 338 - 353, 1965.

## On Nonlightlike Offset Curves in Minkowski 3-Space

Melek ERDOĞDU

Necmettin Erbakan University, Konya, Turkey  
merdogdu@konya.edu.tr

### Abstract

In this study, the offset curves of nonlightlike curves are investigated in three different cases. Then the curvature, torsion and arclength of a given offset curve are expressed in terms of the curvature, torsion of the main curve and constants A and B for each case. Moreover, it is proved that the offset curve constitutes another Bertrand curve.

**Keywords:** Bertrand Curve, Offset Curve, Minkowski Space.

### References

- [1] Arrondo, E., Sendra, J., Sendra, R., *Genus formula for generalized offset curves*, Journal of Pure and Applied Algebra, Vol. **136**, 199-209, 1999.
- [2] Babaarslan, M., Yaylı, Y., *On Space-like constant slope surfaces and Bertrand curves in Minkowski 3-space*, arXiv:1112.1504v6., 2014.
- [3] Ding, Q., Inoguchi, J., *Schrödinger flows, binormal motion for curves and second AKNS-hierarchies*, Chaos Solitons and Fractals, Vol. **21**, 669-677, 2004.
- [4] Inoguchi, J., *Biharmonic curves in Minkowski 3-space*, International Journal of Mathematics and Mathematical Sciences, Vol. **21**, 1365-1368, 2003.

- [5] Maekawa, T., *An overview of offset curves and surfaces*, Computer-Aided Design, Vol. **31**, 165-173, 1999.
- [6] Pham, B., *Offset curves and surfaces: a brief survey*, Computer-Aided Design, Vol. **24**, 223-239, 1992.
- [7] Schief, W. K., *On the integrability of Bertrand curves and Razzaboni surfaces*, Journal of Geometry and Physics, Vol. **45**, 130-150, 2003.

## Weierstrass Representation, Degree and Classes of the Surfaces in the Four Dimensional Euclidean Space

**Erhan GÜLER, Ömer KIŞI, Semra SARAÇOĞLU ÇELİK**  
Bartın University, Faculty of Science, Department of Mathematics  
ergler@gmail.com, eguler@bartin.edu.tr, okisi@bartin.edu.tr,  
ssaracoglu@bartin.edu.tr

### Abstract

In this talk, we present on a minimal surface using Weierstrass representation in the four dimensional Euclidean space. We compute implicit equations, degree and class of the surface.

**Keywords:** 4-space, Weierstrass representation, minimal surface, degree, class.

### Referances

- [1] Eisenhart, L.P. *A Treatise on the Differential Geometry of Curves and Surfaces*, Dover Publications, N.Y. 1909.
- [2] Hoffman, D.A., Osserman, R. *The Geometry of the Generalized Gauss Map*. Memoirs of the AMS, 1980.
- [3] Nitsche, J.C.C. *Lectures on Minimal Surfaces. Vol. 1. Introduction, Fundamentals, Geometry and Basic Boundary Value Problems*. Cambridge Un. Press, Cambridge, 1989.
- [4] Osserman, R. *A Survey of Minimal Surfaces*. Van Nostrand Reinhold Co., New York-London-Melbourne, 1969.
- [5] Weierstrass, K. *Untersuchungen über die flächen, deren mittlere Krümmung überall gleich Null ist*. Monatsber d Berliner Akad., 612-625, 1866.
- [6] Weierstrass, K., *Mathematische Werke*. Vol. 3, Mayer & Muller, Berlin, 1903.

## A New Type Graph and Their Parameters

Merve ÖZDEMİR, Nihat AKGÜNEŞ

Department of Mathematics, Computer Science, Necmettin Erbakan  
University, Konya, Turkey  
4232merve@gmail.com, nakgunes@konya.edu.tr

### Abstract

The graph theory has been improved fastly since it has applications in different fields of science. In this paper a special algebraic graph has been defined then some parameters of this graph have been studied.

**Keywords:** Graph, Graph Parameter.

### Referances

- [1] Gross, J.L. and Yellen, J. , *Handbook of graph theory* , Chapman Hall, CRC Press, 2004.
- [2] Das, K.Ch. , Akgüneş, N. and Çevik, A.S. , *On a graph of monogenic semigroups* , Journal of Inequalities and Applications , 2013, 2013:44.
- [3] Klotz, W. and Sander, T. , *Some properties of unitary cayley graphs* , The Electronic Journal of Combinatorics 14, 2007, R45.

## The Dot Product Graph of Monogenic Semigroup

Büşra ÇAĞAN, Nihat AKGÜNEŞ

Department of Mathematics, Computer Science, Necmettin Erbakan  
University, Konya, Turkey  
bsrcgn@gmail.com, nakgunes@konya.edu.tr

### Abstract

We can identify  $S$  as a cartesian product of finite times a finite semigroup  $S_M^n$  which has some elements like  $\{0, x, x^2, \dots, x^n\}$  Let  $\Gamma(S)$  be a dot product graph whose vertices are the nonzero elements of  $S$ . In this study we are going to analyze some parameters of  $\Gamma(S)$ .

**Keywords:** Dot Product, Monogenic Semigroup, Graph.

### Referances

- [1] Gross, J.L. and Yellen, J. , *Handbook of graph theory* , Chapman Hall, CRC Press, 2004.
- [2] Das, K.Ch. , Akgüneş, N. and Çevik, A.S. , *On a graph of monogenic semigroups* , Journal of Inequalities and Applications , 2013, 2013:44.
- [3] Badawi, A. , *On the dot product graph of a commutative ring* , Communications in Algebra 43, 2015, 43-50.

## Some Number Theoretical Results Related to the Suborbital Graphs for the Congruence Subgroup $\Gamma_0\left(\frac{n}{h}\right)$

Seda ÖZTÜRK

Karadeniz Technical University, Department of Mathematics  
seda.ozturk.seda@gmail.com

### Abstract

In this work, we study the congruence subgroup  $\Gamma_0\left(\frac{n}{h}\right)$  of the Modular group  $\Gamma$  acting transitively on the subset  $\bar{Q}(h)$ . From the suborbital graph  $F(1, n)$  we obtain some interesting number theoretical results, for instance, for all  $n \in \mathbb{N}$ , the numbers  $n(n-4)b^2 - 4$  are not squares.

**Keywords:** Graph Theory, Number Theory.

### Referances

- [1] G. A. Jones, D. Singerman, and K. Wicks, *The Modular Group and Generalized Farey Graphs*, London Mathematical Society Lecture Note Series, Cambridge University Press, **160**, 1991.

## Schur Stabilitiy in Floating Point Arithmetic: Systems with Periodic Coefficients

Ali Osman ÇIBIKDİKEN, Ahmet DUMAN,  
Kemal AYDIN

Necmettin Erbakan University, Department of Computer Engineering,  
Konya, Turkey  
Necmettin Erbakan University, Department of Mathematics and  
Computer Science, Konya, Turkey  
Selçuk University, Department of Mathematics, Konya, Turkey  
aocdiken@konya.edu.tr, aduman@konya.edu.tr, kaydin@selcuk.edu.tr

### Abstract

The representation of the numbers

$$\mathbb{F} = \mathbb{F}(\gamma, p_-, p_+, k) = \{0\} \cup \left\{ z \mid z = \pm \gamma^{p(z)} m_\gamma(z) \right\}, \quad (0.0.5)$$

which are called floating point numbers [2, 13, 17, 18]. Computers use floating point numbers for computing. These numbers are also called computer numbers or machine numbers [6, 14, 19, 21]. If the results of arithmetic operations are elements of set  $\mathbb{F}$ , they are directly stored in the memory. Otherwise they are stored with error [7, 11, 15, 16, 20, 22].

Let  $A_n$  be an  $N$ -dimensional periodic matrix ( $T$ -period) and difference equation system  $x_{n+1} = A_n x_n$ . The matrix  $X_n$  is called the fundamental matrix of the system, and the matrix  $X_T$  is called the monodromy matrix of the system [1, 3, 4, 5, 12]. The matrix  $Y_n = fl(A_{n-1} Y_{n-1})$  is the computed fundamental matrix  $X_N$  in floating point arithmetic. Cauchy problem of difference equation system can be written

$$fl(A_{n-1} Y_{n-1}) = Y_n = A_{n-1} Y_{n-1} + \phi_N; Y_0 = I, \quad (0.0.6)$$

where  $\phi_N$  is computation error of  $(A_{n-1} Y_{n-1})$ .

The effects of floating point arithmetic in the computation of the fundamental matrix  $X_N$  were investigated. Error bounds were obtained for  $\|X_N - Y_N\|$ , where  $A_N \in M_N(D)$ . The obtained results were investigated for Schur stability of the system [8, 9, 10]. These results were supported with numerical examples.

**Keywords:** floating point arithmetic, difference equation, error analysis, Schur stability.

### References

- [1] Agarwal R.P., *Difference Equations and Inequalities*, Pure and Applied Mathematics Series (second ed.), vol. 228, National University of Singapore, 2000.
- [2] Akın, O. and Bulgak H., *Linear difference equations and stability theory*, Selçuk University Research Centre of Applied Mathematics, No. 2, Konya, 1998 (in Turkish).
- [3] Aydın K., *The condition number for the asymptotic stability of the periodic ordinary differential systems*, Ph.D. Thesis, Selçuk University Graduate Natural and Applied Sciences, Konya, (1996) (in Turkish).



- [4] Aydın K., Bulgak H., Demidenko G.V., *Numeric characteristics for asymptotic stability of solutions to linear difference equations with periodic coefficients*, Sib. Math. J., 41 (6), pp. 1005-1014, 2000.
- [5] Aydın K., Bulgak H., Demidenko G.V., *Asymptotic stability of solutions to perturbed linear difference equations with periodic coefficients*, Sib. Math. J., 43 (3), pp. 389-401, 2002.
- [6] Behrooz P., *Computer Architecture: From Microprocessors to Supercomputers*, Oxford University Press (2005).
- [7] Çıbıkdiken A.O., *Error estimation of elementary matrix operations*, Master Thesis, Selçuk University Graduate Natural and Applied Sciences, Konya, 2002 (in Turkish).
- [8] Çıbıkdiken A.O., *The effect of floating point arithmetic on fundamental matrix computation of periodic linear difference equation system*, Ph.D. Thesis, Selçuk University Graduate Natural and Applied Sciences, Konya, 2008 (in Turkish).
- [9] Çıbıkdiken A.O., Aydın K., *Computation of the monodromy matrix in floating point arithmetic with the Wilkinson Model*, Computers & Mathematics with Applications, Volume 67, Issue 5, Pages 1186-1194, 2014.
- [10] Çıbıkdiken A.O., Aydın K., *Computation of monodromy matrix on floating point arithmetic with Godunov Model*, Konuralp Journal of Mathematics, Volume 4, No 1, 2016.
- [11] Dahlquist G., Björck A., *Numerical Mathematics and Scientific Computation*, Vol. 1, SIAM (2008).
- [12] Elaydi S.N., *An Introduction to Difference Equations*, Springer-Verlag, New York 1996.
- [13] Godunov, S.K., Antonov, A.G., Kiriluk, O.P., V.I. Kostin, *Guaranteed Accuracy in Mathematical Computations*, Prentice-Hall, Englewood Cliffs, NJ (1993).
- [14] Goldberg D., *What every computer scientist should know about floating-point arithmetic*, J. ACM Comput. Surv., 23 (1) (1991), pp. 5-48.
- [15] Golub G.H., Van Loan C.F., *Matrix Computations*, (third ed.)The Johns Hopkins University Press, Baltimore (1996).
- [16] Higham N.J., *Accuracy and Stability of Numerical Algorithms*, SIAM (1996).
- [17] Kulisch, U.W., *Mathematical foundation of computer arithmetic*, IEEE Trans. Comput., C-26 (7) (1977), pp. 610-621.
- [18] Kulisch U.W., Miranker W.L., *Computer Arithmetic in Theory and Practice*, Academic Press Inc. (1981).
- [19] Overton M.L., *Numerical Computing with IEEE Floating Point Arithmetic*, SIAM (2001).
- [20] Shampine L.F., Allen Jr. R.C., Pruess S., *Fundamentals of Numerical Computing*, John Wiley & Sons Inc. (1997).
- [21] Sterbenz P.H., *Floating-Point Computation*, Prentice-Hall Inc., Englewood Cliffs, NJ (1974).
- [22] Wilkinson J.H., *Rounding Errors in Algebraic Processes*, Prentice-Hall Inc. (1963).

## $\mathcal{I}$ -Limit Inferior and $\mathcal{I}$ -Limit Superior of Sequences of Sets

Ömer KİŞİ, Fatih NURAY

Bartın University, Faculty of Science, Department of Mathematics  
Afyon Kocatepe University, Faculty of Science, Department of Mathematics

okisi@bartin.edu.tr, omerkisi@hotmail.com, fnuray@aku.edu.tr

### Abstract

In this talk, we extend the concepts of  $\mathcal{I}$ -Limit superior and  $\mathcal{I}$ -Limit inferior for real number sequences to  $\mathcal{I}$ -Limit superior and  $\mathcal{I}$ -Limit inferior for sequences of sets, study their certain properties and establish some basic theorems.

**Keywords:** Statistical convergence,  $I$ -convergence and  $I^*$ -convergence, sequence of sets, Wijsman convergence.

### References

- [1] Baronti, M. and Papini, P. (1986). Convergence of sequences of sets. In methods of functional analysis in approximation theory, ISNM 76, Birkhauser, Basel, 133-155.
- [2] Beer, G. (1985). *On convergence of closed sets in a metric space and distance functions*, Bull. Austral. Math. Soc., **31**, 421-432.
- [3] Fast, H. (1951). *Sur la convergence statistique*, Coll. Math., **2**, 241-244.
- [4] Kiş, Ö. and Nuray, F. (2013). *New Convergence Definitions for Sequence of Sets*, Abstract and Applied Analysis Volume 2013, Article ID 852796, 6 pages <http://dx.doi.org/10.1155/2013/852796>
- [5] Kostyrko, P., Šalát, T. and Wilezyński, W. (2000),  $\mathcal{I}$ -Convergence, Real Analysis Exchange, **26**, 669-680.
- [6] Demirci, H. (2001).  $\mathcal{I}$ -limit superior and  $\mathcal{I}$ -limit inferior, Mathematical Communications, **6**, 165-172.
- [7] Wijsman, R. A. (1965). *Convergence of sequences of convex sets, cones and functions*, Bull. Amer. Math. Soc., **70**, 186-188.

## A Ditopological Fuzzy Structural View of Inverse Systems and Inverse Limits

Filiz YILDIZ

Hacettepe University, Faculty of Science, Department of Mathematics,  
Beytepe, Ankara  
yfiliz@hacettepe.edu.tr

### Abstract

One of the theories defined to develop some complement-free concepts is the *texture (fuzzy structure) theory* and it was constructed in [1] as a point-based setting for the study of classical sets and fuzzy sets. Add to that, the notion of *ditopology* described on a texture is essentially a topology for which there is no a priori relation between the open and closed sets, and thus ditopological fuzzy structures [1] were conceived as a unified setting for the study of fuzzy topology. Especially, some useful relationships with fuzzy topology may be found in [3].

In the previous study [2], the foundations of a corresponding theory of *inverse (projective) systems* and their limits, called *inverse limits* were laid in the category **ifPTex** of plain textures which are special types of textures, and point functions satisfying a compatibility condition, named *w*-preserving. Therefore firstly, a detailed analysis of inverse systems and inverse limits was presented in [2] insofar as the category of plain textures is concerned. Evidently, this theory was constituted as an analogue of the inverse system theory in the classical categories **Set**, **Top** and **Rng** in algebra, simultaneously.

As the main theme of this presentation, a suitable theory of inverse systems and their limits is established for some subcategories of the category **ifPDitop** topological over **ifPTex**, whose objects are ditopological fuzzy structures which have plain texturing and morphisms are bicontinuous **ifPTex**-morphisms. In addition, many useful properties of inverse limits in **ifPDitop** are studied via examples in the context of ditopological fuzzy structures, as natural counterparts of the classical cases.

**Keywords:** Inverse Limit, Fuzzy Topology, Category, Texture, Ditopology.

### Referances

- [1] Brown, L. M., *Ditopological fuzzy structures I*. Fuzzy Systems and A.I. Magazine, 3 (1993)
- [2] Yıldız, F., *Inverse Systems and Inverse Limits in the Category of Plain Textures*, Topology and Its Applications , Vol. 201, pp. 217-234, 2016.
- [3] Yıldız, F. and Brown, L.M., *Extended Real Dicomactness and an Application to Hutton Spaces*, Fuzzy Sets and Systems, 227, 74–95, 2013.

## On Nullnorms on Bounded Lattices

Emel AŞICI

Department of Software Engineering, Faculty of Technology, Karadeniz  
 Technical University, 61830 Trabzon, Turkey  
 emelkalin@hotmail.com

### Abstract

t-operators and nullnorms were introduced in [9], [7] respectively, which are also generalizations of the notions of t-norms and t-conorms. And then in [10], it is pointed out that nullnorms and t-operators are equivalent since they have the same block structures in  $[0, 1]^2$ . Namely, if a binary operator  $F$  is a nullnorm then it is also a t-operator and vice versa.

**Definition 0.0.1** *Let  $(L, \leq, 0, 1)$  be a bounded lattice. A commutative, associative, non-decreasing in each variable function  $F : L^2 \rightarrow L$  is called a nullnorm if there is an element  $a \in L$  such that  $F(x, 0) = x$  for all  $x \leq a$ ,  $F(x, 1) = x$  for all  $x \geq a$ .*

In this study, given a bounded lattice  $L$  and a nullnorm on it, taking into account the properties of nullnorms, we investigate an order induced by nullnorms and equivalence relation on bounded lattice. In this way, we obtain that interesting results.

**Keywords:** Nullnorm, Bounded lattice, Partial order.

**Acknowledgement:** The work on this study was supported by the Research Fund of Karadeniz Technical University, project number FBB-2015-5218.

### References

- [1] Aşıcı, E. and Karaçal, F., *On the T-partial order and properties*, Inf.Sci., Vol. **267**, 323-333, 2014.
- [2] Aşıcı, E. and Karaçal, F., *Incomparability with respect to the triangular order*, Kybernetika, Vol. **52**, 15-27, 2016.
- [3] Birkhoff, G., *Lattice Theory*, 3 rd edition, Providence, 1967.
- [4] Drewniak, J., Drygaś, P. and Rak, E., *Distributivity between uninorms and nullnorms*, Fuzzy Sets Syst., Vol. **159**, 1646-1657, 2008.
- [5] Drygaś, P., *A characterization of idempotent nullnorms*, Fuzzy Sets Syst., Vol. **145**, 455-461, 2004.
- [6] Drygaś, P. and E. Rak, *Distributivity equation in the class of 2-uninorms*, Fuzzy Sets Syst. Vol. **291**, 82-97, 2016.
- [7] Calvo, T., De Baets, B. and Fodor J., *The functional equations of Frank and Alsina for uninorms and nullnorms*, Fuzzy Sets Syst. Vol. **120**, 385-394, 2001.
- [8] Karaçal, F. and Aşıcı, E., *Some notes on T-partial order*, J. Inequal. Appl. **2013**, 219, 2013.
- [9] Mas, M., Mayor, G. and Torrens, J., *T-operators*, Int. J. Uncertain. Fuzz. Knowl.-Based Syst. Vol. **7**, 31-50, 1999.

[10] Mas, M., Mayor, G. and Torrens, J., *The distributivity condition for uni-norms and  $t$ -operators*, Fuzzy Sets Syst. Vol. **128**, 209-225, 2002.

## Ideal Version of Weighted Lacunary Statistical Convergence of Sequences of Order $\alpha$

**Sukran KONCA, Ergin GENÇ, Selman EKİN**

Department of Mathematics, Bitlis Eren University,

13000, Bitlis, Turkey

skonca@beu.edu.tr, ergingenc4444@gmail.com, selmanekin@hotmail.com

### Abstract

In this work, we are interested in ideal version of weighted lacunary statistical convergence of sequences of order  $\alpha$  and we examine some inclusion relations.

**Keywords:**  $I$ -convergence;  $I$ -statistical convergence; weighted lacunary  $I$ -statistical convergence of order  $\alpha$ ; sequence space

### References

- [1] Lahiri, Kumar B. and Das, P.  *$I$  and  $I^*$ -convergence in topological spaces*, Mathematica Bohemica, Vol. **130**, No. 2, 153–160, 2005.
- [2] Kostyrko, P., Macaj, M., Salat T. and Sleziak, M.  *$I$ -convergence and extremal  $I$ -limit points*, Mathematica Slovaca, Vol. **55** No. 4 (2005) 443–464.
- [3] Çolak, R. *Statistical convergence of order  $\alpha$* , Modern Methods in Analysis and Its Applications, New Delhi, India: Anamaya Pub., 121–129, 2010.
- [4] Bhunia, S., Das, P. and Pal, S. *Restricting statistical convergence*, Acta Mathematica Hungarica, **134** No. 1-2, 153–161, 2011.
- [5] Et, M., Çınar M. and Karakas, M. *On  $\lambda$ -statistical convergence of order  $\alpha$  of sequences of function*, Journal of Inequalities and Applications, Vol. **2013** No. 1 (2013) 1–8.
- [6] Şengül, H. and Et, M. *On lacunary statistical convergence of order  $\alpha$* , Acta Mathematica Scientia Series, Vol. **34** No. 2 (2014) 473–482.
- [7] Başarır, M. and Konca, Ş. *On some spaces of lacunary convergent sequences derived by Nörlund-type mean and weighted lacunary statistical convergence*, Arab Journal of Mathematical Sciences, Vol. **20** No. 2 (2014) 250–263.
- [8] Başarır, M. and Konca, Ş. *Weighted lacunary statistical convergence in locally solid Riesz spaces*, Filomat Vol. **28** No. 10 (2014) 2059–2067.

## $AK(S)$ and $AB(S)$ Properties of a $K$ -Space

**Mahmut KARAKUŞ**

Department of Mathematics, Yüzüncü Yıl University, Van, Turkey  
matfonks@gmail.com

### Abstract

A typical sum  $s$  on a  $K$ -space  $S$  has often the representation,

$$s(z) = \lim_{\gamma \in \Gamma} \sum_k u_{\gamma k} z_k, \quad z = (z_k) \in S \quad (0.0.7)$$

where  $\Gamma$  is a directed index set and  $u_\gamma = (u_{\gamma k}) \in \phi$ , the space of finitely non-zero sequences, for each  $\gamma \in \Gamma$ . Let a  $K$ -space  $S$  be equipped with a sum (0.0.7). Then, for each  $x = (x_k)$  and  $\gamma \in \Gamma$ , the sequence  $P_\gamma(x) = \sum_k u_{\gamma k} x_k \delta^k$ , ( $\gamma \in \Gamma$ ) is called the  $\gamma$ th  $S$ -section of  $x$  [2]. Here  $\delta^k$  is the sequence whose  $k$ th component is 1 all the others are 0.

If  $\lambda \supset \phi$  is a  $K$  space, then Boos and Leiger defined the spaces  $\lambda_{AB(S)}$  and  $\lambda_{AK(S)}$  in [?] as

$$\lambda_{AB(S)} = \{x \in \omega | (P_\gamma(x))_{\gamma \in \Gamma} \text{ is a bounded net in } \lambda\},$$

and

$$\lambda_{AK(S)} = \{x \in (\lambda_{AB(S)} \cap \lambda) | \lim_\gamma P_\gamma(x) \text{ exists in } \lambda\}.$$

In this work, we investigate some properties of these spaces and give some theorems related to the duals.

**Keywords:**  $K$ - spaces,  $n$ -th section of a sequence,  $\beta$ -,  $\gamma$ -,  $f$ -duality.

### References

- [1] J. Boos, T. Leiger, Dual pairs of sequence spaces, *Int. J. Math. Math. Sci.*, **28**(2001): 9-23.
- [2] J. Boos, T. Leiger, Dual pairs of sequence spaces II, *Proc. Estonian Acad. Sci. Phys. Math.*, **51**(2002): 3-17.
- [3] J. Boos, T. Leiger, Dual pairs of sequence spaces III, *J. Math. Anal. Appl.*, **324** (2006): 1213-1227.
- [4] J. Boos, Classical and Modern Methods in Summability, *Oxford University Press. New York, Oxford*, 2000.
- [5] M. Buntinas, Convergent and bounded Cesàro sections in FK-spaces, *Math. Zeitschr.*, **121** (1971): 191-200.
- [6] M. Buntinas, On Toeplitz sections in sequence spaces, *Math. Proc. Camb. Phil. Soc.*, **78** (1975), 451-460.
- [7] W. H. Ruckle, An abstract concept of the sum of a numerical series, *Can. J. Math.*, 1970, **22**: 863-874.

# On the Second Homology of the Schützenberger Product of Monoids

Melek YAĞCI

Department of Mathematics, Çukurova University, Adana, 01330, Turkey  
msenol@cu.edu.tr

## Abstract

For two finite monoids  $S$  and  $T$ , we prove that the second integral homology of the Schützenberger product  $S\Diamond T$  is equal to

$$H_2(S\Diamond T) = H_2(S) \times H_2(T) \times (H_1(S) \otimes_{\mathbb{Z}} H_1(T))$$

as the second integral homology of the direct product of two monoids.

This is joint work with Hayrullah Ayık and Leyla Bugay.

(Melek Yağcı is supported by Ç.Ü. BAP.)

**Keywords:** Monoid, Schützenberger product, Second integral homology.

## Referances

- [1] Çevik, A.S., *Minimal but inefficient presentations of the semi-direct products of some monoids*, Semigroup Forum, **66**, 1-17, 2003.
- [2] Çevik, A.S., *The  $p$ -Cockcroft property of the semi-direct products of monoids*, Internat. J. Algebra Comput. **13**, 1-16, 2003.
- [3] Campbell, C.M., Mitchell, J.D., Ruskuc, N., *On defining groups efficiently without using inverses*, Math. Proc. Camb. Phil. Soc. **133**, 31-36, 2002.
- [4] Squier, C., *Word problems and a homological finiteness condition for monoids*, Journal Pure Appl. Algebra, **49**, 201-217, 1987.
- [5] Johnson D.L., *Presentations of Groups*, Cambridge Univ. Press, Cambridge, 1990.
- [6] Ayık, H., Campbell, C.M., O'Connor, J.J., *On the efficiency of the direct products of Monogenic Monoids*, Algebra Colloq. **14**, 279-284, 2007.
- [7] Ayık, H., Campbell, C.M., O'Connor, J.J., Ruskuc, N., *Minimal presentations and efficiency of semigroups*, Semigroup Forum, **60**, 231-242, 2000.
- [8] Ayık, H., Campbell, C.M., O'Connor, J.J., Ruskuc, N., *On the efficiency of finite simple semigroups*, Turkish Journal of Math. **24**, 129-146, 2000.
- [9] Howie, J.M., *Fundamentals of Semigroup Theory*, New York, Oxford University Press, 1995.
- [10] Howie, J.M., Ruskuc, N., *Constructions and presentations for monoids*, Comm. Algebra, **49**, 6209-6224, 1994.

- [11] Guba, V.S., Pride, S.J., *Low dimensional (co)homology of free Burnside monoids*, Journal Pure Appl. Algebra, **108**, 61-79, 1996.  
 [12] Guba, V.S., Pride, S.J., *On the left and right cohomological dimension of monoids*, Bull London Math. Soc. **30**, 391-396, 1998.

## New Sequence Spaces with Respect to a Sequence of Modulus Functions

**Ömer KİŞİ, Semra SARAÇOĞLU ÇELİK, Erhan GÜLER**  
 Bartın University, Faculty of Science, Department of Mathematics  
 okisi@bartin.edu.tr, ssaracoglu@bartin.edu.tr, ergler@gmail.com,  
 eguler@bartin.edu.tr

### Abstract

In this talk, we introduce the notion of  $A^{\mathcal{I}}$ -invariant statistical convergence,  $A^{\mathcal{I}}$ -lacunary invariant statistical convergence with respect to a sequence of modulus functions. We establish some inclusion relations between these spaces under some conditions.

**Keywords:** Lacunary invariant statistical convergence; Invariant statistical convergence, ideal convergence, modulus function.

### References

- [1] H. Nakano, *Concave modulars*, J. Math. Soc. Jpn. 5 (1953) 29-49.  
 [2] T. Bilgin, *Lacunary strong  $A$ -convergence with respect to a modulus*, Mathematica XLVI(4) (2001) 39-46.  
 [3] P. Kostyrko, M. Macaj, T. Salat,  $\mathcal{I}$ -convergence, Real Anal. Exchange, 26 (2) (2000) 669-686.  
 [4] E. Savaş, F. Nuray, On  $\sigma$ -statistically convergence and lacunary  $\sigma$ -statistically convergence. Math. Slovaca, 43 (3), 1993, 309-315.  
 [5] E. Savaş, P. Das, A generalized statistical convergence via ideals, App. Math. Lett. 24, 2011, 826-830.

## TF-Type Hypersurfaces in 4-Space

**Erhan GÜLER, Ömer KİŞİ, Semra SARAÇOĞLU ÇELİK**  
 Bartın University, Faculty of Science, Department of Mathematics



ergler@gmail.com, eguler@bartin.edu.tr, okisi@bartin.edu.tr,  
ssaracoglu@bartin.edu.tr

### Abstract

We study on translation and factorable hypersurfaces in the four dimensional Euclidean space. We calculate implicit algebraic equations of the hypersurfaces.

**Keywords:** 4-space, translation hypersurface, factorable hypersurface, algebraic equation.

### Referances

- [1] Dillen, F., Goemans, W., Van de Woestyne, I. *Translation surfaces of Weingarten type in 3-space*. Bull. Transilv. Univ. Braşov Ser. III 1 (50) 109-122, 2008.
- [2] Do Carmo, M.P. , *Differential Geometry of Curves and Surfaces*, Prentice-Hall, Englewood Cliffs, 1976.
- [3] Liu, H. *Translation surfaces with dependent Gaussian and mean curvature in 3-dimensional spaces*. (Chinese) J. Northeast Univ. Tech. 14 (1), 88-93, 1993.
- [4] Verstraelen, L., Walrave, J., Yaprak, S. *The minimal translation surfaces in Euclidean space*. Soochow J. Math. 20 (1), 77-82, 1994.

## A Note on $q$ -Binomial Coefficients

Özge ÇOLAKOĞLU HAVARE, Hamza MENKEN  
Department of Mathematics, Mersin University, Mersin, Turkey  
ozgecolakoglu@mersin.edu.tr; hmenken@mersin.edu.tr

### Abstract

The  $q$ -calculus has been developing fast. In the present work we study on a  $q$ -extension of binomial coefficients. The infinite sum of  $q$ -extension of binomial coefficients is obtained. Then, by using its infinite sum, we obtain Volkenborn integral value of  $q$ -extension of binomial coefficients.

**Keywords:**  $p$ -adic number, Indefinite sum,  $q$ -analogue of the binomial coefficients, Volkenborn integral.

### Referances

- [1] Araci, S. and Açıkgöz, M., *A note on the values of weighted  $q$ -Bernstein polynomials and weighted  $q$ -Genocchi numbers*, Advances in Differences Equations, 1-9, 2015.
- [2] Foata, D. and Han G., *The  $q$ -series in combinatorics; Permütation Statistics*, Lecture Note, 2004.
- [3] Schikhof, W. H., *Ultrametric Calculus: An Introduction to  $p$ -adic Analysis*, Cambridge University Pres, 1984.
- [4] Volkenborn A., *Ein  $p$ -adisches Integral und seine Anwendungen. I*, Manuscripta Math. 7-4, 341–373, 1972.
- [5] Volkenborn A., *Ein  $p$ -adisches Integral und seine Anwendungen II*, Manuscripta Math. 12, 17-46, 1974.

## On Statistical Convergence of Sequences of $p$ -Adic Numbers

**Mehmet Cihan BOZDAĞ, Hamza MENKEN**  
Mersin University, Mersin-TURKEY  
mehmetcihanbozdag@gmail.com, hmenken@mersin.edu.tr

### Abstract

Let  $p$  be a fixed prime number. By  $\mathbb{Q}_p$  we denote the field of  $p$ -adic numbers, the completion of the rational numbers field  $\mathbb{Q}$  with respect to the  $p$ -adic norm  $|\cdot|_p$ . The concept of statistical convergence was introduced by H. Fast (1951) [1] and R. C. Buck (1953) [2] independently for real or complex sequences. This concept was studied by T.Salat (1980) [3], J. A. Fridy (1985) [4] and many authors. We note that the field of  $p$ -adic numbers  $\mathbb{Q}_p$  is non-Archimedean, means that the ultrametric inequality is valid

$$|x + y|_p \leq \max \left\{ |x|_p, |y|_p \right\}$$

for all  $x, y \in \mathbb{Q}_p$ . In the present work we define the concept of statistical convergence of sequences for  $p$ -adic numbers and give some its properties.

**Keywords:**  $p$ -adic number, statistical convergence of sequence of  $p$ -adic numbers, statistical Cauchy sequence of  $p$ -adic numbers.

### Referances

- [1] H.Fast, Sur la convergence statistique, Colloq. Math., 241-244 (1951).
- [2] R. C Buck, Generalized Asymptotic Density, Amer. J. Math. 75 335–346 (1953)

- [3] T.Salat ,On statistically convergent sequences of real numbers, *Mathematica Slovaca*,vol.30 issue 2, pp. 139-150 (1980).  
 [4] Fridy, J.A., On statistical convergence, *Analysis* 5 (1985),301-313.  
 [5] Schikhof, W. H. , *Ultrametric Calculus*, Cambridge Univ. Press, New York, 2006.

## Lightlike Hypersurface of an Indefinite Kaehler Manifold with a Complex Semi-Symmetric Metric Connection

**Halil İbrahim YOLDAŞ, Erol YAŞAR**

Mersin University, Department of Mathematics Mersin, Turkey  
 hibrahimyoldas@gmail.com, yerol@mersin.edu.tr

### Abstract

In this paper, we study lightlike hypersurface of an indefinite Kaehler manifold admitting a complex semi-symmetric metric connection. We get the equations of Gauss and Codazzi. Then, we give some characterizations of lightlike hypersurface in an indefinite complex space form with a complex semi-symmetric metric connection. Finally, we show that the Ricci tensor of lightlike hypersurface of an indefinite Kaehler manifold with complex semi-symmetric metric connection is not symmetric.

**Keywords:** Lightlike Hypersurface, Indefinite Complex space form, Complex Semi-Symmetric Metric Connection, Levi-Civita connection, Ricci tensor.

## Intuitionistic Fuzzy Fractional Evolution Problem

**M. ELOMARI, S. MELLIANI, L. S. CHADLI**

LMACS, Laboratoire de Mathématiques Appliquées & Calcul Scientifique  
 Sultan Moulay Slimane University, PO Box 523, 23000 Beni Mellal  
 Morocco  
 said.melliani@gmail.com

### Abstract

We introduce the generalized intuitionistic fuzzy derivative, this concept used in order to give a generalized intuitionistic fuzzy Caputo fractional derivative. And we discuss the intuitionistic fuzzy fractional evolution problem.

**Keywords:** Generalized intuitionistic fuzzy Hukuhara difference, Generalized intuitionistic fuzzy derivative, generalized intuitionistic fuzzy Caputo-derivative, intuitionistic fuzzy fractional evolution problem.

### References

- [1] T. Allahviranloo, A. Armand and Z. Gouyandeh, Fuzzy fractional differential equations under generalized fuzzy Caputo derivative, *J. Intell. Fuzzy Systems*, 26 (2014) 1481–1490.
- [2] T. Allahviranloo, A. Armand, Z. Gouyandeh, H. Ghadiri, Existence and uniqueness of solutions for fuzzy fractional Volterra-Fredholm integro-differential equations, *J. Fuzzy Set Valued Analysis*, 2013 (2013) 1–9.
- [3] K. Atanassov, Intuitionistic fuzzy sets, *Fuzzy sets and Systems*, 20(1) (1986) 87–96.
- [4] V. Barbu, *Nonlinear differential equations of monotone types in Banach spaces*. Springer, Monographs in mathematics.
- [5] B. Bede and L. Stefanini, Generalized differentiability of fuzzy-valued functions, *Fuzzy Sets and Systems*, 230 (2013) 119–141.
- [6] H. Brezis and A. Pazy. *Convergence and Approximation of Semigroups of Nonlinear Operators in Banach Spaces*. *J. Funct. Ana.*, 9 (1972) 63–74.
- [7] G. Debreu, *Integration of correspondences*, in *Proc. Fifth Berkeley Symp. Math. Stat. Probab.*, Vol. 2, Part 1 (Univ. California Press, Berkeley, CA, 1967) 351–372.
- [8] M. Elomari, S. Melliani, R. Ettoussi, L. S. Chadli, Intuitionistic fuzzy semigroups, *Notes on Intuitionistic Fuzzy Sets*, 21(1) (2015) 43–50.
- [9] V. Lakshmikantham, S. Leela, *Nonlinear Differential Equations in Abstract Spaces*, Pergamon Press, New York, (1981).
- [10] O. Kaleva, Nonlinear iteration semigroup of fuzzy Cauchy problems, *Fuzzy sets and systems*, 209 (2012) 104–111.
- [11] O. Kaleva, On the convergence of fuzzy sets, *Fuzzy Sets and Systems*, 17 (1985) 53–65.
- [12] O. Kaleva, The Cauchy problem for fuzzy differential equations, *Fuzzy Set Systems*, 35 (1990) 366–389.
- [13] O. Kaleva, Fuzzy differential equations, *Fuzzy Sets Systems*, 24 (1987) 301–317.
- [14] T. Kato. *Nonlinear semigroup and evolution equations*, *J. Math. Soc. Japan.*, 19 (4) (1967).
- [15] N. Kosmatov, Integral equations and initial value problems for nonlinear differential equations of fractional order, *Nonlinear Anal.*, 70 (2009) 2521–2529.
- [16] Y. Komura, Nonlinear semigroup in Hilbert space, *J. Math. Soc. Japan*, 21(1969), 375–507.

- [17] S. Melliani, M. Elomari, L. S. Chadli, R. Ettoussi, Intuitionistic fuzzy metric space, *Notes on Intuitionistic Fuzzy Sets*, 21(1) (2015) 43–53.
- [18] S. Melliani, M. Elomari, L. S. Chadli, R. Ettoussi, Extension of Hukuhara difference in intuitionistic fuzzy set theory, *Notes on Intuitionistic Fuzzy Sets*, 21(4) (2015) 34–47.
- [19] S. Melliani, M. Elomari, L. S. Chadli, R. Ettoussi, Intuitionistic fuzzy metric space, *Notes on Intuitionistic Fuzzy Sets*, 21 (1) (2015) 43–53.
- [20] A. Pazy. *Semigroups of Linear Operators and Applications to Partial Differential Equations*, Springer-Verlag, 1983.
- [21] M. L. Puri and D.A. Ralescu, Differentials for fuzzy functions, *J. Math. Anal. Appl.* 91 (1983) 552–558.
- [22] M. L. Puri and D.A. Ralescu, Fuzzy random variables, *J. Math. Anal. Appl.* 114 (1986) 409–422.
- [23] M. L. Puri and D.A. Ralescu, Fuzzy random variables, *J. Math. Anal. Appl.* 114 (1986) 409–422.
- [24] K. Roshdi, M. Al Horani, A. Yousef, M. Sababheh, A new Definition Of Fractional Derivative, *J. Comput. Appl. Math.*, 264 (2014) 65–70.

## A General Tableaux Method for Contact Logics Interpreted over Intervals

**Philippe BALBIANI, Çiğdem GENCER,  
Zafer ÖZDEMİR**

Institut de recherche en informatique de Toulouse CNRS - Université de  
Toulouse

Department of Mathematics and Computer Science Sabancı University,  
Faculty of Engineering and Natural Sciences

Department of Mathematics and Computer Science Istanbul Kültür  
University - Faculty of Science and Letters

ozdemir.zafer@yahoo.com.tr

### Abstract

In this paper, we focus our attention on tableau methods for contact logics interpreted over intervals on the reals. Contact logics provide a natural framework for representing and reasoning about regions in several areas of computer science such as geological information systems, artificial intelligence and etc. In this paper, we focus our attention on tableau methods for contact logics interpreted over intervals on the reals. Contact logics provide a natural framework for representing and reasoning about regions in several areas of computer science such as geological information systems, artificial intelligence and etc.[1,3,4]. However,

while various tableau methods have been developed for classical logic, modal logics and intuitionistic logic, not much work has been done on tableau methods for contact logics [1,2]. We develop a general tableau method for contact logic interpreted over intervals. In this paper we give sound and complete tableaux-based decision procedures for contact logics. Developing such tableaux-based decision procedures, we obtain new decidability/complexity results.

### References

- [1] Balbiani, Philippe, Tinko Tinchev, and Dimiter Vakarelov. "Modal logics for region-based theories of space." *Fundamenta Informaticae* 81.1-3 (2007): 29-82.
- [2] Galton, Antony. "The mereotopology of discrete space." *International Conference on Spatial Information Theory*. Springer Berlin Heidelberg, 1999.
- [3] Galton, A. "Qualitative Spatial Change". Oxford University Press." (2000).
- [4] Wolter, Frank, and Michael Zakharyashev. "Spatial representation and reasoning in RCC-8 with Boolean region terms." *Proceedings of the 14th European Conference on Artificial Intelligence (ECAI 2000)*.

## Solving Intuitionistic Fuzzy Differential Equations with Linear Differential Operator by Adomain Decomposition Method

Suvankar BISWAS, Sanhita BANERJEE,  
Tapan Kumar ROY

Department of Mathematics, Indian Institute of Engineering Science and  
Technology, Shibpur, Howrah-711103, West Bengal, India  
suvo180591@gmail.com

### Abstract

In this paper we presented intuitionistic fuzzy differential equation with linear differential operator which can be of any order and it also involves nonlinear functional. So our solution procedure gives the solutions of a large area of problems involving intuitionistic fuzzy differential equations. Adomain decomposition method (ADM) has been used to find the approximate solution. Note that we used ADM which gives solution even for some nonlinear problems that can't be solved by classical methods. We have given two numerical examples and by comparing the numerical results obtain from ADM with the exact solution, we have studied their accuracy.

### Referances

- [1] G. Adomain, Stochastic systems analysis, in “Applied Stochastic Processes”, Academic Press, New York (1980) 1-18.
- [2] G. Adomain, Solving Frontier Problems of Physics: The Decomposition Method, Kluwer Academic Publishers, Dordrecht, 1994.
- [3] G. Adomain, A review of the decomposition method in applied mathematics, Journal of Mathematical Analysis and Applications 135 (1988) 501-544.
- [4] G. Adomain, Convergent series solution of nonlinear equations, Journal of Computational and Applied Mathematics 11 (1984) 225-230.
- [5] G. Adomain, On Green’s function in higher order stochastic differential equations, Journal of Mathematical Analysis and Applications 88 (1982) 604-606.
- [6] E. Babolian, H. Sadeghi, Sh. Javadi, Numerically solution of fuzzy differential equations by Adomain method, Applied Mathematics and Computation 149 (2004) 547-557.
- [7] M. Paripour, E. Hajilou, A. Hajilou, H. Heidari, Application of Adomain decomposition method to solve hybrid fuzzy differential equations, Journal of Taibah University for Science 9 (2015) 95-103.
- [8] L. Wang and S. Guo, Adomain method for second-order fuzzy differential equation, World Academy of Science, Engineering and Technology Vol: 5 (2011) 04-23.
- [9] L. Zadeh, Toward a generalized theory of uncertainty (GTU) – an outline, Information Sciences 175 (2005) 1-40.
- [10] O. Kaleva, Fuzzy differential equations, Fuzzy sets and Systems 24 (1987) 301-317.
- [11] B. Bede, I. J. Rudas, A. L. Bencsik, First order linear fuzzy differential equations under generalized differentiability, Information Sciences 177 (2007) 1648-1662.
- [12] Y. Chalco-Cano, H. Roman-Flores, Comparison between some approaches to solve fuzzy differential equations, Fuzzy Sets and Systems 160 (2009) 1517-1527.
- [13] Z. Ding, M. Ma, A. Kandel, Existence of the solutions of fuzzy differential equations with parameters, Information Sciences 99 (1997) 205-217.
- [14] S. Seikkala, On the fuzzy initial value problem, Fuzzy Sets and Systems 24 (1987) 319-330.
- [15] M. T. Mizukoshi, L. C. Barros, Y. Chalco-Cano, H. Roman-Flores, R. C. Bassanezi, Fuzzy differential equations and extension principle, Information Sciences 177 (2007) 3627-3635.
- [16] T. Allahviranloo, N. A. Kiani, N. Motamedi, Solving fuzzy differential equations by differential transformation method, Information Sciences 179 (2009) 956-966.
- [17] R. Rodriguez-Lopez, Monotone method for fuzzy differential equations, Fuzzy Sets and Systems 159 (2008) 2047-2076.
- [18] B. Ghazanfari, A. Shakerami, Numerical Solutions of fuzzy differential equations by extended Runge-Kutta-like formulae of order 4, Fuzzy Sets and Systems

189 (2011) 74-91.

- [19] C. Wu, S. Song, E. S. Lee, Approximate solutions, existence and uniqueness of the Cauchy problem of fuzzy differential equations, *Journal of Mathematical Analysis and Applications* 202 (1996) 629- 644.
- [20] J. J. Buckley, Thomas Feuring, Fuzzy differential equations, *Fuzzy sets and Systems* 110 (2000) 43-54.
- [21] C. Wu, S. Song, Existence theorem to the Cauchy problem of fuzzy differential equations under compactness- type conditions, *Information Sciences* 108 (1998) 123-134.
- [22] A. Khastan, R. Rodriguez-Lopez, On periodic solutions to first order linear fuzzy differential equations under differential inclusions' approach, *Information Sciences* 322 (2015) 31-50.
- [23] S. Song, C. Wu, Existence and uniqueness of Solutions to the Cauchy problem of fuzzy differential equations, *Fuzzy Sets and Systems* 110 (2000) 55-67.
- [24] M. Friedman, M. Ma, A. Kandel, Numerical solutions of fuzzy differential and integral equations, *Fuzzy Sets and Systems* 106 (1999) 35-48.
- [25] M. Mosleh, M. Otadi, Approximate solution of fuzzy differential equations under generalized differentiability, *Applied Mathematical Modelling* 39 (2015) 3003-3015.
- [26] V. M. Cabral, L. C. Barros, Fuzzy differential equation with completely correlated parameters. *Fuzzy Sets and Systems* 265 (2015) 86-98.
- [27] K. T. Atanassov, Intuitionistic fuzzy sets. VII ITKR's session, Sofia (deposited in Central Science and Technical Library of the Bulgarian Academy of Sciences 1697/84) (1983).
- [28] K. T. Atanassov, Intuitionistic fuzzy sets. *Fuzzy Sets Syst.* 20 (1986) 87–96.
- [29] K. T. Atanassov, Operators over interval valued intuitionistic fuzzy sets. *Fuzzy Sets Syst.* 64(2) (1994) 159–174.
- [30] M. Nikolova, N. Nikolov, C. Cornelis, G. Deschrijver, Survey of the research on intuitionistic fuzzy sets. *Adv. Stud. Contempor. Math.* 4(2) (2002) 127–157.
- [31] S. K. De, R. Biswas, , A. R. Roy, An application of intuitionistic fuzzy sets in medical diagnosis. *Fuzzy Sets Syst.* 117(2) (2001) 209–213.
- [32] M. H. Shu, C. H. Cheng, J. R. Chang, Using intuitionistic fuzzy sets for fault-tree analysis on printed circuit board assembly. *Microelectron. Reliab.* 46(12) (2006) 2139–2148.
- [33] D. F. Li, C. T. Cheng, New similarity measures of intuitionistic fuzzy sets and application to pattern recognitions. *Pattern Recognit. Lett.* 23 (2002) 221–225.
- [34] J. Ye, Multicriteria fuzzy decision-making method based on a novel accuracy function under interval valued intuitionistic fuzzy environment. *Expert Syst. Applicat.* 36 (2009) 6899–6902.
- [35] D. F. Li, Multiattribute decision making models and methods using intuitionistic fuzzy sets. *J. Comput. Syst. Sci.* 70 (2005) 73–85.
- [36] A. Kharal, Homeopathic drug selection using intuitionistic fuzzy sets. *Homeopathy* 98(1) (2009) 35–39.



- [37] M. Oberguggenberger, S. Pittschmann, Differential equations with fuzzy parameters. *Math. Mod. Syst.* 5 (1999) 181–202.
- [38] J. Casanovas, F. Rossell, Averaging fuzzy biopolymers. *Fuzzy Sets Syst.* 152 (2005) 139–158.
- [39] M .Z. Ahmad, B. De Baets, A predator–prey model with fuzzy initial populations. In: *Proceedings of the 13th IFSA World Congress and 6th European Society of Fuzzy Logic and Technology Conference, IFSA-EUSFLAT (2009)*.
- [40] L. C. Barros, R. C. Bassanezi, P. A. Tonelli, Fuzzy modelling in population dynamics. *Ecol. Model.* 128 (2000) 27–33.
- [41] M. S. El Naschie, From experimental quantum optics to quantum gravity via a fuzzy Khler manifold. *Chaos Solitons Fractals* 25 (2005) 969–977.
- [42] S. P. Mondal, T. K. Roy, First order linear non homogeneous ordinary differential equation in fuzzy environment. *Math. Theory Model.* 3(1) (2013) 85–95.
- [43] Z. Hassan, A.V. Kamyad, A. A. Heydari, Fuzzy modeling and control of hiv infection. *Comput. Math. Methods Med.* (2012). doi:10.1155/2012/893474
- [44] S. P. Mondal, S. Banerjee, T.K. Roy, First order linear homogeneous ordinary differential equation in fuzzy environment. *Int. J. Pure Appl. Sci. Technol.* 14(1) (2013) 16–26.
- [45] A. Bencsik, B. Bede, J. Tar, J. Fodor, Fuzzy differential equations in modeling hydraulic Differential servo cylinders. In: *Third Romanian-Hungarian joint symposium on applied computational intelligence (SACI), Timisoara, Romania (2006)*.
- [46] V. Nirmala and S. C. Pandian, Numerical Approach for Solving Intuitionistic Fuzzy Differential Equation under Generalised Differentiability Concept, *Applied Mathematical Sciences*, Vol. 9, 2015, no. 67, 3337 – 3346.
- [47] R. Ettoussi, S. Melliani, M. Elomari and L. S. Chadli, Solution of intuitionistic fuzzy differential equations by successive approximations method, *Notes IFS* 21 (2015) No. 2,51–62.
- [48] S. Melliani, M. Elomari, M. Atraoui and L. S. Chadli, Intuitionistic fuzzy differential equation with nonlocal condition, *Notes on Intuitionistic Fuzzy Sets* 21(2015) No. 4, 58–68.
- [49] Sankar Prasad Mondal and Tapan Kumar Roy, System of Differential Equation with Initial Value as Triangular Intuitionistic Fuzzy Number and its Application, *Int. J. Appl. Comput. Math* 1 (2015) 449– 474.

## Involution Matrices of $\frac{1}{4}$ -Quaternions

**Murat BEKAR**

Necmettin Erbakan University, Department of Mathematics-Computer  
Sciences, 42090, Konya, Turkey

mbekar@konya.edu.tr

**Abstract**

In this work, we consider the  $\frac{1}{4}$ -quaternion algebra and give the matrix representations of the involution and anti-involution maps obtained by this algebra.

**References**

- [1] Bekar M., and Yayli Y., Involution Matrices of Real Quaternions, Caspian J. of Math. Sci. Vol. 5, No. 1, 7-16, 2016.
- [2] Jafari M. , On The Properties of Quasi-Quaternion Algebra, Commun. Fac. Sci. Univ. Ank. Series A1 Vol. 63, No. 1, 1-10, 2014.
- [3] Rosenfeld B.A., Geometry of Lie Groups, Kluwer Academic Publishers, Dordrecht , 1997.
- [4] Ercan Z., Yuce S., On properties of the Dual Quaternions, European J. of Pure and Appl. Math., Vol. 4, No. 2, 142-146, 2011.

## Topological Full Groups of Cantor Minimal Systems

**Anıl OZDEMİR**

Tobb University of Economics and Technology, Department of  
Mathematics  
anilozdemir@etu.edu.tr

**Abstract**

In this work, we study topological full groups of Cantor minimal systems. In recent years, this subject has been very popular since it supplies a connection between dynamical systems and group theory. We will investigate the relationship between conjugation of dynamical systems and isomorphism of their topological full groups. Moreover, topological full groups provide the first examples of finitely generated, simple and amenable groups. We will survey the ideas behind the proofs of these facts.

**Keywords:** Cantor Space, Topological Full Group, Simple Grup, Amenable Group.

**References**

- [1] Matui Hiroki. Some remarks on topological full groups of Cantor minimal systems, *Internat.J. Math.*, (2006), 231-251.
- [2] S. Bezuglyi and K. Medynets. Full Groups, Flip Conjugacy, and Equivalence of Cantor Minimal Systems, *Colloquium Mathematicum*, (2008).
- [3] K. Juschenko, N. Monod. Cantor systems, piecewise translations and simple amenable, *Annals of Mathematics* (2012).

## Global Stability to Nonlinear Neutral Differential Equations of First Order

**Ramazan YAZGAN, Cemil TUNÇ**

Yüzüncü Yıl University, Faculty of Science, Department of Mathematics,  
Van, Turkey

ryazgan503@gmail.com, cemtunc@yahoo.com

### Abstract

In this paper, we study globally asymptotically stability of zero solution to a nonlinear neutral differential of first order. The technique of the proof involves the fixed point method. By this way, we extend and improve some recent works in the literature.

**Keywords:** Fixed point theorem, globally asymptotically stability, neutral differential equation, first order.

### References

- [1] R. P. Agarwal and S. R. Grace, Asymptotic Stability of Differential systems of neutral type, *Appl.Math Letter* 13 (2000), 9-15.
- [2] C. Tunc, Exponential stability to a neutral differential equation of first order with delay, *Ann. Differential Equations* 29 (2013), 253-256.
- [3] Y. N Raffoul, Stability in neutral nonlinear differential equations with functional delays using fixed-point theory. *Math Comput Model* 40 (2004), 691-700.
- [4] T. A. Burton, *Stability by fixed point theory for functional differential equation*. New York: Dover, 2006.

## A Study on the Cartesian Product of a Special Graphs

**Nihat AKGÜNEŞ, Ahmet Sinan ÇEVİK**

nakgunes@konya.edu.tr, sinan.cevik@selcuk.edu.tr

### Abstract

Our main aim in this presentation is to extend these studies over  $\Gamma(S_M)$  to the cartesian product. In here,  $\Gamma(S_M)$  is a graph of monogenic semigroup  $S_M = \{x, x^2, x^3, \dots, x^n\}$  with zero. In detail, we will investigate some important graph parameters for the cartesian product of any two (not necessarily different) graphs  $\Gamma(S_{M_1})$  and  $\Gamma(S_{M_2})$ .

**Keywords:** Monogenic semigroup graph, Graph product.

### Referances

- [1] N. Akgunes, K. C. Das, and A. S. Cevik, Some properties on the tensor product of graphs obtained by monogenic semigroups. *Applied Mathematics and Computation* 235, 2014, pp. 352-357.  
 [2] K. C. Das, N. Akgunes, A. S. Cevik, On a graph of monogenic semigroup, *Journal of Inequalities and Applications*. 2013:44, 2013, pp.1-13.

## Unit Dual Lorentzian Sphere and Tangent Bundle of Lorentzan Unit 2-Sphere

**Murat BEKAR, Yusuf YAYLI**

Necmettin Erbakan University, Department of Mathematics-Computer Sciences,42090 Konya, Turkey  
 Ankara University, Department of Mathematics, 06100 Ankara, Turkey  
 mbekar@konya.edu.tr, yayli@science.ankara.edu.tr

### Abstract

The purpose of this study is two-fold, firstly to recall some basic concepts and notions of unit dual Lorentzian sphere. Secondly, to define a one-to-one relationship between the unit dual Lorentzian sphere and tangent bundle of Lorentzian unit 2-sphere.

### Referances

- [1] Yayli, Y., Caliskan, A. and Ugurlu, H. H., The E. Study Maps of Circles on Dual Hyperbolic and Lorentzian Unit Spheres  $H^2$  and  $S^2$ , *Mathematical Proceedings of the Royal Irish Academy*, Vol. 102, No. A1, 37-47, 2002.

[2] Hathout, F., Bekar, M. and Yayli, Y., N-legendre and N-slant curves in the unit tangent bundle of surfaces, Kuwait Journal of Science, (Accepted on 2015-In Press).

## On the Inverse Problem for a Sturm-Liouville Equation with Discontinuous Coefficient

**Volkan ALA, Khanlar R. MAMEDOV**

Department of Mathematics, Mersin University, Mersin, Turkey  
volkanala@mersin.edu.tr, hanlar@mersin.edu.tr

### Abstract

In this study, it is investigated nonhomogenous boundary value problem. During the solution it is encountered Sturm-Liouville problem with piecewise continuous coefficients and that contained eigenvalue parameter. One transmission condition, which given by as relations between the right and left hand limit of the solution at the point of discontinuity are added to the boundary conditions. We examined some spectral properties of the problem. The numeric solutions of eigenvalues are obtained. According to the spectral datas the inverse problem are researched.

**Keywords:** Discontinuous Sturm-Liouville Problem, Inverse Problem, Transmission Condition.

### Referances

- [1] A. N. Tikhonov and A. A. Samarskii, Equations of Mathematical Physics [in Russian], Nauka, Moscow (1977).  
[2] A.M.Akhtyamov, On the uniqueness of the solution of an inverse spectral problem. Diferential Equations, v.39, No.8, 2003, pp.1061-1066.

## Calculation and Analysis of Electronic Parameters of Electroluminescent Device Cells Through I-V Based Modeling

**Ali Kemal HAVARE, Özge ÇOLAKOĞLU HAVARE**  
Toros University Electrical and Electronics Engineering Department,  
Mersin, Turkey

Department of Mathematics, Mersin University, Mersin, Turkey  
alikelmal.havere@toros.edu.tr, ozgecolakoglu@mersin.edu.tr

### Abstract

Light-emitting electrochemical cells (LECs) is one of the simplest kinds of electroluminescent devices. LEC is constituted to an organic single layer structure that was sandwiched between a cathode and an anode. In this study we calculated theoretically of the electronic parameter of LECs device through I-V based modeling. The LEC diode electronic parameters as the ideality factor  $n$  and barrier height  $\phi_b$  were obtained using a method developed by Cheung and confirmed by Werner. The net current of a LEC device is due to the thermionic emission and it can be expressed as

$$I = I_0 \exp\left(\frac{q(V - IR_s)}{nkT}\right)$$

where  $V$  is applied voltage and saturation current  $I_0$  is defined as

$$I_0 = AA^*T^2 \exp\left(-\frac{q\phi_b}{kT}\right).$$

**Keywords:** LECs device, Electronic parameters, I-V modeling.

### References

- [1] Cheung, S. and Cheung, N., *Extraction of Schottky diode parameters from forward current-voltage characteristic*, Applied Physics Letters, 49, 85, 1986.  
[2] Werner, J. H., *Schottky barrier and pn-junction I/V plots — Small signal evaluation*, Applied Physics A, 47, 3, 291–300, 1988.

## On Some Properties of Sum Spaces

**Mahmut KARAKUŞ**  
Department of Mathematics, YYU  
matfonks@gmail.com

### Abstract

A sum is a continuous linear functional  $s$  defined on a  $K$ -space  $\lambda \supset \phi$  (space of finitely non-zero sequences) such that,  $s(z) = \sum_k z_k$ ,  $\forall z = (z_k) \in \phi$ . A  $K$ -space

$\lambda$  is called a sum space if and only if  $\lambda \supset \phi$  and  $\lambda^f = \lambda^\lambda$ , where  $\lambda^f = \{(f(\delta^k)) : f \in \lambda'\}$  and  $\lambda^\lambda$  is the set of all sequences  $x$  such that  $xy \in \lambda$ ,  $\forall y \in \lambda$  [4,6]. Here  $\delta^k$  is the sequence whose  $k$ th component is 1 all the others are 0,  $xy = (x_k y_k)$  for  $x = (x_k), y = (y_k)$  and  $\lambda'$  is the space of continuous linear functionals on  $\lambda$ .

An FK space  $\lambda \supset \phi$  is generalized semiconservative FK space if  $\lambda^f \subset \lambda^{\lambda^2}$ , where  $\lambda^{\lambda^2} = \lambda^{\lambda\lambda} = (\lambda^\lambda)^\lambda$ .

In this work, we give some definitions and theorems related with sum spaces and generalized semiconservative FK spaces.

**Keywords:** FK spaces,  $\beta$ - dual,  $f$ - dual, Semiconservative FK spaces.

### References

- [1] J. Boos, T. Leiger, Dual pairs of sequence spaces. *Int. J. Math. Math. Sci.*, **28**(2001): 9-23.
- [2] J. Boos, T. Leiger, Dual pairs of sequence spaces II, *Proc. Estonian Acad. Sci. Phys. Math.*, **51**(2002): 3-17.
- [3] J. Boos, T. Leiger, Dual pairs of sequence spaces III, *J. Math. Anal. Appl.*, **324** (2006): 1213-1227.
- [4] J. Boos, Classical and Modern Methods in Summability, *Oxford University Press. New York, Oxford*, 2000.
- [5] D. J. H. Garling, The  $\beta$ - and  $\gamma$ -duality of sequence spaces, *Proc. Camb. Phil. Soc.*, **63** (Jan. 1967), 963-981.
- [6] W. H. Ruckle, An abstract concept of the sum of a numerical series, *Can. J. Math.*, 1970, **22**: 863-874.

## On $q^\lambda$ and $q_0^\lambda$ Invariant Sequence Spaces

**Mahmut KARAKUŞ**  
 Department of Mathematics, YYU  
 matfonks@gmail.com

### Abstract

Invariant sequence spaces are very helpful for investigations of the duality of sequence spaces. For instance, if the sequence space  $X$  satisfies the condition  $\ell_\infty.X = X$  then its  $\alpha$ -,  $\beta$ - and  $\gamma$ - duals are same [4]. Garling [1] investigated  $B$ - and  $B_0$ - invariant sequence spaces and Buntinas [2] introduced and investigated  $q$ - and  $q_0$ - invariant sequence spaces and recently, Grosse- Erdmann [3] studied on  $\ell_1$  invariant sequence spaces.

In this work, we define  $q^\lambda$  and  $q_0^\lambda$  invariant sequence spaces,  $X$  with  $q^\lambda.X = X$  and  $q_0^\lambda.X = X$ , respectively. and give some related theorems.

**Keywords:** K- spaces,  $\lambda$ -boundedness and  $\lambda$ -convergence of a sequence,  $\beta$ -,  $\gamma$ -,  $f$ - duality.

### References

- [1] D. J. H. Garling, *On topological sequence spaces*, Proc. Camb. Phil. Soc., 63 (1967), 997-1019.
- [2] M. Buntinas, *Convergent and bounded Cesro sections in FK-spaces*, Math. Zeitschr., 121 (1971), 191-200.
- [3] K.-G. Grosse-Erdmann, *On  $\ell_1$  invariant sequence spaces*, J. Math. Anal. Appl., **262**(2001), 112-132.
- [4] J. Boos, *Classical and Modern Methods in Summability*, Oxford University Press. New York, Oxford, 2000.

## Some New Results on a Graph of Monogenic Semigroup

**Nihat AKGÜNEŞ**

Department of Mathematics-Computer Sciences, Necmettin Erbakan  
University, Konya, Turkey  
nakgunes@konya.edu.tr

### Abstract

In [2], it has been defined a new graph  $\Gamma(S_M)$  on monogenic semigroups  $S_M$  (with zero) having elements  $\{0, x, x^2, x^3, \dots, x^n\}$ . Many researchers have been working on this area after that work, for example [1,3,4]. As a continues study of these studies, in this paper, it will be investigated define some new graph parameters (such as covering number, accessible number, Zagreb indices, ect.) for monogenic semigroup graph  $\Gamma(S_M)$ .

### References

- [1] N. Akgunes, K. C. Das, A. S. Cevik, Topological indices on a graph of monogenic semigroups, Topics in Chemical Graph Theory. Topics in Chemical Graph Theory, Mathematical Chemistry Monographs, University of Kragujevac and Faculty of Science Kragujevac, Kragujevac, No.16a, 2014, pp. 1-21.
- [2] K. C. Das, N. Akgunes, A. S. Cevik, On a graph of monogenic semigroup, Journal of Inequalities and Applications. 2013:44, 2013, pp.1-13.



[3] K. C. Das, Proof of conjectures on the distance signless Laplacian eigenvalues of graphs. *Linear Algebra and its Applications*, 467, 2015, pp. 100-115.

[4] K. C. Das, N. Akgunes, M. Togan, A. Yurttas, I. N. Cangul, A. S. Cevik, On the first Zagreb index and multiplicative Zagreb coindices of graphs, *Analele Stiintifice ale Universitatii Ovidius Constanta*, 24(1), 2016, pp. 153-176.

## Computational Solution of Katugampola Conformable Fractional Differential Equations Via RBF Collocation Method

**Fuat USTA**

Department of Mathematics, Duzce University, Konuralp Campus, 81620,  
Duzce, Turkey  
fuatusta@duzce.edu.tr

### Abstract

In conjunction with the development of fractional calculus, conformable derivatives and integrals has been widely used a number of scientific areas. In this talk, we provide a numerical scheme to solve Katugampola conformable fractional differential equations via radial basis function (RBF) collocation technique. In order to confirm our numerical scheme, we present some numerical experiments results.

### Referances

[1] U. Katugampola, A new fractional derivative with classical properties, *ArXiv:1410.6535v2*.

[2] E. J. Kansa, Multiquadrics a scattered data approximation scheme with applications to computational fluid-dynamics. i. surface approximations and partial derivative estimates, *Computers and Mathematics with Applications*, 19(8-9), 1990, 127-145.

[3] C. Franke, R. Schaback, Solving partial differential equations by collocation using radial basis functions, *Applied Mathematics and Computation*, 93, 1998, 73-82.

## Some Integral Inequalities Via Conformable Calculus

**Fuat USTA, Mehmet Zeki SARIKAYA**

Department of Mathematics, Duzce University, Konuralp Campus, 81620,  
Düzce, Turkey  
fuatusta@duzce.edu.tr, sarikayamz@gmail.com

### Abstract

The purpose of this talk is making generalization of Gronwall, Volterra and Pachpatte type inequalities for conformable differential equations. Then we provide some upper or lower bound for fractional derivatives and integrals with the help of Katugampola definition for conformable calculus. These results are extensions of some existing Gronwall, Volterra and Pachpatte type inequalities in the previous studies.

### References

- [1] U. Katugampola, A new fractional derivative with classical properties, ArXiv:1410.6535v2.
- [2] T. Abdeljawad, On conformable fractional calculus, Journal of Computational and Applied Mathematics 279 (2015) 57{66.
- [3] M.Z. Sarikaya, Gronwall type inequality for conformable fractional integrals, 2016, preprint.

## Weighted Ostrowski, Chebyshev and Grüss Type Inequalities for Conformable Fractional Integrals

**Hüseyin BUDAK, Fuat USTA, Mehmet Zeki SARIKAYA**  
Department of Mathematics, Duzce University, Konuralp Campus, 81620,  
Düzce, Turkey

hsyn.budak@gmail.com, fuatusta@duzce.edu.tr, sarikayamz@gmail.com

### Abstract

In this presentation, we have obtained weighted versions of Ostrowski,  $\tilde{C}$ ebysev and Grüss type inequalities for conformable fractional integrals. In accordance with this purpose we have used Katugompola type conformable fractional integrals. The present study confirms previous findings and contributes additional evidence that provide the bounds for more general functions.

### Referances

- [1] U. Katugampola, A new fractional derivative with classical properties, ArXiv:1410.6535v2.
- [2] D. R. Anderson, Taylor's formula and integral inequalities for conformable fractional derivatives, Contributions in Mathematics and Engineering, in Honor of Constantin Caratheodory, Springer, to appear.
- [3] A. M. Ostrowski, Über die absolutabweichung einer di erentiebaren funktion von ihrem integralmittelwert, Comment. Math. Helv. 10(1938), 226-227..e 1 No. 1 pp. 48-53, 2013.
- [4] P. L. Ceby sev, Sur less expressions approximatives des integrales de nies par les autres prisesentre les memes limites, Proc. Math. Soc. Charkov, 2, 93-98, 1882.
- [5] G. Gruss, Über das maximum des absoluten Betrages von  $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx$ , Mat.Z., 39, 215-226, 1935.

## Bifurcation and Stability Analysis of a Discrete-Time Predator-Prey Model

Gökçe SUCU  
sucugokce@gmail.com

### Abstract

We consider discrete-time Leslie Model. We first determine its non-negative fixed point. Later on, we study local stability of the fixed point and determine the conditions on the parameters to show the existence of flip bifurcation by taking the step-size as a bifurcation parameter. Analytical results are also supported by some numerical simulation. Moreover, using Center Manifold Theory, we show the existence of flip bifurcation and its properties.

## Applications of Hermite-Hadamard Inequalities for $\mathbb{B}$ -Convex Functions and $\mathbb{B}^{-1}$ -Convex Functions

**Gabil ADILOV, İlknur YEŞİLCE**  
Akdeniz University, Antalya, Turkey  
Mersin University, Mersin, Turkey  
gabiladilov@gmail.com, ilknuryesilce@gmail.com

### Abstract

Hermite-Hadamard Inequality that is expressed in the following form

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(t) dt \leq \frac{1}{2}(f(a) + f(b))$$

was proven by Hermite in [1] and then, ten years later, Hadamard rediscovered in [2] (for the historical consideration see also [3]). Next, Hermite-Hadamard Inequalities for different kinds of functions were examined in numerous article [4,5].

$\mathbb{B}$ -convex functions were introduced and studied in [6].  $\mathbb{B}^{-1}$ -convex functions were defined and examined in [7]. Hermite-Hadamard Inequalities for  $\mathbb{B}$ -convex Functions and  $\mathbb{B}^{-1}$ -convex Functions were introduced in [8].

In this work, we give the applications of Hermite-Hadamard Inequalities for  $\mathbb{B}$ -convex functions and  $\mathbb{B}^{-1}$ -convex functions.

**Keywords:** Hermite-Hadamard Inequalities,  $\mathbb{B}$ -convex functions,  $\mathbb{B}^{-1}$ -convex functions.

### References

- [1] Ch. Hermite (1883). Sur deux limites d'une integrale define, *Mathesis*. 3, 82.
- [2] J. Hadamard (1893). Etude sur les proprietes des fonctions entieres et en particulier d'une fonction consideree par Riemann, *Journal des Mathematiques Pures et Appliquees*, 58, 171-215.
- [3] S.S. Dragomir, C.E.M. Pearce (2000). *Selected Topics on Hermite-Hadamard Inequalities and Applications*, RGMIA Monographs, Victoria University.
- [4] H. Kavurmaci, M. Avci, M.E. Ozdemir (2011). New Inequalities of Hermite-Hadamard Type for Convex Functions with Applications, *Journal of Inequalities and Applications*, 2011:86, doi: 10.1186/1029-2011-86.

- [5] M.E. Ozdemir, E. Set, M.Z. Sarikaya (2011). Some New Hadamard Type Inequalities for Co-Ordinated  $m$ -Convex and  $(\alpha, m)$ -Convex Functions, *Hacettepe Journal of Mathematics and Statistics*, 40, no. 2, 219-229.
- [6] Kemali, S., Yesilce, I., Adilov, G.,  $\mathbb{B}$ -convexity,  $\mathbb{B}^{-1}$ -convexity and Their Comparison, *Numerical Functional Analysis and Optimization*, Vol. 36, No. 2, pp. 133-146, 2015.
- [7] G. Adilov and I. Yesilce (2017).  $\mathbb{B}^{-1}$ -convex Functions. *Journal of Convex Analysis*. 24(2).
- [8] Adilov, G., Yesilce, I., The Hermite-Hadamard Inequalities for  $\mathbb{B}$ -convex and  $\mathbb{B}^{-1}$ -convex Functions, (submitted).

## Operations and Extension Principle under T-Intuitionistic Fuzzy Environment

**Arindam GARAI, Palash MANDAL, Tapan Kumar ROY**

\*Department of Mathematics, Sonarpur Mahavidyalaya, West Bengal,  
India, Pin 700149.

Department of Mathematics, IEST, Shibpur, Howrah, West Bengal,  
India, Pin 711103

fuzzy\_arindam@yahoo.com, palashmandalmbss@gmail.com,  
roy\_t.k@yahoo.co.in

### Abstract

In this paper, we introduce necessary definitions and related theorems on intuitionistic fuzzy T-set theory. In existing crisp set theory, characteristic functions, defined by two valued logic, can take values: zero and one only. In fuzzy set theory, introduced by Zadeh (1965), membership functions can take any value in closed unit interval . And, in intuitionistic fuzzy set theory, introduced by K. T. Atanassov (1986), both membership functions and non-membership functions can take suitable values in closed unit interval . But, we may observe that in those existing theories, we have to assign same membership value unity to elements, even if belongingness of one element is more certain than other to the subsets. In other words, certainly belongingness and more certainly belongingness of elements to subsets of universal sets are treated at par in all these existing theories, including classical crisp set theory. Similarly, in non-membership functions of intuitionistic fuzzy sets, zero is assigned as non-membership values to elements, both not belonging and not belonging certainly to subsets of universal sets. In order to overcome these limitations of existing theories, in 2015, we proposed intuitionistic fuzzy T-set theory, in which real numbers are suitably assigned to membership and non-membership functions. In this paper, we further introduce necessary definitions and related theorems

on intuitionistic fuzzy  $T$ -set theory. Those may be considered as generalizations of existing definitions and theorems from existing fuzzy and/or intuitionistic fuzzy set theory. In particular, we have generalized the concepts of extension principle and alpha, beta cut under existing intuitionistic fuzzy environment to  $T$ -intuitionistic fuzzy environment. Moreover, we have discussed some associated results under  $T$ -intuitionistic fuzzy environment. Finally, conclusions and future research directions are drawn.

**Keywords:** Intuitionistic fuzzy sets,  $T^{(+)}$ -characteristic functions,  $T^{(-)}$ -characteristic functions, intuitionistic Fuzzy  $T$ -sets,  $T$ -extension principle.

## On Generalized Double Statistical Convergence of Order $\alpha$ in Intuitionistic Fuzzy $N$ - Normed Spaces

**Ekrem SAVAŞ**

Istanbul Ticaret University, Department of Mathematics, Sötlüce,  
Istanbul, Turkey  
ekremsavas@yahoo.com, esavas@iticu.edu

### Abstract

In the present paper, we introduce and study the notion  $\mathcal{I}$ -double statistical convergence and ideal  $\lambda$ -double statistical convergence of order  $\alpha$  with respect to the intuitionistic fuzzy  $n$ -normed space, briefly  $IFnNS$ , also we examine the relationship between these classes.

### Referances

- [1] Al. Narayanan, S. Vijayabalaji, Fuzzy  $n$ -normed linear space, Int. J. Math. Sci., 24, 3963-3977, 2005.
- [2] Al. Narayanan, S. Vijayabalaji, N. Thillaigovindan, Intuitionistic fuzzy bounded linear operators, Iran. J. Fuzzy Systems, 4, 89-101, 2007.
- [3] N. Thillaigovindan, S. A. Shanthi, Y. B. Jun, On lacunary statistical convergence in intuitionistic fuzzy  $n$ -normed linear space, Annals of Fuzzy Math. and Inf., 1(2), 119-131, 2011.
- [4] E. Savas,  $\lambda$ -statistical convergence in intuitionistic fuzzy 2-normed space, Appl. Math. Inf. Sci., 9(1), 501-505, 2015

## On Feng Qi-Type Integral Inequalities for Conformable Fractional Integrals

Abdullah AKKURT, M. Esra YILDIRIM\*,  
Hüseyin YILDIRIM

Department of Mathematics, Faculty of Science and Arts, University of  
Kahramanmaraş Sütçü İmam, 46100, Kahramanmaraş, Turkey

\*Department of Mathematics, Faculty of Science, University of  
Cumhuriyet, 58140, Sivas, Turkey

abdullahmat@gmail.com, mesra@cumhuriyet.edu.tr, hyildir@ksu.edu.tr

### Abstract

In the last few decades, much significant development of integral inequalities had been established. Recall the famous integral inequality of Feng Qi type [1, 2, 3]:

$$\int_a^b (f(t))^{n+2} dt \geq \left( \int_a^b f(t) dt \right)^{n+1} \quad (0.0.8)$$

where  $f \in C^n(a, b)$ ,  $f^{(i)} \geq 0$ ,  $0 \leq i \leq n$ ,  $f^{(n)} \geq n!$ ,  $n \in \mathbb{R}$ .

In this study, we establish the generalized Qi-type inequality involving conformable fractional integrals.

Firstly we give a important integral inequality which is generalized Qi inequality. Finally, we obtain several inequalities related these inequalities using the conformable fractional integral [4,5].

**Keywords:** Integral Inequalities, Special Functions, Fractional Calculus, Conformable Fractional Integral.

### Referances

- [1] F. Qi, Several integral inequalities, J. Inequal. Pure and Appl. Math. 1 (2000) Art. 19.
- [2] A. Akkurt and H. Yildirim, Genelleştirilmiş Fractional Integraller İçin Feng Qi Tipli Integral Eşitsizlikleri Üzerine, Fen Bilimleri Dergisi, 1(2), (2014).
- [3] Q.A. Ngô, D.D. Thang, T.T. Dat and D.A. Tuan, Note on an integral inequality, J. Inequal. Pure Appl. Math., 7(4) (2006), Art. 120.
- [4] T. Abdeljawad, On conformable fractional calculus, Journal of Computational and Applied Mathematics 279 (2015) 57–66.
- [5] R. Khalil, M. Al horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, Journal of Computational Applied Mathematics, 264 (2014), 65-70.